

## Cytomegalovirus Antigenemia: Clinical Correlations in Transplant Recipients and in Persons with AIDS

TONY MAZZULLI,<sup>1†</sup> ROBERT H. RUBIN,<sup>1</sup> MARY JANE FERRARO,<sup>2</sup> RICHARD T. D'AQUILA,<sup>1</sup>  
SANDRA A. DOVEIKIS,<sup>2</sup> BENJAMIN R. SMITH,<sup>1</sup> T. HAUW THE,<sup>3</sup> AND MARTIN S. HIRSCH<sup>1\*</sup>

*Infectious Disease Unit<sup>1</sup> and Division of Microbiology,<sup>2</sup> Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts 02114, and Department of Clinical Immunology, University Hospital, Groningen, The Netherlands<sup>3</sup>*

Received 25 March 1993/Returned for modification 21 May 1993/Accepted 20 July 1993

**We evaluated a rapid immunoperoxidase technique for the detection of cytomegalovirus (CMV) antigenemia in peripheral blood neutrophils of 56 transplant recipients (117 specimens) and 36 persons with AIDS (59 specimens). Antigenemia was 92% sensitive and 98% specific for the detection of clinical CMV infection in transplant recipients and 100% sensitive and 86% specific in persons with AIDS. Overall, CMV antigenemia was a more rapid and sensitive method for the detection of clinical CMV infection than either shell vial culture or conventional tube culture of blood.**

Infection with human cytomegalovirus (CMV) continues to be a major cause of morbidity and mortality in immunocompromised patients. Twenty to 60% of recipients of all types of allografts and 25 to 90% of patients with AIDS may develop active CMV infection (2, 15). With the introduction of effective anti-CMV therapies, early and rapid diagnosis is essential for proper management, because the benefit of therapy is substantially increased by early administration (7).

The detection of CMV in blood specimens (CMV viremia) has been shown to be the most specific indicator of significant CMV infection (10, 16). However, conventional tube cultures (CCs) of blood may not yield results for several weeks. Although shell vial culture (SV) techniques have shortened the detection time for CMV infection considerably (5, 6), they may be negative in 37% or more of blood specimens that are positive by CC (11).

Recently, a diagnostic technique was developed that allows for the rapid and quantitative detection of CMV antigenemia in peripheral blood polymorphonuclear leukocytes (PMNLs) (1, 3, 13, 18, 19, 21-23). This technique uses a pool of monoclonal antibodies directed against what was initially thought to be CMV immediate early antigen but is now recognized as being a lower-matrix phosphoprotein (pp65) (8, 12). Using these monoclonal antibodies, we evaluated an immunoperoxidase staining technique for the diagnosis of CMV antigenemia in transplant recipients and human immunodeficiency virus-infected patients, comparing the results with those from SV and CC of blood. The results were correlated with the presence or absence of clinical CMV disease.

Between December 1991 and December 1992, blood specimens from all organ transplant recipients with unexplained fever 3 weeks to 3 months posttransplantation and persons with AIDS with CD4<sup>+</sup> lymphocyte counts of <100 cells per mm<sup>3</sup> were examined for the presence of CMV antigenemia and viremia. Interpretation of the antigenemia assay slides was made independent of the results of SV, CC, and the

clinical status of the patients. Transplant recipients were deemed to have clinical CMV infection if they had no other cause of their fever identified, had evidence of CMV by conventional culture or tissue biopsy, and responded to ganciclovir therapy. Persons with AIDS were deemed to have clinical CMV infection if they had signs or symptoms (e.g., retinitis) suggestive of CMV disease and conventional culture or tissue biopsy evidence of CMV. Among the transplant recipients, patients receiving ganciclovir therapy for  $\geq 24$  h at the time the blood specimen was drawn were excluded from the study.

Preparation and staining of PMNL cytopspins were carried out in accordance with methods described by van der Bij et al. (22), with slight modification. Cytopspin preparations were made with 100  $\mu$ l of a suspension of  $2 \times 10^6$  cells per ml (Shandon cytopspin III; Shandon/Lipshaw, Pittsburgh, Pa.), and only monoclonal antibodies C10 and C11 (22) were used. Specimens known to contain CMV antigen-positive cells were used as positive controls. Negative control PMNL slides were prepared from healthy donors.

For SV and CC, 5 ml of EDTA-treated blood was allowed to sediment by gravity and the leukocyte-rich plasma was removed and centrifuged. The pellet was washed twice with minimal essential media (Bio Whittaker, Walkersville, Md.) and then was resuspended in 2 ml of minimal essential medium with 10% fetal bovine serum. CCs with human embryonic lung fibroblast (HEL 299) cells (American Type Culture Collection) were inoculated with 1 ml of the cell suspension. Three SVs, prepared weekly in-house, with MRC-5 cells (Viomed Laboratories, Minneapolis, Minn.) were inoculated with 0.3 ml of the cell suspension and centrifuged at  $700 \times g$  for 40 min at 20 to 30°C. CCs were maintained for 4 weeks (or longer if there was evidence of CMV-induced cytopathic effect). Coverslips from two SVs were fixed, stained (with fluorescein isothiocyanate-labelled CMV monoclonal antibody [Syva MicroTrak, Palo Alto, Calif.]), and examined at 16 to 20 h, and the third SV was examined at 48 h. Any specimen that was SV negative but showed the characteristic cytopathic effect in CC was confirmed with the fluorescein isothiocyanate-labelled monoclonal antibody.

One hundred seventy-six specimens from 56 transplant recipients (117 specimens) and 36 persons with AIDS (59

\* Corresponding author.

† Present address: Department of Microbiology, Mount Sinai Hospital, Toronto, Ontario M5G 1X5, Canada.

TABLE 1. Comparison of antigenemia assay, SV, and CC of blood for detection of clinical CMV infection<sup>a</sup>

Assay	Transplant recipients				Persons with AIDS			
	Clinical CMV infection (no. positive/no. tested)	No clinical CMV infection (no. positive/no. tested)	Sensitivity (%)	Specificity (%)	Clinical CMV infection (no. positive/no. tested)	No clinical CMV infection (no. positive/no. tested)	Sensitivity (%)	Specificity (%)
Antigenemia	45/49	1/68	92	98	23/23	5/36	100	86
SV	9/40	0/57	23	100	4/10	0/10	40	100
CC	24/41	0/62	59	100	14/21	2/34	67	94

<sup>a</sup> Values are for specimens tested.

specimens) were evaluated for antigenemia. For 49 specimens, SV was not performed, and for 18 of these specimens CC was not performed. For an additional 10 SV specimens, the inoculum was toxic for the fibroblast monolayer, making them uninterpretable. Of the transplant recipients, there were 19 liver, 19 kidney, 8 lung, and 10 heart allografts. Thirty-five percent of transplant recipients and 30% of the persons with AIDS were women.

Table 1 shows that in both transplant recipients and persons with AIDS, the antigenemia assay was highly sensitive (92 to 100%) and specific (86 to 98%) for the detection of clinical CMV infection. The sensitivities of SVs and CCs were very low (23 to 40% and 59 to 67%, respectively). CC was positive a mean of 24 days (median, 20 days; range, 7 to 57 days) after the specimen was drawn.

The sensitivities of the antigenemia assay and SV for the detection of CMV viremia by CC were 92 and 41%, respectively, for transplant recipients and 100 and 44%, respectively, for patients with AIDS. The specificities were 78 and 100%, respectively, for transplant recipients and 74 and 100%, respectively, for patients with AIDS (Table 2).

The mean number of antigen-positive cells per slide in the six specimens from patients without clinical CMV infection was 2.1 cells (range, 1 to 4; median, 2), compared with a mean of 76 antigen-positive cells in the 68 specimens from patients with clinical CMV infection (range, 1 to 600; median, 14) ( $P < 0.005$ , Mann-Whitney U test). The transplant recipient in whom the antigenemia assay was positive, but who had no evidence of clinical CMV infection, was CMV seropositive and had received a liver from a CMV-seropositive donor 30 days prior to the antigenemia test. Of the five patients with AIDS with a positive antigenemia test but no evidence of clinical CMV infection at the time of the test,

two patients had a blood culture positive for CMV, two other patients had positive blood cultures on a subsequent specimen, and the final patient developed CMV colitis within 2 months of the positive antigenemia result.

Of the four transplant recipients with negative antigenemia but evidence of clinical CMV infection, all had negative SVs, two had positive CCs (on days 10 and 57 of culture), and three had positive antigenemia detected on a blood specimen drawn within 10 days of the first negative assay. Three of these patients had fever and leukopenia, and the fourth had fever and elevated liver function tests. Forty-five specimens from 26 transplant recipients with 36 episodes of clinical CMV infection were antigenemia positive. Of these 26 patients, 15 (21 episodes) had fever and leukopenia (28 antigen-positive specimens, 14 positive CCs of blood for CMV, 8 positive urine and/or throat cultures for CMV, 2 biopsies of the gastric antrum positive for CMV), 7 (11 episodes) had fever and elevated liver function tests with or without leukopenia (13 antigen-positive specimens, 4 positive CCs of blood for CMV, 6 histologic and/or culture-positive liver biopsies for CMV, 6 positive urine and/or throat cultures), and 4 (4 episodes) had fever and pneumonitis (4 antigen-positive specimens, 4 positive CCs of blood for CMV, 2 bronchoalveolar lavage specimens positive for CMV).

Antigenemia was positive in all 23 specimens from 13 persons with AIDS who had clinical CMV infection. Of these 13 persons, 9 had CMV retinitis (15 antigen-positive specimens, 9 positive CCs of blood for CMV), 2 had CMV colitis (4 antigen-positive specimens, 1 positive CC of blood for CMV), 1 had CMV pneumonitis (1 antigen-positive specimen, 1 positive blood culture, and 1 positive bronchoalveolar lavage culture for CMV), and 1 had unexplained fever (3 antigen-positive specimens, 3 positive CCs of blood for CMV). In all cases of CMV retinitis, the diagnosis was made by an ophthalmologist. The diagnosis of CMV colitis was based on endoscopic findings and histologic examination of biopsy specimens.

In nine transplant recipients (11 episodes of clinical CMV infection) from whom a blood specimen was examined during or immediately after ganciclovir therapy, antigenemia levels fell rapidly and corresponded to clinical improvement. In two persons with AIDS and CMV retinitis, a rapid rise in antigenemia level while receiving ganciclovir therapy paralleled the development of progressive CMV disease.

The results of our study show that an antigenemia assay for the detection of CMV in PMNLs is a sensitive and rapid test for the diagnosis of CMV infection and disease. The antigenemia assay provided positive results in 27 specimens with negative SV and CC results. Clinical CMV infection was present in all but four of these specimen donors at the time. Others have also reported greater sensitivity of the

TABLE 2. Comparison of antigenemia assay and SV with CC for detection of positive CMV blood culture

Assay result <sup>a</sup>		CC			
		No. of transplant recipients:		No. of persons with AIDS:	
		Positive	Negative	Positive	Negative
Antigenemia	+	22	17	16	10
	-	2	62	0	29
SV	+	9	0	4	0
	-	13	75	5	11

<sup>a</sup> The sensitivity and specificity of the antigenemia assay for the detection of CMV viremia were 92 and 78%, respectively, in transplant recipients and 100 and 74%, respectively, in persons with AIDS. The sensitivity and specificity of the SV for the detection of CMV viremia were 41 and 100%, respectively, in transplant recipients and 44 and 100%, respectively, in persons with AIDS.

antigenemia assay compared with both SV and CC techniques (4, 14). Only one study thus far has reported poor results with an antigenemia assay, detecting no CMV-positive transplant recipients (9). The poor results in this study may have been due to less frequent testing than in other studies, delays in processing specimens, differences in host immunocompetence, and the use of different monoclonal antibodies (9).

Patients in our study who were antigenemia positive and had evidence of clinical CMV infection had a mean number of antigen-positive cells per slide that was significantly higher than those patients who had no evidence of clinical CMV infection. Others have also reported that the presence or absence of CMV disease is closely related to the number of antigen-positive cells (18, 20). However, as in our study, some patients with clinical CMV infection may have low levels of antigenemia, suggesting that host factors have an important role in determining the clinical impact of a given level of antigenemia.

The quantitative nature of the antigenemia assay may give an estimate of viral load, and this may be useful for monitoring patients before, during, and after therapy (18, 19). Serial testing with the antigenemia assay of patients at risk for CMV infection and disease may allow the detection of important changes in the antigenemia level. Early positive or rising antigenemia levels may signal the onset of active CMV disease and allow early preemptive therapy to be initiated, particularly in transplant recipients (18, 19). In our study, transplant recipients with clinical CMV infection treated with ganciclovir showed a rapid decline in antigenemia levels which paralleled clinical improvement. A persistently high or rising level of antigenemia despite appropriate CMV therapy may signal progressive CMV disease or the development of viral resistance. Further study, however, is needed to determine the significance of persistently elevated or rising antigenemia levels during or immediately after therapy (4).

The CMV antigenemia assay is relatively simple to perform and may be completed in 5 to 6 h (22). Interpretation of slides, however, may be difficult if there is a great deal of background endogenous peroxidase staining (usually the result of eosinophils in the preparation) or if the morphology of the PMNLs is distorted (17). Despite these potential problems, the CMV antigenemia assay appears to give accurate results, making it useful for the rapid detection of CMV disease in blood specimens from patients at risk. Because the antigenemia assay may occasionally miss some cases of clinical CMV infection, repeat testing should be considered for patients for whom there is strong clinical suspicion of active CMV infection. As was noted in our study, the antigenemia assay was positive on a repeat specimen drawn within 10 days of the initial negative result in three of four patients with clinical CMV infection. The lower level of sensitivity and lengthy time (an average of 24 days in our study) before CC of blood for CMV becomes positive makes it clinically less useful than the antigenemia assay in managing patients at risk for CMV disease.

Tony Mazzulli is supported by a Medical Research Council of Canada Fellowship. This work was also supported by grants NIH CA35020 and CA54741.

We thank Mary Ellen Halliwell and Lucy Chou for technical support and Teri Flynn, Meg White-Guthro, Mary Sullivan, Maureen Doran, Angela Delvecchio, Nesli Basgoz, Jay Fishman, Steve Boswell, and Howard Heller for providing clinical specimens and information on patients.

## REFERENCES

- Drew, W. L., V. I. Cafaro, R. C. Minor, and D. Jekic-McMullen. 1992. Detection of cytomegalovirus (CMV) antigenemia using the CMV-vue antigenemic assay, abstr. 1427. Program Abstr. 32nd Intersci. Conf. Antimicrob. Agents Chemother.
- Dworkin, R., and W. L. Drew. 1989. Therapy for herpesvirus infections in AIDS. *Curr. Opin. Infect. Dis.* 3:108-112.
- Erice, A., M. A. Holm, P. C. Gill, S. Henry, C. L. Dirksen, D. L. Dunn, R. P. Hillam, and H. H. Balfour, Jr. 1992. Cytomegalovirus (CMV) antigenemia assay is more sensitive than shell vial cultures for rapid detection of CMV in polymorphonuclear blood leukocytes. *J. Clin. Microbiol.* 30:2822-2825.
- Gerna, G., D. Zipeto, M. Parea, M. G. Revello, E. Silini, E. Percivalle, M. Zavattoni, P. Grossi, and G. Milanese. 1991. Monitoring of human cytomegalovirus infections and ganciclovir treatment in heart transplant recipients by determination of viremia, antigenemia, and DNAemia. *J. Infect. Dis.* 164:488-498.
- Gleaves, C. A., C. F. Lee, L. Kirsch, and J. D. Meyers. 1987. Evaluation of a direct fluorescein-conjugated monoclonal antibody for detection of cytomegalovirus in centrifugation culture. *J. Clin. Microbiol.* 25:1548-1550.
- Gleaves, C. A., T. F. Smith, E. A. Shuster, and G. R. Pearson. 1985. Comparison of standard tube and shell vial cell culture techniques for the detection of cytomegalovirus in clinical specimens. *J. Clin. Microbiol.* 21:217-221.
- Goodrich, J. M., M. Mori, C. A. Gleaves, C. Du Mond, M. Cays, D. F. Ebeling, W. C. Buhles, B. DeArmond, and J. D. Meyers. 1991. Early treatment with ganciclovir to prevent cytomegalovirus disease after allogeneic bone marrow transplantation. *N. Engl. J. Med.* 325:1601-1607.
- Grefte, J. M. M., B. T. F. van der Gun, S. Schmolke, M. van der Giessen, W. J. van Son, B. Plachter, G. Jahn, and T. H. The. 1991. The lower matrix protein pp65 is the principal viral antigen present in peripheral blood leukocytes during an active cytomegalovirus infection. *J. Gen. Virol.* 73:2923-2932.
- Miller, H., E. Rossier, R. Milk, and C. Thomas. 1991. Prospective study of cytomegalovirus antigenemia in allograft recipients. *J. Clin. Microbiol.* 29:1054-1055.
- Myers, J. D., P. Ljungman, and L. D. Fisher. 1990. Cytomegalovirus excretion as a predictor of cytomegalovirus disease after marrow transplantation: importance of cytomegalovirus viremia. *J. Infect. Dis.* 162:373-380.
- Paya, C. V., A. D. Wold, and T. F. Smith. 1987. Detection of cytomegalovirus infections in specimens other than urine by the shell vial assay and conventional tube cell cultures. *J. Clin. Microbiol.* 25:755-757.
- Revello, M. G., E. Percivalle, A. Di Matteo, F. Morini, and G. Gerna. 1992. Nuclear expression of the lower matrix protein of human cytomegalovirus in peripheral blood leukocytes of immunocompromised viremic patients. *J. Gen. Virol.* 73:437-442.
- Revello, M. G., E. Percivalle, M. Zavattoni, M. Parea, P. Grossi, and G. Gerna. 1989. Detection of human cytomegalovirus immediate early antigen in leukocytes as a marker of viremia in immunocompromised patients. *J. Med. Virol.* 29:88-93.
- Revello, M. G., M. Zavattoni, E. Percivalle, P. Grossi, and G. Gerna. 1989. Correlation between immunofluorescent detection of human cytomegalovirus immediate early antigens in polymorphonuclear leukocytes and viremia. *J. Infect. Dis.* 160:159-160.
- Rubin, R. H. 1988. Infections in the patient after renal and liver transplantation, p. 557-583. *In* R. H. Rubin and L. S. Young (ed.), *Clinical approach to infections in the immunocompromised patient*, 2nd ed. Plenum Press, New York.
- Salmon, D., F. Lacassin, M. Harzic, C. Lepout, C. Perronne, F. Bricaire, F. Brun-Vezinet, and J. L. Vilde. 1990. Predictive value of cytomegalovirus viraemia for the occurrence of CMV organ involvement in AIDS. *J. Med. Virol.* 32:160-163.
- The, T. H., W. van der Bij, A. P. van den Berg, M. van der Giessen, J. Weits, H. G. Sprenger, and W. J. van Son. 1990. Cytomegalovirus antigenemia. *Rev. Infect. Dis.* 12(Suppl. 7): S737-S744.
- Van den Berg, A. P., I. J. Klompaker, E. B. Haagsma, A. Scholten-Sampson, C. M. A. Bijleveld, J. Schirm, M. van der

- Giessen, M. J. H. Slooff, and T. H. The. 1991. Antigenemia in the diagnosis and monitoring of active cytomegalovirus infection after liver transplantation. *J. Infect. Dis.* **164**:265–270.
19. Van den Berg, A. P., W. van der Bij, W. J. van Son, J. Anema, M. van der Giessen, J. Schirm, A. M. Tegzess, and T. H. The. 1989. Cytomegalovirus antigenemia as a useful marker of symptomatic cytomegalovirus infection after renal transplantation—a report of 130 consecutive patients. *Transplantation* **48**:991–995.
  20. Van den Berg, A. P., W. J. van Son, N. M. Jiwa, W. van der Bij, J. Schirm, M. van der Giessen, and T. H. The. 1990. Recent advances in the diagnosis of active CMV infection after organ transplantation. *Transplant. Proc.* **22**:226–228.
  21. van der Bij, W., J. Schirm, R. Torensma, W. J. van Son, A. M. Tegzess, and T. H. The. 1988. Comparison between viremia and antigenemia for detection of cytomegalovirus in blood. *J. Clin. Microbiol.* **26**:2531–2535.
  22. van der Bij, W., R. Torensma, W. J. van Son, J. Anema, J. Schirm, A. M. Tegzess, and T. H. The. 1988. Rapid immunodiagnosis of active cytomegalovirus infection by monoclonal antibody staining of blood leucocytes. *J. Med. Virol.* **25**:179–188.
  23. van der Bij, W., W. J. van Son, A. P. M. van der Berg, A. M. Tegzess, R. Torensma, and T. H. The. 1989. Cytomegalovirus (CMV) antigenemia: rapid diagnosis and relationship with CMV-associated clinical syndromes in renal allograft recipients. *Transplant. Proc.* **21**:2061–2064.