

## *Staphylococcus lugdunensis*: Report of a Case of Peritonitis and an Easy-To-Perform Screening Strategy

NORBERT SCHNITZLER,<sup>1</sup> RAINER MEILICKE,<sup>1</sup> GEORG CONRADS,<sup>1</sup> DARIUS FRANK,<sup>2</sup>  
AND GERHARD HAASE<sup>1\*</sup>

*Institute of Medical Microbiology and National Reference Center for Streptococci,<sup>1</sup> and Department of Internal Medicine II,<sup>2</sup> University Hospital RWTH Aachen, D-52057 Aachen, Germany*

Received 3 September 1997/Returned for modification 17 November 1997/Accepted 8 December 1997

**We report on a severe case of peritonitis due to *Staphylococcus lugdunensis*. The clinical course resembled an infection due to *S. aureus* more than one due to other coagulase-negative staphylococci. Therefore, we strongly recommend identification and propose an easy-to-perform procedure for screening of this pathogen.**

Staphylococcal peritonitis is an important cause of morbidity in continuous ambulatory peritoneal dialysis (CAPD) patients (8). While coagulase-negative staphylococci (CNS) are the most common cause of peritonitis in CAPD patients, *Staphylococcus aureus* is the leading cause of more serious infections, often necessitating removal of the catheter (3, 8). *S. lugdunensis* was first described in 1988 (5) and is isolated mostly as a causative agent of skin and soft-tissue infections (4, 5, 7). The clinical course of infections due to *S. lugdunensis* is known to resemble that of infections due to *S. aureus* (7, 14). In recent years, this pathogen has been reported to cause a wide variety of more serious infections, including brain abscess, sepsis, chronic osteomyelitis, and infective endocarditis (5, 7, 11, 14). Only in one report, dealing with the frequency of CNS as causative agents of CAPD-associated peritonitis, was it mentioned that 3 of 127 isolates of CNS derived from 106 episodes of CAPD peritonitis in 46 patients had been identified as *S. lugdunensis* (12). That report did not detail the clinical findings, diagnosis, therapy, or outcome of the *S. lugdunensis*-associated cases. Since then, to the best of our knowledge, no further case of peritonitis due to *S. lugdunensis* has been reported. In the present case, *S. lugdunensis* was recovered in pure culture from the exit site, the tunnel abscess, and the peritoneal fluid of a CAPD patient soon after catheter implantation. The clinical course required intensive-care observation and therapy.

A 36-year-old man with a diagnosis of end-stage renal disease was admitted to our hospital for implantation of a disconnect catheter allowing therapy by CAPD. Due to rolling up of the catheter inside the peritoneum and subsequent problems with drainage of the peritoneal fluid, CAPD was stopped 7 days after the implantation. The patient developed abdominal pain, fever (39.2°C), and purulent drainage and erythema of the skin at the catheter-epidermal interface. Ultrasound examination of the abdomen showed an abscess approximately 3 cm in diameter within the catheter tunnel. Pathological laboratory findings were as follows: leukocytes, 19,300/μl of peripheral blood and 1,650/μl of peritoneal fluid; C-reactive protein, 0.153 g/liter of serum. For microbiological investigations, peritoneal fluid and abscess material were aspirated and a smear was taken from the exit site. Since Gram staining of all three specimens showed the presence of gram-positive cocci, antibiotic treatment was started with intraperitoneal adminis-

tration of vancomycin (2 g). Despite this antibiotic treatment, the body temperature further increased to 39.8°C and the cell count in the peritoneal fluid increased to 2,100/μl. Due to this rapid deterioration, the patient had to be admitted to the intensive-care unit. The catheter and the abscess were removed a few hours later by surgery. Peritoneal irrigation of all four peritoneal quadrants was performed, and the antibiotic treatment was switched to intravenous vancomycin (4 × 0.5 g/day) and rifampin (600 mg/day). The postoperative course was uneventful, and hemodialysis was started by means of a Shaldon catheter. Irrigation was stopped after 3 days, and another 6 days later, a Cimono-Brescia shunt was introduced at the right forearm, allowing continuation of hemodialysis. Antibiotic therapy with vancomycin and rifampin was continued for 14 days. Afterwards, the patient was discharged in good condition.

All specimens taken for microbiological investigations yielded pure cultures of CNS. The bacteria were identified as *S. lugdunensis* by ATB 32 STAPH (bioMérieux, La Balmes-les-Grottes, France), by a negative tube coagulase test using rabbit plasma (bioMérieux), by positive test results for clumping factor using human plasma, and also with the Staphaurex Plus (Murex Diagnostics) and the Pastorex Staphplus (Sanofi Pasteur) agglutination tests. *S. lugdunensis* was identified according to the criteria of Freney et al. (5) and Hébert (6). All of the *S. lugdunensis* strains tested yielded identical biochemical reaction patterns in the ATB 32 STAPH system (code 567134700). Disk diffusion testing was performed according to National Committee for Clinical Laboratory Standards performance standards (1) and showed that all of the isolates were susceptible to penicillin, ampicillin, augmentin, cefotiam, cefotaxime, gentamicin, doxycycline, erythromycin, cotrimoxazole, clindamycin, ofloxacin, rifampin, and vancomycin.

The frequency of a microorganism being found to be responsible for a disease depends heavily on the availability of an easy-to-perform diagnostic method with high discriminatory power. In routine laboratory testing, *S. lugdunensis* is most often confused with *S. aureus* or with other CNS (4, 5, 6). Like *S. aureus*, *S. lugdunensis* may be clumping factor positive and, in that case, shows a positive reaction with the commonly used agglutination tests for *S. aureus*, as did our isolates. A positive pyrrolidonyl-arylamidase reaction (PYR), missing utilization of mannitol, a negative tube coagulase test, and ornithine decarboxylation turned out to be discriminative and easy-to-perform tests for presumptive identification of clumping factor-positive *S. lugdunensis*. Within the group of PYR-positive CNS occurring in humans, i.e., *S. haemolyticus*, *S. schleiferi*, *S. xylo-*

\* Corresponding author. Mailing address: Institute of Medical Microbiology and National Reference Center for Streptococci, University Hospital Aachen, D-52057 Aachen, Germany. Phone: 49 241 8089515. Fax: 49 241 8888-483. E-mail: haase@alpha.imib.rwth-aachen.de.

