

## Presence and Characterization of Extraintestinal Pathogenic *Escherichia coli* Virulence Genes in F165-Positive *E. coli* Strains Isolated from Diseased Calves and Pigs

Hojabr Dezfulian,<sup>1</sup> Isabelle Batisson,<sup>1</sup> John M. Fairbrother,<sup>1</sup> Peter C. K. Lau,<sup>2</sup>  
Atef Nassar,<sup>1</sup> George Szatmari,<sup>3</sup> and Josée Harel<sup>1\*</sup>

*Groupe de Recherche sur les Maladies Infectieuses du Porc, Faculté de Médecine Vétérinaire,<sup>1</sup> and Département de Microbiologie et Immunologie, Faculté de Médecine,<sup>3</sup> Université de Montréal, Saint-Hyacinthe, and Environmental Biotechnology Sector, Biotechnology Research Institute, National Research Council of Canada, Montreal,<sup>2</sup> Québec, Canada*

Received 11 January 2002/Returned for modification 22 September 2002/Accepted 15 December 2002

The virulence genotype profile and presence of a pathogenicity island(s) (PAI) were studied in 18 strains of F165-positive *Escherichia coli* originally isolated from diseased calves or piglets. On the basis of their adhesion phenotypes and genotypes, these extraintestinal pathogenic strains were classified into three groups. The F165 fimbrial complex consists of at least two serologically and genetically distinct fimbriae: F165<sub>1</sub> and F165<sub>2</sub>. F165<sub>1</sub> is encoded by the *foo* operon (*pap*-like), and F165<sub>2</sub> is encoded by *fol* (*sfa* related). Strains in group 1 were *foo* and *fol* positive, strains in group 2 were *foo* and *afa* positive, and strains in group 3 were *foo* positive only. The strains were tested for the presence of virulence genes found mainly in extraintestinal pathogenic *E. coli* (ExPEC) strains. Although all the strains were positive for the *papA* variant encoding F11 fimbriae *incD*, *traT*, and *papC*, the prevalence of virulence genes commonly found in PAIs associated with ExPEC strains was highly variable, with strains of group 2 harboring most of the virulence genes tested. *papG* allele III was detected in all strains in group 1 and in one strain in group 3. All other strains were negative for the known alleles encoding PapG adhesins. The association of virulence genes with tRNA genes was characterized in these strains by using pulsed-field gel electrophoresis and DNA hybridization. The insertion site of the *foo* operon was found at the *pheU* tRNA locus in 16 of the 18 strains and at the *selC* tRNA locus in the other 2 strains. Furthermore, 8 of the 18 strains harbored a high-pathogenicity island which was inserted in either the *asnT* or the *asnV/U* tRNA locus. These results suggest the presence of one or more PAIs in septicemic strains from animals and the association of the *foo* operon with at least one of these islands. F165-positive strains share certain virulence traits with ExPEC, and most of them are pathogenic in piglets, as tested in experimental infections.

*Escherichia coli* is a frequent cause of intestinal and extraintestinal diseases in humans and animals. Typical extraintestinal infections include urinary tract infections, newborn meningitis, polyserositis, and septicemia. All these groups of pathogenic *E. coli* strains have been called extraintestinal pathogenic *E. coli* (ExPEC). The recognized virulence factors of ExPEC include diverse adhesins (e.g., P fimbriae, S/F1C fimbriae, F165 fimbriae, Afa/Dr adhesins, and type 1 fimbriae), toxins (e.g., hemolysin, cytotoxic necrotizing factor, and cytolethal distending toxin), surface antigens (e.g., group II and group III capsules and lipopolysaccharide), invasins (e.g., an invasin responsible for invasion of brain endothelium [IbeA, also called Ibe10]), iron uptake systems (e.g., the aerobactin system), and secretion systems (e.g., type III secretion systems). These virulence factors facilitate colonization and invasion of the host, avoidance or disruption of host defense mechanisms, injury to host tissues, and/or stimulation of a noxious host inflammatory response (34, 50).

Fimbrial antigen complex F165, which exhibits mannose-resistant hemagglutination, is found mostly on *E. coli* strains isolated from piglets and calves with septicemia and/or diarrhea (19) and from humans with septicemia (11). Most F165-positive *E. coli* isolates are not enterotoxigenic or hemolytic; produce aerobactin; are resistant to the bactericidal effects of serum; do not produce verotoxin; and are negative for fimbrial antigens F4, F5, F41, and F6 (24). However, most F165-positive isolates, whether of intestinal or extraintestinal origin, induce septicemia and polyserositis but rarely induce diarrhea or significant enteric lesions in experimentally infected newborn piglets (17, 18). The F165 fimbrial complex consists of at least two serologically and genetically distinct fimbriae: F165<sub>1</sub> and F165<sub>2</sub> (23, 25). The F165<sub>1</sub>A major fimbrial subunit of F165, which is encoded by the *foo* operon, is closely related to that of P fimbriae of serotype F11 but bears a class III G adhesin similar to the P-related (Prs) adhesin of F13 fimbriae, with specificity for the galactose-*N*-acetyl- $\alpha$ -(1-3)-galactose-*N*-acetyl (GalNac-GalNac) moiety (17, 40). F165<sub>2</sub> fimbriae, which are encoded by the *fol* operon, are closely related to F1C fimbriae (26). F165-positive strains possess many of the attributes of ExPEC strains that cause septicemia.

Pathogenicity islands (PAIs), which are large clusters of virulence genes in the bacterial chromosome, have been identi-

\* Corresponding author. Mailing address: Groupe de Recherche sur les Maladies Infectieuses du Porc, Faculté de Médecine Vétérinaire, Université de Montréal, C.P. 5000, Saint-Hyacinthe, Québec J2S 7C6, Canada. Phone: (450) 773-8521, ext. 8233. Fax: (450) 778-8108. E-mail: harelj@MEDVET.UMontreal.CA.

fied in many different bacterial pathogens (22). The PAIs of uropathogenic (UPEC) human strains were the first to be described in *E. coli*. At least four PAIs are present in the genome of UPEC strain 536. PAI I<sub>536</sub>, inserted next to the *selC* tRNA locus, and PAI II<sub>536</sub>, inserted next to the *leuX* tRNA locus, respectively encode the hemolysin and the Prs fimbrial adhesin. On the other hand, PAI III<sub>536</sub> is inserted next to the *thrW* tRNA locus and encodes the S fimbrial adhesin (14). PAI IV<sub>536</sub> is inserted next to the *asnT* tRNA locus and carries the *fyuA* (ferrin yersiniabactin uptake) and *irp1* (iron-repressible protein) through *irp5* genes originally found in the high-pathogenicity islands (HPIs) of various *Yersinia* species (10). Two PAIs were described in UPEC strain J96 and encode the hemolysin and P or Prs fimbrial adhesins. They are PAI I<sub>J96</sub>, which is inserted next to the *pheV* tRNA locus, and PAI II<sub>J96</sub>, which also encodes cytotoxic necrotizing factor type 1 and which is inserted next to the phenylalanine-specific *pheR* (also known as *pheU*) tRNA locus (22).

The population structure of *E. coli*, as represented by the *E. coli* reference (ECOR) collection (46), is thought to be clonal, since it comprises four major clonal groups called A, B1, B2, and D (28). Most ExPEC isolates belong to ECOR group B2 and, to a lesser extent, ECOR group D (5). Recently, it was shown that ECOR group B2 and D strains more often carry certain virulence genes, including *hly*, *pap*, *sfa*, and *kps*, than strains from the other ECOR groups (8). These studies suggest that a cluster of *E. coli* strains acquired virulence genes by horizontal transfer, thus defining a highly virulent group. The link between the B2 phylogenetic group and virulence in ExPEC was recently confirmed in studies with experimentally infected mice (48). In this study we have found that F165-positive strains from pigs and calves share virulence genes found in ExPEC strains and that one or more PAIs are present. Certain virulence attributes found in F165 strains can be associated with pathogenicity, as tested in our model. Although F165-positive strains have been isolated from clinical cases, they belong to the groups, according to the phylogenetic scheme proposed by other investigators (13, 24, 28), that are less virulent than human ExPEC strains. Moreover, the ExPEC genes of animal F165-positive *E. coli* strains were characterized for their relationship with PAIs.

## MATERIALS AND METHODS

**Bacterial strains.** The 18 F165-positive *E. coli* isolates used in this study were obtained from the *Escherichia coli* Laboratory of the Faculté de Médecine Vétérinaire, Université de Montréal, Saint-Hyacinthe, Québec, Canada. These isolates were originally isolated from calves or piglets and were isolated either from the intestines of animals with septicemia or from extraintestinal tissues of animals with septicemia. All isolates were from different animals and farms. Other strains and/or plasmids used as controls for assay development included *E. coli* strains MG1655, 536 (*fyuA irp1 irp2*), CFT073, and J96 (*papA papC papG* alleles I and III *sfa/foc fimA kpsMT III hlyC cnf1*); *E. coli* O157:H7 (*E-hlyA espP etpD katP*); *Yersinia pseudotuberculosis* IP32637 (for detection of the FyuA protein); and *E. coli* strains IA2 (*papG* allele II), CS31A (*clpG fl7c-A*), JM109/pAH1010 (*nfa* allele I), pKT107 (*traT*), HB101/pILL 1194 (*afa-7*), and HB101/pILL 1224 (*afa-8*). The strains were stored at  $-70^{\circ}\text{C}$  in Luria broth plus 15% glycerol until they were ready for use. The *E. coli* strains were routinely grown overnight in Luria-Bertani (LB) broth with shaking at  $37^{\circ}\text{C}$ .

**Detection of virulence genes and rapid phylogenetic analysis by PCR.** The primers used for amplification of the virulence genes were derived from different sources or were designed from available nucleotide sequences (Table 1). All strains were tested for the presence of the three alleles of *papG* and 11 variants of *papA* by established specific PCR assays (30, 36). The virulence determinants

examined in this work were chosen because of their association with *E. coli* strains causing extraintestinal infections. The strains were screened for PCR products specific for genes encoding adhesins: P (*pap*), S/F1C (*sfa*), type 1 (*fimA*), F17c (*fl7c-A*), and Afa/Dr (*afa*) adhesins; M blood group antigen-specific M fimbriae (*bma*); nonfimbrial adhesin type 1 (*nfa*); and Iha (*iha*) nonhemagglutinating adhesin (55). The toxin genes screened for were those encoding alpha-hemolysin (*hly*), *cnf*, and *E-hlyA* (53) and *cdtB* (cytolethal distending toxin) (34). The siderophore systems screened for included aerobactin (*iucD*), yersiniabactin (*fyuA*), and *iroN*<sub>*E. coli*</sub>, a novel catechol siderophore receptor (33). The *cvaC* gene, which encodes colicin V (34), and the *iss* and *traT* genes, which encode proteins that increase resistance to serum (6, 12), were included in this study, as were other genes found in ExPEC, such as the gene for invasion of brain endothelium (*ibeA* or *ibe10*) (34) and capsular polysaccharide synthesis genes *kpsMT* II (e.g., K1, K5, and K12) and *kpsMT* III (e.g., K10 and K54) (34). Proteases encoding the *ompT* (outer membrane protein T), *tsh* (15), and *espP* (53) genes were also included in the study.

Bacterial DNA was released by the boiling method. PCR assays were carried out with the reagents and by the protocols supplied by the manufacturer (Pharmacia). The total reaction mixture volume was 50  $\mu\text{l}$ , which contained 10  $\mu\text{l}$  of supernatant from the boiled bacteria, the appropriate oligonucleotide primers at concentrations of 0.5  $\mu\text{M}$  each, 200  $\mu\text{M}$  (each) deoxynucleotide triphosphate, 5  $\mu\text{l}$  of  $10\times$  PCR buffer (Pharmacia), and 2.5 U of *Taq* DNA polymerase (Pharmacia). The reactions with all the reaction mixtures included an initial denaturation at  $94^{\circ}\text{C}$  for 5 min and a final cycle of primer extension at  $72^{\circ}\text{C}$  for 7 min. The thermocycler reaction conditions for each primer pair were calculated on the basis of the annealing temperature and the length of the product size. PCR-amplified DNA was analyzed on 0.8 to 1% agarose gels by electrophoresis.

The phylogenetic group to which the *E. coli* strains belonged was determined by a PCR-based method, as described by Clermont et al. (13). Briefly, a two-step triplex PCR was performed directly with 3  $\mu\text{l}$  of each of the bacterial lysates. The primer pairs used were ChuA.1-ChuA.2, YjaA.1-YjaA.2, and TspE4C2.1-TspE4C2.2 (Table 1). The PCR steps were as follows: denaturation for 4 min at  $94^{\circ}\text{C}$ ; 30 cycles of 5 s at  $94^{\circ}\text{C}$ , 10 s at  $59^{\circ}\text{C}$ , and 30 s at  $72^{\circ}\text{C}$ ; and a final extension step of 5 min at  $72^{\circ}\text{C}$ . The data from the three amplifications resulted in assignment of the strains to phylogenetic groups as follows: *chuA* positive and *yjaA* positive, group B2; *chuA* positive and *yjaA* negative, group D; *chuA* negative and TspE4.C2 positive, group B1; *chuA* negative and TspE4.C2 negative, group A (13).

DNA probes specific for *pheU* tRNA, *thrW* tRNA, *selC* tRNA, *leuX* tRNA, *meiV* tRNA, *afaE-8*, *fimA*, *papC*, *sfaDE*, *fl7c-A*, *iucD*, *irp2*, *fyuA*, *traT*, *iss*, *cvaC*, *hlyC*, *ehlyA*, *cnf*, *cdt3*, and *cdt4* were generated by PCR with suitable primer pairs (Table 1). Following amplification, the PCR products were run on a 1% agarose gel. Appropriate fragments were cut from the gel, concentrated by ethanol precipitation, and/or purified with the Qiaquick PCR purification kit (Qiagen, Inc. Chatsworth, Calif.) and then radiolabeled with [ $\alpha$ - $^{32}\text{P}$ ]dCTP by using a random priming kit for labeling of oligonucleotides (Pharmacia LKB Biotechnology Inc., Baie d'Urfé, Québec, Canada), according to the instructions of the manufacturer.

**Detection of FyuA by immunoblotting.** The positive control bacterial strain used for PCR amplification of *fyuA* and for detection of the FyuA protein was *Y. pseudotuberculosis* IP32637 (kindly provided by Elisabeth Carniel). For this purpose, bacteria were grown overnight in LB broth with shaking at  $28^{\circ}\text{C}$  (*Yersinia*) or  $37^{\circ}\text{C}$  (*E. coli*) (54). Iron-restricted medium was prepared by the addition of 2,2'-dipyridyl to a final concentration of 50  $\mu\text{M}$ , and iron-rich medium was prepared by adding 150  $\mu\text{M}$   $\text{FeCl}_3$ . The same quantity of bacteria of each strain was sonicated, the cell debris was removed, and the supernatant was centrifuged at  $20,000\times g$  for 1 h at  $4^{\circ}\text{C}$ . The pellet fraction was incubated for 1 h in 1% sodium dodecyl sulfate (SDS) at room temperature and centrifuged again (at  $20,000\times g$  for 1 h). The supernatant was dialyzed against water and subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis in a 12% polyacrylamide gel. The gel was electroblotted onto a nitrocellulose membrane, and FyuA was detected with anti-FyuA antiserum kindly provided by J. Heesemann. Goat anti-rabbit antibody conjugated to horseradish peroxidase was used as the secondary antibody, and the reaction was developed with hydrogen peroxide and 4-chloro-1-naphthol as the substrate.

**PFGE and Southern hybridization.** The pulsed-field gel electrophoresis (PFGE) and Southern hybridization procedure used was a modified method of Gautom (21). The bacteria were grown overnight at  $37^{\circ}\text{C}$  in LB broth and centrifuged at  $1,500\times g$  for 15 min at  $4^{\circ}\text{C}$ . The cells were washed in SE (75 mM NaCl, 25 mM EDTA [pH 7.5]) and centrifuged, and the optical density of the cells at a wavelength of 600 nm was adjusted to between 1.60 and 1.80 in cold TE buffer (10 mM Tris-HCl, 1 mM EDTA [pH 8.0]). For each bacterial strain, 500  $\mu\text{l}$  of suspension was added to 500  $\mu\text{l}$  of prewarmed ( $60^{\circ}\text{C}$ ) 1.5% low-melting-

TABLE 1. PCR primers and conditions used in this study

Gene	Primer pair (5'-3') <sup>a</sup>	Annealing temp (°C)	Predicted size (bp)	Reference or source
<i>papC</i>	F GACGGCTGTACTGCAGGGTGTGGCG R. ATATCCTTTCTGCAGGGATGCAATA	60	328	39
<i>sfaDE</i>	F CGGAGGAGTAATTACAAACCTGGCA R. CTCCGGAGAACTGGGTGCATCTTAC	60	410	39
<i>afaBC</i> III	F GCTGGGCAGCAAACCTGATAACTCTC R CATCAAGCTGTTTGTTCGTCGCCG	60	793	39
<i>afaE-7</i>	F GCTAAATCAACTGTTGATGTT R. GGACAATCCAAATGGCGAATTA	65	618	38
<i>afaE-8</i>	F CTAACCTGCCATGCTGTGACAGTA R. TTATCCCCTGCGTAGTTGTGAATC	65	302	38
<i>clpG</i>	F. GGGCGCTTCTCTCCTTCAAC R CGCCCTAATTGCTGGCGAC	55	402	4
20K ( <i>f17c-A</i> )	F GCAGAAAATTCAATTTATCCTTGG R CTGATAAGCGATGGTGTAAATTAAC	55	537	4
<i>fimA</i>	<i>fimA</i> 05 GTTGATCAAACCGTTCAG <i>fimA</i> 16 AATAACGCGCCTGGAACG	55	331	42
<i>bmaE</i>	F ATGGCGCTAACTTGCCATGCTG R AGGGGGACATATAGCCCCCTTC	60	507	34
<i>nfaE</i>	F GCTTACTGATTCTGGGATGGA R CGGTGGCCGAGTCATATGCCA	60	559	34
<i>iha</i>	F. CTGGCGGAGGCTCTGAGATCA R TCCTTAAGCTCCCGCGGCTGA	55	827	33
<i>iucD</i>	F AAGTGTCGATTTTATTGGTGTGA R CCATCCGATGTCAGTTTTCTG	60	760	27
<i>fyuA</i>	<i>FyuA</i> . 1A. GGCGGCGTGCGCTTCTCGCA <i>FyuArp</i> CGCAGTAGGCACGATGTTGTA	60	209	1
<i>irp1</i>	F GCGATGTTTAACCCCGATT R TGCCTGGAAACCCTGAGACT	55	1,691	1
<i>irp2</i>	<i>Irp2</i> . 15: GTTGCTGTCCATCAAGCACG <i>Irp2</i> . 18 GCCGGAAAGCCTGGCCTTTA	60	1,243	1
<i>iroN</i>	F AAGTCAAAGCAGGGTTGCCCCG R GACGCCGACATTAAGACGCAG	55	665	33
<i>kpsMT</i> II	F GCGCATTTGCTGATACTGTTG R CATCCAGACGATAAGCATGAGCA	60	272	34
<i>kpsMT</i> III	F TCCTCTTGCTACTATTCCCCCT R AGGCGTATCCATCCCTCCTAAC	60	392	34
<i>traT</i>	F GGTGTGGTGCATGAGCACAG R CACGGTTCAGCCATCCCTGAG	60	290	34
<i>iss</i>	F TCACATAGGATTCTGCCG R AGAAATCAAAAGGTGGCC	50	607	This study
<i>cvaC</i>	F CACACACAAACGGGAGCTGTT R CTTCGCCGACGATAGTTCCAT	55	680	34
<i>ompT</i>	<i>ompT</i> -f ATCTAGCCGAAGAAGGAGGC <i>ompT</i> -r CCCGGTCCATAGTGTTCATC	60	559	32
<i>espP</i>	<i>espP</i> -A AAACAGCAGGCACTTGAACG <i>espP</i> -B GGAGTCGTCAGTCAGTAGAT	56	1,830	9

Continued on following page

TABLE 1—Continued

Gene	Primer pair (5'-3') <sup>a</sup>	Annealing temp (°C)	Predicted size (bp)	Reference or source
<i>ibe10</i>	F AGGCAGGTGTGCGCCGCGTAC R TGGTGCTCCGGCAAACCATGC	60	170	34
<i>hlyC</i>	F AGGTTCTTGGGCATGTATCCT R TTGCTTTGCAGACTGCAGTGT	60	556	5
E- <i>hlyA</i>	F GGTGCAGCAGAAAAAGTTGTAG R TCTCGCTGATAGTGTGGTA	57	1,551	52
<i>cnf</i>	F TTATATAGTCGTCAAGATGGA R CACTAAGCTTTACAATATTGAC	50	636	47
<i>tsh</i>	tsh-1 GGTGGTGCCTGGAGTGG tsh-2 AGTCCAGCGTGATAGTGG	53	640	15
<i>chuA</i>	F GACGAACCAACGGTCAGGAT R TGCCGCCAGTACCAAAGACA	59	279	13
<i>yjaA</i>	F TGAAGTGTGAGGAGACGCTG R ATGGAGAATGCGTTCCTCAAC	59	211	13
TspE4 C2	F GAGTAATGTCGGGGCATTCA R CGCGCAACAAAGTATTACG	59	152	13
<i>int</i>	F TCCCTTACCGACGCAAAAATCC R TGCTTCCAGATAATCCGACCAC	58	1,203	37
<i>asnT-int</i>	IntSB ATCGCTTTGCGGGCTTCTAGGT AsnTSB GAACGGCGGACTGTTAAT	60	1,393	1
<i>asnV-int</i>	F GACAGCAAAACAAACAAAAA R TGCTTCCAGATAATCCGACCAC	60	1,500	37
<i>asnU-int</i>	F TTTTCGCTGTAAAGATGTGCC R TGCTTCCAGATAATCCGACCAC	60	1,500	37
IS100	F ATTGATCCACCGTTTTACTC R CGAACGAAAGCATGAAACAA	60	963	1
<i>selC</i> tRNA	F GAGCGAATATTCCGATATCTGGTT R CCTGCAAATAAACACGGCGCAT	60	527	45
<i>phcU</i> tRNA	F TTC AGA AAA TCT CAT CAG TCG C R CAG AAA CAC AGA AAA GAA GCG A	60	475	This study
<i>thrW</i> tRNA	F TGT TTA CGT TAA CGC CTC TAC G R TGA GCT AAT TTG TTC GAG CTT T	60	586	This study
<i>leuX</i> tRNA	F TGC TGA AAA TTT CAG CAC TTA G R ATT TTT TGC TTT CCC TCA TAA C	55	520	This study
<i>metV</i> tRNA	F GGT AAA AAA AAG GTT GCA TGA A R TAA AAA TCA AGTTGA ACA GGC C	55	628	This study
<i>afaE-8</i> and <i>pheV</i>	F GATTCACTACTACGCAGGGG R ATTTGATTGACGAGACGAGGGCGAA	65	1,800	38
<i>afaE-8</i> and <i>pheU</i>	F GATTCACTACTACGCAGGGG R CCGAACTCAACCAGATTCTCCCC	65	1,850	38

<sup>a</sup> F, forward; R, reverse.

point agarose (Sigma), and the mixture was immediately poured into a block former (plug mold). After solidification, the plugs were incubated overnight at 50°C in 1 ml of lysis buffer (1% [wt/vol] *N*-laurylsarcosine, 0.5 M EDTA [pH 9.5]) supplemented with 1 mg of proteinase K per ml. To inactivate the proteinase K, the plugs were then washed three times (each time 1 h) in TE buffer supplemented with 1 mM phenylmethylsulfonyl fluoride (PMSF) at room temperature

and were run for 45 min at 60 V in 0.5× TBE buffer (44.5 mM Tris-borate, 12.5 mM EDTA [pH 8.3]). This premigration step removed degraded or extrachromosomal DNA from the gel and, thus, strongly reduced the smear background in PFGE (7). Macrorestriction of genomic DNA embedded in agarose blocks with restriction enzymes *Xba*I and *Sfi*I was done as described by Birren and Lai (7). The DNA was separated on an AutoBase machine (Mandel Scientific Com-

TABLE 2. Virulence genotypes of 18 F165-positive *E. coli* strains isolated from swine and cattle<sup>a</sup>

Strain	Group	Adhesins								Iron uptake					Serum resistance			Proteases	
		<i>papC</i>	<i>sfaDE</i>	<i>afaE-8</i>	<i>clpG</i>	20K	<i>fimA</i>	<i>bmaE</i>	<i>iha</i>	<i>iucD</i>	<i>fyuA</i>	<i>irp1</i>	<i>irp2</i>	<i>iroN</i>	<i>kpsMT III</i>	<i>traT</i>	<i>iss</i>	<i>cvaC</i>	<i>ompT</i>
5131	1	+	+	-	-	-	+	-	+	-	-	-	+	-	+	+	+	-	-
4787	1	+	+	-	-	-	+	-	+	-	-	-	+	-	+	+	+	+	-
6389	1	+	+	-	-	-	+	-	+	-	-	-	+	-	+	+	-	+	-
1776	1	+	+	-	-	-	+	-	+	-	-	-	+	-	+	-	-	-	-
393	1	+	+	-	-	-	+	-	+	-	-	-	+	-	+	-	-	+	-
2325	2	+	-	+	+	-	+	+	+	+	+	+	-	-	+	-	-	+	-
3292	2	+	-	+	-	+	+	+	+	+	+	+	-	-	+	+	+	+	-
7867	2	+	-	+	-	-	+	+	+	+	+	+	-	-	+	+	+	+	-
3863	2	+	-	+	-	-	+	+	+	+	+	+	-	-	+	+	+	+	-
1401	2	+	-	+	+	+	+	+	+	+	+	+	-	-	+	-	-	+	+
3984	2	+	-	+	-	-	+	+	+	-	-	-	+	-	+	+	-	+	-
215	2	+	-	+	+	-	-	+	+	-	-	-	-	+	+	-	-	+	+
2313	3	+	-	-	-	-	+	-	+	-	-	-	-	-	+	-	-	-	-
2878	3	+	-	-	+	-	+	-	+	-	-	-	-	-	+	-	-	-	+
1195	3	+	-	-	-	-	+	-	+	+	+	+	-	-	+	-	+	+	-
1616	3	+	-	-	-	-	+	-	+	+	-	+	+	-	+	+	+	-	-
3373	3	+	-	-	-	-	+	-	+	+	+	+	-	-	+	-	+	+	-
3719	3	+	-	-	-	-	+	-	+	-	-	-	+	-	+	+	+	+	-

<sup>a</sup> The genotypes shown are for virulence factors detected by PCR and/or probe hybridization. + and -, presence and absence of a gene, respectively. *papC*, gene responsible for pilus assembly that is found in the central region of *pap* operon; *sfa/focDE*, genes found in the central region of *sfa* (S fimbriae) and *foc* (F1C fimbriae) operons; *afaE-8*, afimbrial adhesin gene, *clpG*, gene for structural subunit of CS31A surface antigen; 20K, F17C fimbria; *fimA*, gene for type 1 fimbrial major subunit; *bmaE*, gene for blood group M-specific adhesin; *iha*, IrgA (iron-regulated gene A of *Vibrio cholerae*) homologue adhesin; *iucD*, gene that encodes a membrane-bound enzyme synthesizing N<sub>6</sub>-hydroxylysine, the first product of the aerobactin biosynthesis pathway; *iroN*, gene for a novel catecholate siderophore receptor; *fyuA*, gene for *Yersinia* siderophore receptor (ferric yersiniabactin uptake); *irp1* and *irp2*, genes for iron-repressible high-molecular-weight proteins involved in the production of yersiniabactin; *kpsMT III*, gene for group III capsular polysaccharide synthesis (e.g., K3, K10, and K54); *traT*, gene for surface exclusion, serum survival (outer membrane protein); *iss*, gene for increased serum survival; *cvaC*, colicin V, located on conjugative plasmids (*traT*, *iss*, and antimicrobial resistance); *ompT*, gene for outer membrane protein T (protease); *espP*, gene for *E. coli* serine protease.

pany Ltd.), which uses the principle of zero integrated field electrophoresis, a variation of the field inversion gel electrophoresis configuration, which gives sharper bands and minimum band inversion (7). PFGE was performed with a 0.8% agarose gel in 1× TBE buffer at room temperature without cooling or circulation of buffer. PFGE Marker I (Roche Diagnostics Corporation, Laval, Quebec, Canada), a bacteriophage lambda DNA digest consisting of a ladder (20 bands) of increasing size from 48.5 kb (monomer) to approximately 1,000 kb (20-mer), was included as a DNA size standard. The digested genomic DNA separated in agarose gels was transferred to positively charged nylon membranes (Immobilon-Ny<sup>+</sup>; Millipore Corporation, Bedford, Mass.) in accordance with the instructions of the manufacturer and hybridized under stringent conditions as described by Sambrook et al. (51).

**Determination of plasmid profiles by nuclease S1 treatment and PFGE.** To determine the molecular sizes of the bacterial high-molecular-weight plasmids accurately, plasmid profiles were determined by using nuclease S1 treatment followed by PFGE, as described elsewhere (3). Half of each plug was rinsed with 200 µl of 1× nuclease S1 buffer (50 mM NaCl, 30 mM sodium acetate [pH 4.5], 5 mM ZnSO<sub>4</sub>) at room temperature for 20 min before being incubated with 1 U of *Aspergillus oryzae* nuclease S1 (Sigma, St. Louis, Mo.) in 200 µl of 1× nuclease S1 buffer at 37°C for 45 min. The reaction was stopped by adding 10 µl of 0.5 M EDTA (pH 8.0). Electrophoresis was performed on a CHEF DR II PFGE apparatus (Bio-Rad Laboratories, Hercules, Calif.) in 0.5× TBE buffer at 14°C. The gels were then run at pulse ramp times ranging from 2 to 35 s for 20 h at a constant voltage of 200 V. For size determination, a 48.5-kb bacteriophage lambda DNA ladder (Roche Diagnostics Corporation, Laval, Quebec, Canada) was used as a molecular size marker. After electrophoresis, the gels were stained with ethidium bromide and were photographed with an AlphaImager 2000 camera (Alpha Innotech Corporation, San Leandro, Calif.).

**Experimental infections.** Strains were tested for pathogenicity by experimental infection of colostrum-deprived newborn piglets. When the piglets were 2 days old, they were intragastrically given 1 ml of an overnight culture in tryptic soy broth containing approximately 10<sup>9</sup> CFU of *E. coli* ml<sup>-1</sup> diluted in 20 ml of 0.1% peptone water. The pigs were observed for clinical signs and were necropsied when moribund or at 4 days after inoculation if no clinical signs were observed. Bacteriological and pathological examinations were done as described previously (17, 18).

RESULTS

**Virulence factor profiles including profiles for *papG* and *papA* alleles.** A total of 18 F165-positive septicemic *E. coli* strains from animals were originally selected from among a group of isolates found to be positive with probes for *pap*, *sfa*, or *afa* on colony hybridization. Only single copies of the *pap*-, *sfa*-, and *afa*-related sequences were found among these isolates (23, 41). These isolates had been divided into three groups on the basis of their adhesion phenotypes and genotypes. Isolates in group 1 were *pap* and *sfa* positive, isolates in group 2 were *pap* and *afa* positive, and isolates in group 3 were *pap* positive only (Table 2). Among 11 recognized variants of *papA*, the major pilin gene of *E. coli* P fimbriae, which are termed F7-1, F7-2, and F8 to F16 (36), only the F11 *papA* variant was detected in all 18 strains. The PapG adhesin occurs in three known molecular variants, encoded by alleles I to III of the corresponding *papG* gene (29). Allele III was detected in all strains in group 1, all of which belonged to serogroup O115, and in one strain in group 3, which belonged to serogroup O9. All of these strains were of porcine origin. Although the strains in groups 2 and 3 were positive for the *pap* operon, they were negative for all three of the *papG* alleles tested. All strains were positive for *iucD*, *traT*, and *papC* and were negative for cytotoxin genes commonly found in PAIs associated with ExPEC strains (*cnf*, *hlyC*). All strains were negative for *afaE-3*, *afaE-7*, *nfaE*, *cdt3*, *cdt4*, *E-hlyA*, *ibeA* or *ibe10*, *kpsMT II*, and *tsh*.

Strains in group 1 (*pap* and *sfa* positive) were also positive for *fimA* and *iroN*. Some strains were also positive for *iss*, *cvaC*,




FIG. 1. (a) Macrorestriction patterns of *Xba*I-digested genomic DNA from F165-positive strains analyzed by PFGE. (b and c) Southern blot of F165-positive strains digested with *Xba*I and separated by PFGE (b, hybridization with a *pheU* tRNA gene-specific probe; c, hybridization with a *papC*-specific probe). The asterisks on the right sides of certain fragments indicate the fragments that hybridized with both probes. Lanes: M, bacteriophage  $\lambda$  (PFGE ladder); 1, 5131; 2, 4787; 3, 1776; 4, 2325; 5, 3292; 6, 7867; 7, 3863; 8, 215; 9, 2878; 10, 1195; 11, 1616; 12, 3373; and 13, 3719.

and *ompT* (Table 2). Strains in group 2 (*pap* and *afa* positive) harbored the chromosomal *afaE-8* operon, which is specific for human and bovine septicemia and which seems to be very similar to the M agglutinin of uropathogenic strains from humans (38). Strains in group 2 possessed most of the virulence factors for which tests were conducted. They were all positive for *bmaE*, *iucD*, *traT*, and *ompT*. Three strains in this group (strains 2325, 1401, and 215) were positive for *clpG*, which encodes capsule-like antigen CS31A (4). Two strains (strains 3292 and 1401) were positive for the F17c (also known as 20K) fimbria. Four strains were positive for all the iron uptake systems for which tests were conducted. Four strains (strains 3292, 7867, 3863, and 3984) were positive for *iss*, and three strains (strains 3292, 7867, and 3863) were positive for *cvaC* (colicin V), which is carried by conjugative plasmids. Strain 215 was the only strain in group 2 that was positive for *kpsMT* III and negative for *fimA*. Although *iucD* and *traT* were found in strains in group 3 (*pap* positive), the presence of virulence genes was more variable. Strain 2878 was the only strain in this group that was positive for *clpG*. Interestingly, *espP* (*E. coli* serine protease), which is found in most enterohemorrhagic strains (9), was detected in three bovine F165-positive strains in groups 2 and 3 (Table 2).

**Locations of PAIs encoding fimbrial operons.** Many of the PAIs identified are associated with tRNA loci (22). To study the regions containing the *foo* (*pap*-like), *fol* (*sfa*-related), and *afaE-8* operons, DNA from PFGE separations was transferred to membranes and was hybridized with different tRNA gene-specific probes. For most strains (16 of 18 F165-positive strains), the band that hybridized with the probe for the *foo* operon cohybridized with the *pheU* tRNA gene-specific probe. This suggests that for these strains the insertion site of the putative PAIs encompassing the *foo* operon may be within the *pheU* locus. In two other strains the band that hybridized with the probe for the *foo* operon cohybridized with the *selC* tRNA gene-specific probe. This suggests that for these two strains, the insertion site of the PAIs encompassing the *foo* operon may be within the *selC* locus. To establish whether the virulence genes are linked in the same segment of the chromosome,

suggesting their linkage in a PAI, the PFGE bands generated by digestion with two restriction enzymes (*Xba*I and *Sfi*I) were hybridized with probes specific for *papC*, *sfaDE*, *afaE-8*, *fimA*, and *F17c-A*. None of these genes were located on a single restriction fragment, regardless of the restriction enzyme used, suggesting that either there are unknown virulence genes within these putative PAIs or they only contain fimbrial operons as a sole virulence factor.

Sets of PCR primers that could amplify the intact regions containing the tRNA genes were used to confirm that some fimbrial operons in F165-positive strains were inserted within or near tRNA loci (Table 1). We investigated whether the insertion sites of the fimbrial operons in the F165-positive isolates resided within the *pheU*, *thrW*, *selC*, *leuX*, and *metV* tRNA loci (data not shown). Our results indicated that, for most strains, the *pheU* locus was not amplified and that for two strains in group 3 the *selC* locus was not amplified. This suggests that in these isolates these tRNA loci were rearranged or disrupted. Different primer pairs covering the phenylalanine-specific tRNA genes (*pheU* and *pheV*) and the *afa-8* gene were used to determine the insertion site of the *afa-8* PAI (Table 1). In contrast to a recent study (38) on the *afa-8* PAI, the insertion site of this *afa-8* PAI in our group 2 strains was not detected in any of the *phe* tRNA loci.

For 16 of 18 strains, the *pheU* tRNA gene-specific probe hybridized with the fragment that contained *foo*, which is similar to PAI II<sub>196</sub>, which is also inserted next to the *pheU* locus, at 94 min of the *E. coli* K-12 chromosomal map (22) (Fig. 1). Furthermore, the *pheU* tRNA gene-specific probe did not hybridize with the fragment containing the *afa-8* operon. For only two group 2 strains (strains 2325 and 3292), the *selC* tRNA gene-specific probe hybridized with the fragment that contains the *foo* operon, which is similar to PAI I<sub>536</sub> (22). Hybridization with *foo* (*pap*-like) and *fol* (*sfa*-related) gene-specific probes for strains in group 1 showed that these two operons are not physically linked. Although *thrW* tRNA was disrupted in all strains in group 1 that were positive for both *foo* and *fol*, the *thrW* tRNA gene-specific probe did not hybridize with the fragment containing the *foo* or *fol* operon. This is different

from the finding for *E. coli* 536, in which an *sfa* operon homologous to *tot* was inserted next to *thrW* tRNA (PAI III<sub>536</sub>) (14). For the group 2 strains, hybridization with *foo* and *afaE-8* gene-specific probes suggested that these two operons are not on the same fragment.

The PFGE patterns, which were heterogeneous among the different strains, were similar within the particular groups. PFGE analysis demonstrated that the most closely related strains were within the same group but were also of the same origin.

**Presence of HPI in F165-positive strains.** The HPI-specific genes (*irp2*, *irp1*, and *fyuA*) were found in 8 of 18 strains (only 7 of the 8 strains had all three genes); 5 of these strains belonged to group 2, and 3 of these strains belonged to group 3 (Table 2). Strain 1616 was positive for both *irp2* and *fyuA* but was negative for *irp1*. In *Yersinia* spp. and *E. coli*, the site of HPI integration into the bacterial chromosome shares homology with the P4 bacteriophage attachment site and is located within an *asn* tRNA locus (10). To determine whether the HPIs of these strains were bordered by an *asn* tRNA gene locus, a set of primers whose sequences covered a portion of the *int* and *asn* genes was used (Table 1). The insertion site of the HPI in six of eight strains was next to the *asnT* tRNA gene, as demonstrated by PCR with primer pairs specific for *asnT* and *int*. In agreement with a recent report (1), the size of the PCR product was smaller than that in *Y. pseudotuberculosis* (1,100 bp instead of the expected 1,393 bp) in two strains (strains 2325 and 3292), suggesting that a deletion of approximately 300 bp had taken place in this region. Interestingly, in one strain (strain 1401), amplification by PCR produced a band of the expected size and two other smaller bands (1,200 and 1,100 bp). Two strains (strains 1195 and 1616) did not yield any amplification product with three different *asn*-specific primers (specific for *asnT*, *asnV*, and *asnU*) or the *int*-specific primer. All of these strains harboring HPIs were positive for IS100, as demonstrated by use of specific primers.

Certain *E. coli* strains (up to 3% of *irp2*-positive strains) carry a truncated *fyuA-irp* gene cluster, with deletions proceeding from *fyuA* or from both *fyuA* and *irp1* to the 3' end of the HPI. Such spontaneous deletions involve only the 3' part of the *fyuA-irp* gene cluster and abrogate expression of the yersiniabactin receptor (54). In order to analyze the expression of *fyuA* in the eight strains carrying the *Yersinia* HPI, immunoblotting of outer membrane proteins was performed. One specific band of about 67 kDa for FyuA was detected in seven of the eight HPI-positive strains (data not shown).

**Phylogenetic grouping results.** Phylogenetic analysis studies have suggested that *E. coli* is composed of four main phylogenetic groups (groups A, B1, B2, and D) and that virulent ExPEC strains mainly belong to groups B2 and D, whereas commensal strains belong to groups A and B1. Recently, Clermont et al. (13) described a simple and rapid phylogenetic grouping technique based on a triplex PCR. The results of the method, which uses a combination of two genes (*chuA* and *yjaA*) and an anonymous DNA fragment, showed a good correlation with those of reference methods (13). Examination of the strains by the triplex PCR method demonstrated that the strains in group 1, all of which were O115 and positive for *foo* and *tot*, belong to phylogenetic group B1, whereas all the strains in groups 2 and 3 belong to phylogenetic group A.

**High-molecular-mass plasmid profiles.** In a previous study (24) it was shown that F165-positive strains carried several low-molecular-mass plasmids. In order to study the high-molecular-mass plasmids in 18 F165-positive strains, plasmid profiles were determined by using nuclease S1 treatment followed by PFGE. We identified plasmids with molecular masses ranging from 10 to 245 kb. Most of the strains that were positive by hybridization with probes specific for the *iucD*, *traT*, and *iss* genes harbored at least two high-molecular-mass plasmids of 90 to 245 kb. These plasmids were more frequently found in group 2 strains than in group 1 and 3 strains (Fig. 2.).

**Pathogenicities of strains in pigs.** The virulence of the F165-positive group 1 strains in piglets was high (Table 3). All O115 strains caused 100% mortality in pigs within 48 h of inoculation, with manifestation of gross lesions of polyserositis on necropsy. Pathogenicity was variable in strains of groups 2 and 3 (Table 3). Of four group 2 strains tested, three O11 or O9 strains caused 100% mortality in pigs within 48 h of inoculation, with manifestation of gross lesions of polyserositis on necropsy, whereas an O101 strain was nonpathogenic. Of three group 3 strains tested, an O9 strain caused mortality in pigs, whereas two O15 strains were nonpathogenic. All HPI-positive strains tested were pathogenic in piglets. On the other hand, the HPI-negative strains in group 1 tested were pathogenic, whereas those in groups 2 and 3 were less pathogenic or nonpathogenic in piglets. The strains in group 1 belong to phylogenetic group B1, whereas all strains in groups 2 and 3 belong to phylogenetic group A. No obvious relationship between the presence of the genes and pathogenicity in piglets was observed for any of the other virulence determinants.

## DISCUSSION

We have previously defined three phenotypic and genotypic groups among F165-positive animal *E. coli* strains on the basis of the expression of three different fimbrial operons. Strains in group 1 were positive for *foo* (*pap*-like) and *tot* (*sfa*-related) genes, and strains in group 2 or 3 were *foo* and *afa* positive and *foo* positive only, respectively (41). The results of this study suggest the presence of one or more PAIs (*foo*, *tot*, *afaE-8*, HPI) in septicemic strains isolated from animals. Moreover, the tRNA gene locus associated with some PAIs was also determined. The P-like operons were located near the *pheU* or *selC* tRNA gene and HPI sequences near the *asnT* tRNA gene. The insertion site of the HPI in six of eight strains was next to the *asnT* tRNA gene. The insertion site of *afa-8* PAI in group II strains was not detected in any of the *phe* tRNA loci, which is in contrast to the findings in a recent study by Lalioui and Le Bouguenec (38) on the *afa-8* PAI. The insertion site of *tot*, which belongs to the Sfa/F1C family was not next to *thrW* tRNA, unlike the *sfa* operon in *E. coli* 536.

The linkage of the *foo* operon to the *pheU* tRNA gene in 16 strains and to the *selC* tRNA gene in 2 others, the low G+C content of this operon (41 to 43%, compared to a 50.8% content in this operon of *E. coli* K-12 [data not shown]), and its presence in pathogenic strains indicates that this operon is part of a PAI. Moreover, the F165<sub>1</sub> major fimbrial subunit (encoded by the *foo* operon) is closely related to that of P fimbria serotype F11 but bears a class III G adhesin similar to the Prs adhesin of F13 fimbriae (40) and may represent a new member




FIG. 2. Plasmid profiles of 14 F165-positive isolates, determined by nuclease S1 treatment followed by PFGE analysis. Black asterisks, plasmid bands hybridizing with the *traT*-specific probe; white asterisks, plasmid bands hybridizing with the *cvaC*-specific probe. Lanes: M, bacteriophage  $\lambda$  (PFGE ladder); 1, 5131; 2, 4787; 3, 1776; 4, 2325; 5, 3292; 6, 7867; 7, 3863; 8, 1401; 9, 215; 10, 2878; 11, 1195; 12, 1616; 13, 3373; and 14, 3719.

of the family of P fimbriae. Although all F165-positive strains possessed *papC*, only six, all being of porcine origin, had a PCR-detectable *papG* allele corresponding to *papG* allele III. The fact that most porcine strains were positive for allele III whereas all bovine strains were negative for all three known alleles could suggest that there are unknown *papG* alleles among these isolates. The observed predominance of *papG* allele III among the porcine isolates (six of nine) in the present study is consistent with the previously demonstrated predominance of *papG* allele III in canine fecal *E. coli* isolates (35). The *papG* allele III is also the most prevalent *papG* allele among *E. coli* isolates from humans with cystitis and in some reports is prevalent among isolates from patients with bacteremia and pyelonephritis (29, 31). Among the 11 *PapA* variants tested, only the F11 *papA* allele was detected in all 18 strains. In addition to the F165 fimbria, which was detected in the strains tested in this study, which were isolated from pigs and calves with septicemia, interestingly, F11 is the predominant serotype of P fimbriae expressed by *E. coli* strains implicated in avian colisepticemia (16). This suggests that F11 fimbriae are frequently expressed by pathogenic *E. coli* strains isolated from animals with septicemia. Marklund and coworkers (43) have proposed that *E. coli* acquired the *pap* locus after the speciation of *E. coli* and have suggested that the different *pap* genes could have been acquired by horizontal gene transfer. Moreover, they proposed that the recent genetic exchanges involving the entire fimbrial gene clusters have occurred in response to selection pressures exerted by the host.

In addition to *pap*, the F165-positive strains contained multiple virulence-associated genes characteristic of ExPEC

strains. These included *sfa/focDE*, *afaE-8*, *F17c-A*, *clpG*, *bmaE*, *iucD*, *fyuA*, *traT*, *iss*, *ompT*, and *kpsMT III*. Interestingly, no toxin genes commonly associated with ExPEC strains were present in these strains. The secreted proteases are virulence factors in many bacterial and nonbacterial pathogens. In our study, the EspP protease was found in three F165-positive strains. EspP is associated with enterohemorrhagic *E. coli*, induces cytopathic effects, has been shown to cleave coagulation factor V, and may thus promote intestinal hemorrhage (9). This is the first report on the presence of the *espP* gene in ExPEC strains. The absence of many genes that characterize ExPEC strains such as strains J96, 536, and CFT073 may indicate that in F165-positive strains these PAIs could potentially acquire additional virulence genes over time or that the virulence genes within these putative PAIs were deleted. Furthermore, these three strains are isolates from human infections, as opposed to animal infections, and are associated with different pathologies (e.g., strain J96 was from a patient with pyelonephritis). This could reflect a difference in selective pressure that results in this ExPEC strain with specific characteristics.

The majority of PAIs detected in enterobacteria are specific for particular species or even pathotypes. The HPI, first described in pathogenic *Yersinia* (10), has been detected in many enterobacterial species and pathotypes, including both enteroaggregative *E. coli* and ExPEC (54). In addition, more than 30% of *E. coli* isolates from the normal intestinal microflora also carry this island (54). The functional core of the HPI in *Yersinia* consists of 12 genes (*irp1* to *irp9*, *ybtA*, *fyuA*, and *intB*). The mobility of the HPI elements may be associated with an intact integrase gene located at the left junction of the HPI.



TABLE 3. Pathogenicities of F165-positive strains

Group and strain	Source	Serogroup	No. of piglets inoculated	No. of piglets manifesting:		
				Mortality	Macro lesions <sup>a</sup>	Time of death after inoculation (h)
Group 1						
5131	Porcine intestine	O115:K“V165”	2	2	2	36–48
4787	Porcine intestine	O115:K“V165”	4	4	4	36–48
6389	Porcine intestine	O115:K“V165”	2	2	2	36–48
1776	Porcine intestine	O115:K“V165”	2	2	2	36–48
393	Porcine intestine	O115:K“V165”	2	2	2	40–48
Group 2						
2325	Bovine extraintestine	O11:KF12	2	2	2	48
3292	Bovine intestine	O15,O35,O11	NT <sup>b</sup>	NT	NT	NT
7867	Bovine intestine	O9:K28	2	2	2	48
3863	Bovine intestine	O9:K28	NT	NT	NT	NT
1401	Bovine intestine	O9:K28	2	2	1	20–30
3984	Porcine intestine	O101:K32	2	0	0	0
215	Bovine intestine	O?	NT	NT	NT	NT
Group 3						
2313	Bovine extraintestine	O15:KRVC383	2	0	1	0
2878	Bovine intestine	O15:KRVC383	2	0	0	0
1195	Porcine intestine	O9:K28	2	1	1	20
1616	Bovine intestine	O9:K28	NT	NT	NT	NT
3373	Porcine intestine	O9:K28	NT	NT	NT	NT
3719	Porcine intestine	O9:K28	NT	NT	NT	NT

<sup>a</sup> Macro lesions of polyserositis.

<sup>b</sup> NT, not tested.

The *fyuA*, *irp1*, and *irp2* genes were detected in all of the HPI-positive F165-positive strains. However, the yersiniabactin receptor *FyuA* was expressed in only seven of eight strains. The HPIs of F165-positive strains have features in common with the HPI elements of other enterobacteria, including pathogenic yersiniae. We showed that in six F165-positive *E. coli* strains, the HPI is inserted at the *asnT* tRNA gene. The *asnT* locus in *Yersinia* is linked to a gene whose sequence is highly homologous with that of a phage-derived integrase determinant termed *intB*. Furthermore, a copy of the IS100 insertion sequence present in the *Yersinia pestis* HPI (at the 5' end) is also present in the HPIs of F165-positive *E. coli* strains. The HPI elements code for a particular iron uptake system, termed yersiniabactin. The question arises as to whether the HPI elements in *E. coli* indeed represent PAIs and whether they contribute to the survival of the strains in certain ecological niches. Most strains of the family *Enterobacteriaceae* carry genes that encode high-affinity iron uptake systems, such as the enterobactin and aerobactin systems. Thus, the reason for a third iron uptake system in *E. coli*, like that encoded by the HPI, remains to be clarified. It is conceivable that the bacteria use different iron uptake systems, depending on the environment or the stage of infection. However, siderophores not only are involved in providing the microorganisms with iron but also can affect the cellular immune system (e.g., by suppression of T-cell proliferation) (54). This dual role of siderophores may contribute to the pathogenicities of members of the family *Enterobacteriaceae* and as a cause of extraintestinal infections. All HPI-positive F165 *E. coli* strains tested were pathogenic in piglets. Nevertheless, the HPI is not essential for some F165-positive strains, as shown by its absence from group 1 strains, which were pathogenic.

Four main phylogenetic groups (groups A, B1, B2, and D) were described by Herzer et al. (28) on the basis of examination of the 72 strains in the ECOR collection (46) by comparison of several genetic markers by multilocus enzyme electrophoresis. It may be relevant to these observations that isolates belonging to ECOR subgroup A have a significantly smaller genome than those belonging to either subgroup B2 or subgroup D, which have the largest genomes among natural isolates of *E. coli* (8). The *E. coli* genome varies from 4.5 to 5.5 Mb (49). For example, *E. coli* MG1655 has a genome of 4.63 Mb, and *E. coli* J96, which belongs to phylogenetic group B2, has a genome of 5.12 Mb (49). We determined the genome size of strain 4787, which belongs to phylogenetic group B1, by PFGE using three different restriction enzymes (*SfiI*, *NotI*, and *XbaI*) (data not shown). The genome of strain 4787 is estimated to be about 4.9 Mb, which is larger than that of *E. coli* K-12 but smaller than that of pathogenic strain J96.

Although F165-positive strains have been isolated from clinical cases and most of them are pathogenic in experimental infections, they fall into groups A and B1 of the phylogenetic scheme proposed by Herzer et al. (28). In a study by Picard et al. (48), commensal strains belonging to phylogenetic groups A and B1 were shown to be devoid of virulence determinants and did not kill mice. Some strains of phylogenetic groups A, B1, and D were able to kill the mice and possessed virulence determinants. Strains of the B2 phylogenetic group are highly virulent and possess the highest level of virulence determinants. Certain F165-positive strains possess virulence attributes such as P-like fimbriae, HPIs, O serotypes, and phylogenetic groups that can be associated with pathogenicity, as tested in our model. F165-positive strains in phylogenetic group A possessed more virulence factors than those in phy-

logenetic group B1, but the strains in group B1 were more virulent in piglets than the strains in group A. It is not known why strains in our group 1 (which belong to phylogenetic group B1) are more virulent in our experimental model, despite the presence of fewer virulence markers than the numbers of virulence markers in strains of the two other groups. It is possible that some virulence genes have not yet been discovered or that these strains may contain a new PAI(s), or their virulence could be related to the presence of "black holes," genomic deletions that may enhance pathogenicity (44). The roles of the K"V165" O-antigen capsule and F165<sub>1</sub> fimbriae in bacterial pathogenicity were examined by comparing the pathogenicities of mutants in experimental infections. The F165<sub>1</sub> fimbriae and the K"V165" O-antigen capsule were strongly associated with bacterial survival in the extraintestinal organs and in the bloodstreams of infected piglets and with virulence. Fimbrial antigen F165<sub>1</sub> promotes adherence of bacteria to porcine polymorphonuclear leukocytes in vitro but enhances resistance to phagocytosis. O-antigen capsule K"V165" is required for resistance to the bacterial effects of serum and for resistance to phagocytosis by porcine polymorphonuclear leukocytes in vitro (20). It is also possible that certain factors, such as O type (e.g., O115, O78) and K serotype (K1), are important in key steps for survival in the blood and resistance to bacterial killing by the host, i.e., bacterial killing in the presence of complement and phagocytes in extraintestinal tissues and fluids. A recent study suggests that extracellular polysaccharides as well as type 1 fimbriae are preeminent virulence determinants in uropathogenic *E. coli* strains in the murine urinary tract (2).

Thus, F165-positive strains that possess such key pathogenic traits but that lack certain virulence markers found in ExPEC strains would be relatively more pathogenic than commensal organisms but less pathogenic or pathogenic in fewer situations than the strains in phylogenetic group B2. Nevertheless, our study has shown that strains belonging to phylogenetic groups B1 and A (such as F165-positive strains) are not merely commensal but might be considered opportunistic pathogens. Comparison of the presence of virulence factors (Table 2), the ability of the strains to cause disease (pathogenicity) (Table 3), and the phylogenetic group suggest that the pathogenic origin of the strains in group 1 could be related to other virulence factors which are unidentified and/or could be related to host-dependent factors.

#### ACKNOWLEDGMENTS

This work was supported in part by a grant (grant OGP0025120) to J.H. from the Natural Sciences and Engineering Research Council of Canada and grant 0214 from the Formation de Chercheurs et l'Aide à la Recherche.

We thank Charles Dozois (INRS, Institut Armand Frappier, Québec, Canada) for critical review of this article. Control strains and/or plasmids were kindly provided by James R. Johnson, Charles Dozois, Jorg Hacker, Gabriele Blum-Oehler, Ulrich Dobrindt, Harry L. T. Mobley, Thomas A. Russo, Lisa K. Nolan, Elisabeth Carniel, Juergen Heesemann, and Chantal LeBouguenec. Special thanks go to Héléne Bergeron, Céline Forget, and Ramin Behdjani for technical assistance.

#### REFERENCES

- Bach, S., A. de Almeida, and E. Carniel. 2000. The *Yersinia* high-pathogenicity island is present in different members of the family *Enterobacteriaceae*. *FEMS Microbiol. Lett.* **183**:289–294.
- Bahrani-Mougeot, F. K., E. L. Buckles, C. V. Lockett, J. R. Hebel, D. E. Johnson, C. M. Tang, and M. S. Donnenberg. 2002. Type 1 fimbriae and extracellular polysaccharides are preeminent uropathogenic *Escherichia coli* virulence determinants in the murine urinary tract. *Mol. Microbiol.* **45**:1079–1093.
- Barton, B. M., G. P. Harding, and A. J. Zuccarelli. 1995. A general method for detecting and sizing large plasmids. *Anal. Biochem.* **226**:235–240.
- Bertin, Y., C. Martin, J. P. Girardeau, P. Pohl, and M. Contrepois. 1998. Association of genes encoding P fimbriae, CS31A antigen and EAST 1 toxin among CNF1-producing *Escherichia coli* strains from cattle with septicemia and diarrhea. *FEMS Microbiol. Lett.* **162**:235–239.
- Bingen, E., B. Picard, N. Brahimi, S. Mathy, P. Desjardins, J. Elion, and E. Denamur. 1998. Phylogenetic analysis of *Escherichia coli* strains causing neonatal meningitis suggests horizontal gene transfer from a predominant pool of highly virulent B2 group strains. *J. Infect. Dis.* **177**:642–650.
- Binns, M. M., J. Mayden, and R. P. Levine. 1982. Further characterization of complement resistance conferred on *Escherichia coli* by the plasmid genes *traT* of R100 and *iss* of ColV, I-K94. *Infect. Immun.* **35**:654–659.
- Birren, B. W., and E. H. C. Lai. 1993. Pulsed field gel electrophoresis: a practical guide. Academic Press, Inc., San Diego, Calif.
- Boyd, E. F., and D. L. Hartl. 1998. Chromosomal regions specific to pathogenic isolates of *Escherichia coli* have a phylogenetically clustered distribution. *J. Bacteriol.* **180**:1159–1165.
- Brunder, W., H. Schmidt, M. Frosch, and H. Karch. 1999. The large plasmids of Shiga-toxin-producing *Escherichia coli* (STEC) are highly variable genetic elements. *Microbiology* **145**:1005–1014.
- Carniel, E. 1999. The *Yersinia* high-pathogenicity island. *Int. Microbiol.* **2**:161–167.
- Cherifi, A., M. Contrepois, B. Picard, P. Goulet, J. de Rycke, J. Fairbrother, and J. Barnouin. 1990. Factors and markers of virulence in *Escherichia coli* from human septicemia. *FEMS Microbiol. Lett.* **58**:279–283.
- Chuba, P. J., M. A. Leon, A. Banerjee, and S. Palchaudhuri. 1989. Cloning and DNA sequence of plasmid determinant *iss*, coding for increased serum survival and surface exclusion, which has homology with lambda DNA. *Mol. Gen. Genet.* **216**:287–292.
- Clermont, O., S. Bonacorsi, and E. Bingen. 2000. Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Appl. Environ. Microbiol.* **66**:4555–4558.
- Dobrindt, U., G. Blum-Oehler, T. Hartsch, G. Gottschalk, E. Z. Ron, R. Funfstuck, and J. Hacker. 2001. S-fimbria-encoding determinant *sfa* (I) is located on pathogenicity island III (536) of uropathogenic *Escherichia coli* strain 536. *Infect. Immun.* **69**:4248–4256.
- Dozois, C. M., M. Dho-Moulin, A. Bree, J. M. Fairbrother, C. Desautels, and R. Curtiss III. 2000. Relationship between the Tsh autotransporter and pathogenicity of avian *Escherichia coli* and localization and analysis of the Tsh genetic region. *Infect. Immun.* **68**:4145–4154.
- Dozois, C. M., J. Harel, and J. M. Fairbrother. 1996. P-fimbriae-producing septicaemic *Escherichia coli* from poultry possess *fel*-related gene clusters whereas *pap*-hybridizing P-fimbriae-negative strains have partial or divergent P fimbrial gene clusters. *Microbiology* **142**:2759–2766.
- Fairbrother, J., J. Harel, C. Forget, C. Desautels, and J. Moore. 1993. Receptor binding specificity and pathogenicity of *Escherichia coli* F165-positive strains isolated from piglets and calves and possessing *pap* related sequences. *Can. J. Vet. Res.* **57**:53–55.
- Fairbrother, J. M., A. Broes, M. Jacques, and S. Lariviere. 1989. Pathogenicity of *Escherichia coli* O115:K"V165" strains isolated from pigs with diarrhea. *Am. J. Vet. Res.* **50**:1029–1036.
- Fairbrother, J. M., S. Lariviere, and R. Lallier. 1986. New fimbrial antigen F165 from *Escherichia coli* serogroup O115 strains isolated from piglets with diarrhea. *Infect. Immun.* **51**:10–15.
- Fairbrother, J. M., and M. Ngeleka. 1994. Extraintestinal *Escherichia coli* infections in pigs, p. 221–236. In C. L. Gyles (ed.), *Escherichia coli* in domestic animals and humans. CAB International, Wallingford, United Kingdom.
- Gautom, R. K. 1997. Rapid pulsed-field gel electrophoresis protocol for typing of *Escherichia coli* O157:H7 and other gram-negative organisms in 1 day. *J. Clin. Microbiol.* **35**:2977–2980.
- Hacker, J., and J. B. Kaper. 2000. Pathogenicity islands and the evolution of microbes. *Annu. Rev. Microbiol.* **54**:641–679.
- Harel, J., F. Daigle, S. Maiti, C. Desautels, A. Labigne, and J. M. Fairbrother. 1991. Occurrence of *pap*-, *sfa*-, and *afa*-related sequences among F165-positive *Escherichia coli* from diseased animals. *FEMS Microbiol. Lett.* **66**:177–182.
- Harel, J., J. Fairbrother, C. Forget, C. Desautels, and J. Moore. 1993. Virulence factors associated with F165-positive *Escherichia coli* strains isolated from piglets and calves. *Vet. Microbiol.* **38**:139–155.
- Harel, J., C. Forget, J. Saint-Amand, F. Daigle, D. Dubreuil, M. Jacques, and J. Fairbrother. 1992. Molecular cloning of a determinant coding for fimbrial antigen F165(1), a Prs-like fimbrial antigen from porcine septicaemic *Escherichia coli*. *J. Gen. Microbiol.* **138**:1495–1502.
- Harel, J., M. Jacques, J. M. Fairbrother, M. Bosse, and C. Forget. 1995. Cloning of determinants encoding F165(2) fimbriae from porcine septicaemic *Escherichia coli* confirms their identity as FIC fimbriae. *Microbiology* **141**:221–228.

27. Herrero, M., V. de Lorenzo, and J. B. Neilands. 1988. Nucleotide sequence of the *iucD* gene of the pColV-K30 aerobactin operon and topology of its product studied with *phoA* and *lacZ* gene fusions. *J. Bacteriol.* **170**:56–64.
28. Herzer, P. J., S. Inouye, M. Inouye, and T. S. Whittam. 1990. Phylogenetic distribution of branched RNA-linked multicopy single-stranded DNA among natural isolates of *Escherichia coli*. *J. Bacteriol.* **172**:6175–6181.
29. Johnson, J. R. 1998. *papG* alleles among *Escherichia coli* strains causing urosepsis: associations with other bacterial characteristics and host compromise. *Infect. Immun.* **66**:4568–4571.
30. Johnson, J. R., and J. J. Brown. 1996. A novel multiply primed polymerase chain reaction assay for identification of variant *papG* genes encoding the Gal(alpha 1–4)Gal-binding PapG adhesins of *Escherichia coli*. *J. Infect. Dis.* **173**:920–926.
31. Johnson, J. R., J. J. Brown, and J. N. Maslow. 1998. Clonal distribution of the three alleles of the Gal(alpha 1–4)Gal-specific adhesin gene *papG* among *Escherichia coli* strains from patients with bacteremia. *J. Infect. Dis.* **177**:651–661.
32. Johnson, J. R., T. T. O'Bryan, D. A. Low, G. Ling, P. Delavari, C. Fasching, T. A. Russo, U. Carlino, and A. L. Stell. 2000. Evidence of commonality between canine and human extraintestinal pathogenic *Escherichia coli* strains that express *papG* allele III. *Infect. Immun.* **68**:3327–3336.
33. Johnson, J. R., T. A. Russo, P. I. Tarr, U. Carlino, S. S. Bilge, J. C. Vary, Jr., and A. L. Stell. 2000. Molecular epidemiological and phylogenetic associations of two novel putative virulence genes, *iha* and *iroN<sub>E. coli</sub>*, among *Escherichia coli* isolates from patients with urosepsis. *Infect. Immun.* **68**:3040–3047.
34. Johnson, J. R., and A. L. Stell. 2000. Extended virulence genotypes of *Escherichia coli* strains from patients with urosepsis in relation to phylogeny and host compromise. *J. Infect. Dis.* **181**:261–272.
35. Johnson, J. R., A. L. Stell, P. Delavari, A. C. Murray, M. Kuskowski, and W. Gaastra. 2001. Phylogenetic and pathotypic similarities between *Escherichia coli* isolates from urinary tract infections in dogs and extraintestinal infections in humans. *J. Infect. Dis.* **183**:897–906.
36. Johnson, J. R., A. L. Stell, F. Scheutz, T. T. O'Bryan, T. A. Russo, U. B. Carlino, C. Fasching, J. Kavle, L. Van Dijk, and W. Gaastra. 2000. Analysis of the F antigen-specific *papA* alleles of extraintestinal pathogenic *Escherichia coli* using a novel multiplex PCR-based assay. *Infect. Immun.* **68**:1587–1599.
37. Karch, H., S. Schubert, D. Zhang, W. Zhang, H. Schmidt, T. Olschlager, and J. Hacker. 1999. A genomic island, termed high-pathogenicity island, is present in certain non-O157 Shiga toxin-producing *Escherichia coli* clonal lineages. *Infect. Immun.* **67**:5994–6001.
38. Lalioui, L., and C. Le Bouguenec. 2001. *afa-8* gene cluster is carried by a pathogenicity island inserted into the tRNA (Phe) of human and bovine pathogenic *Escherichia coli* isolates. *Infect. Immun.* **69**:937–948.
39. Le Bouguenec, C., M. Archambaud, and A. Labigne. 1992. Rapid and specific detection of the *pap*, *afa*, and *sfa* adhesin-encoding operons in uropathogenic *Escherichia coli* strains by polymerase chain reaction. *J. Clin. Microbiol.* **30**:1189–1193.
40. Maiti, S. N., L. DesGroseillers, J. M. Fairbrother, and J. Harel. 1994. Analysis of genes coding for the major and minor fimbrial subunits of the Prs-like fimbriae F165 (1) of porcine septicemic *Escherichia coli* strain 4787. *Microb. Pathog.* **16**:15–25.
41. Maiti, S. N., J. Harel, and J. M. Fairbrother. 1993. Structure and copy number analyses of *pap*-, *sfa*-, and *afa*-related gene clusters in F165-positive bovine and porcine *Escherichia coli* isolates. *Infect. Immun.* **61**:2453–2461.
42. Marc, D., and M. Dho-Moulin. 1996. Analysis of the *fim* cluster of an avian O2 strain of *Escherichia coli*: serogroup-specific sites within *fimA* and nucleotide sequence of *fimL*. *J. Med. Microbiol.* **44**:444–452.
43. Marklund, B. I., J. M. Tennent, E. Garcia, A. Hamers, M. Baga, F. Lindberg, W. Gaastra, and S. Normark. 1992. Horizontal gene transfer of the *Escherichia coli pap* and *prs* pili operons as a mechanism for the development of tissue-specific adhesive properties. *Mol. Microbiol.* **6**:2225–2242.
44. Maurelli, A. T., R. E. Fernandez, C. A. Bloch, C. K. Rode, and A. Fasano. 1998. “Black holes” and bacterial pathogenicity: a large genomic deletion that enhances the virulence of *Shigella* spp. and enteroinvasive *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **95**:3943–3948.
45. McDaniel, T. K., K. G. Jarvis, M. S. Donnenberg, and J. B. Kaper. 1995. A genetic locus of enterocyte effacement conserved among diverse enterobacterial pathogens. *Proc. Natl. Acad. Sci. USA* **92**:1664–1668.
46. Ochman, H., and R. K. Selander. 1984. Standard reference strains of *Escherichia coli* from natural populations. *J. Bacteriol.* **157**:690–693.
47. Oswald, E., P. Pohl, E. Jacquemin, P. Lintermans, K. Van Muylem, A. D. O'Brien, and J. Mainil. 1994. Specific DNA probes to detect *Escherichia coli* strains producing cytotoxic necrotising factor type 1 or type 2. *J. Med. Microbiol.* **40**:428–434.
48. Picard, B., J. S. Garcia, S. Gouriou, P. Duriez, N. Brahimi, E. Bingen, J. Elion, and E. Denamur. 1999. The link between phylogeny and virulence in *Escherichia coli* extraintestinal infection. *Infect. Immun.* **67**:546–553.
49. Rode, C. K., L. J. Melkerson-Watson, A. T. Johnson, and C. A. Bloch. 1999. Type-specific contributions to chromosome size differences in *Escherichia coli*. *Infect. Immun.* **67**:230–236.
50. Russo, T. A., and J. R. Johnson. 2000. Proposal for a new inclusive designation for extraintestinal pathogenic isolates of *Escherichia coli*: ExPEC. *J. Infect. Dis.* **181**:1753–1754.
51. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
52. Schmidt, H., L. Beutin, and H. Karch. 1995. Molecular analysis of the plasmid-encoded hemolysin of *Escherichia coli* O157:H7 strain EDL 933. *Infect. Immun.* **63**:1055–1061.
53. Schmidt, H., C. Geitz, P. I. Tarr, M. Frosch, and H. Karch. 1999. Non-O157:H7 pathogenic Shiga toxin-producing *Escherichia coli*: phenotypic and genetic profiling of virulence traits and evidence for clonality. *J. Infect. Dis.* **179**:115–123.
54. Schubert, S., A. Rakin, H. Karch, E. Carniel, and J. Heesemann. 1998. Prevalence of the “high-pathogenicity island” of *Yersinia* species among *Escherichia coli* strains that are pathogenic to humans. *Infect. Immun.* **66**:480–485.
55. Tarr, P. I., S. S. Bilge, J. C. Vary, Jr., S. Jelacic, R. L. Habeeb, T. R. Ward, M. R. Baylor, and T. E. Besser. 2000. *Iha*: a novel *Escherichia coli* O157:H7 adherence-conferring molecule encoded on a recently acquired chromosomal island of conserved structure. *Infect. Immun.* **68**:1400–1407.