

Letter to the Editor

Isolation and Characterization of a Nonfluorescent Strain of *Legionella parisiensis*

Legionella species are gram-negative environmental bacteria that replicate within amoebae and other protozoa. After transmission to humans, pneumonia, influenza-like illness, or asymptomatic infection may occur (1). So far, 50 *Legionella* spp. have been described (www.bacterio.cict.fr). About half of them are associated with illness in humans. The most important species is *L. pneumophila*. The remaining *Legionella* spp. account for about 5 to 10% of clinical cases, mostly in immunosuppressed patients (1).

Several *Legionella* spp. exhibit blue-white fluorescence under long-wavelength UV light. Some investigators have proposed placing these species in the genus *Fluoribacter* within the family *Legionellaceae*. However, recent studies using 16S rRNA gene analysis confirm that only one genus in this family, *Legionella*, exists. Nevertheless, the blue-white species formed clusters within this genus when 16S rRNA, *mip* (macrophage infectivity potentiator), and *rpoB* (RNA polymerase subunit beta) DNA sequences were analyzed (2, 5). The species within this cluster are *L. anisa*, *L. bozemanii*, *L. cherrii*, *L. dumoffii*, *L. gormanii*, *L. parisiensis*, *L. gratiana*, *L. steigerwaltii*, and *L. tucsoniensis*.

Legionella parisiensis was first isolated from environmental samples. In 1997, Lo Presti et al. (3) obtained the first strain of this species from a liver transplant patient with pneumonia. Here, we report on a nonfluorescent *L. parisiensis* strain that was isolated from an immunosuppressed patient with pneumonia in Germany.

This strain showed the typical ground glass colony appearance on buffered charcoal yeast extract (BCYE) agar, did not grow on sheep blood agar, and reacted with the non-*L. pneumophila* reagent in a commercially available latex agglutination kit (Oxoid, Wesel, Germany). According to the manufacturer, this reagent contains rabbit antisera against *L. anisa*, *L. bozemanii*, *L. dumoffii*, *L. gormanii*, *L. jordanis*, *L. longbeachae*, and *L. micdadei*. By using rabbit antisera against these species prepared in our laboratory, the strain reacted strongly to rabbit anti-*L. bozemanii* antiserum and weakly to anti-*L. anisa* antiserum. In contrast, none of the colonies isolated exhibited the blue-white fluorescence under UV light (Fig. 1) typical for strains of the blue-white fluorescent group.

To define the species of this strain exactly, we determined the DNA sequences of the 16S rRNA, *mip*, and *rpoB* genes (2, 5). Comparison of the nucleotide sequences with those in the National Center for Biotechnology Information database by using the standard nucleotide Basic Local Alignment Search Tool (BLASTn) revealed nearly complete homology to *L. parisiensis*: 16S rRNA, 100%; *mip*, 99%; and *rpoB*, 100%. Based on *mip* sequences, the next related species were *L. tucsoniensis* (95%), *L. bozemanii* (95%), and *L. anisa* (94%). The *rpoB* sequences revealed *L. bozemanii* (94%) and *L. anisa* (90%) as the closest related species. The BLAST analysis of the 16S rRNA sequence shows agreement with *L. anisa* (97%), *L. bozemanii* (97%), and *L. dumoffii* (97%).

Furthermore, analysis of bacterial cell wall fatty acids was performed by using the Microbial Identification System (MIDI, Newark, Del.) with the Hewlett-Packard 9890 chro-

matograph and the MIDI software package Aerobe Clin40 database, version 4.0. For this analysis, three colonies were grown and analyzed in separate runs. Good library comparisons—i.e., a similarity index (SI) of >0.6 and separation from the second choice of >0.1—were obtained for *L. parisiensis* (SI = 0.791 ± 0.04). Similarities were indicated for *L. gormanii* (SI = 0.588), *L. anisa* (SI = 0.587), and *L. bozemanii* (SI = 0.326). Thus, the strain could be unambiguously assigned to the species *L. parisiensis*.

There is no information available describing which genes are involved in the blue-white fluorescent phenotype. Furthermore, the relevance of these genes for virulence is unknown. Originally *L. parisiensis* was not known to cause pneumonia in humans. In 1996, O'Connell et al. (4) showed that the type strain of *L. parisiensis* is able to multiply in macrophage-like cells. We infected *Acanthamoeba castellanii* with our strain of *L. parisiensis*. Within 24 h, our strain multiplied approximately 1,000-fold. This result showed that the genes responsible for blue-white fluorescence seem to be not essential for intracellular multiplication. The fact that we isolated this nonfluorescent *L. parisiensis* strain from a patient argues for some virulence properties of this strain.

In general, the identification of *Legionella* spp. in the clinical laboratory is based on phenotypic properties: i.e., colony morphology, cysteine dependence, and serotyping. But, as shown here, this might be misleading. Due to the serological cross-reactivity among *Legionella* spp., definitive species identifica-




FIG. 1. Growth of *L. parisiensis* (colonies 1 to 6) and *L. bozemanii* (colonies 6 and 7) on BCYE agar after 3 days at 37°C in 5% CO₂ when examined under long-wavelength UV light.

tion must be based on sequence analysis of *Legionella* genes. Thus, it might be possible that in different laboratories, strains exist that have been misclassified on the basis of serotyping results. If our strain had shown the blue-white phenotype, we would not have determined the exact species by sequencing the *mip* gene. Finally, it must be pointed out that all *Legionella* spp. are susceptible to macrolides, quinolones, or rifampin. Therefore, the choices of antibiotic therapy are identical for all *Legionella* spp.

Nucleotide sequence accession number. The nucleotide sequences have been submitted to GenBank EMBL under accession no. AJ601373, AJ601374, and AJ601375 for the *mip*, *rpoB*, and 16S rRNA genes, respectively.

We are grateful to Jutta Paasche for technical assistance.

This study was supported by the Federal Ministry of Education and Research of Germany, Network of Competence in Medicine CAPNetz.

REFERENCES

1. **Fields, B. S., R. F. Benson, and R. E. Besser.** 2002. *Legionella* and Legionnaires' disease: 25 years of investigation. *Clin. Microbiol. Rev.* **15**:506–526.
2. **Ko, K. S., H. K. Lee, M.-Y. Park, K.-H. Lee, Y.-J. Yun, S.-Y. Woo, H. Miyamoto, and Y.-H. Kook.** 2002. Application of RNA polymerase β -subunit gene (*rpoB*) sequences for the molecular differentiation of *Legionella* species. *J. Clin. Microbiol.* **40**:2653–2658.
3. **Lo Presti, F., S. Riffard, F. Vandenesch, M. Reyrolle, E. Ronco, P. Ichai, and J. Etienne.** 1997. The first clinical isolate of *Legionella parisiensis*, from a liver transplant patient with pneumonia. *J. Clin. Microbiol.* **35**:1706–1709.
4. **O'Connell, W. A., L. Dhand, and N. P. Cianciotto.** 1996. Infection of macrophage-like cells by *Legionella* species that have not been associated with disease. *Infect. Immun.* **64**:4381–4384.
5. **Ratcliff, R. M., J. A. Lanser, P. A. Manning, and M. W. Heuzenroeder.** 1998. Sequence-based classification scheme for the genus *Legionella* targeting the *mip* gene. *J. Clin. Microbiol.* **36**:1560–1567.

Liane Igel
Jürgen Herbert Helbig
Paul Christian Lück*
Institute of Clinical Microbiology and Hygiene
Fiedlerstr. 42
01307 Dresden, Germany

*Phone: 49 0351 4586580
 E-mail: Christian.Lueck@mailbox.tu-dresden.de