

Use of Single Nucleotide Polymorphisms in the *plcR* Gene for Specific Identification of *Bacillus anthracis*

W. Ryan Easterday,¹† Matthew N. Van Ert,¹† Tatum S. Simonson,¹ David M. Wagner,¹
Leo J. Kenefic,¹ Christopher J. Allender,¹ and Paul Keim^{1,2*}

Department of Biological Sciences, Northern Arizona University, Flagstaff, AZ 86011-5640,¹ and The Translational Genomics Research Institute (TGen), 445 N. Fifth Street, Phoenix, AZ 85004²

Received 7 September 2004/Returned for modification 13 October 2004/Accepted 27 October 2004

A TaqMan-minor groove binding assay designed around a nonsense mutation in the *plcR* gene was used to genotype *Bacillus anthracis*, *B. cereus*, and *B. thuringiensis* isolates. The assay differentiated *B. anthracis* from these genetic near-neighbors and determined that the nonsense mutation is ubiquitous across 89 globally and genetically diverse *B. anthracis* strains.

The genetic similarities among the pathogenic, spore-forming soil bacteria *Bacillus cereus*, *B. thuringiensis*, and *B. anthracis* have resulted in the suggestion that they be considered members of the same species (3). Interestingly, these bacteria exhibit phenotypic differences and express virulence in diverse ways. *B. cereus* and *B. thuringiensis* are opportunistic pathogens in mammals due to the secretion of nonspecific virulence factors, such as hemolysins, the expression of which is regulated by the transcriptional activator PlcR (8). In *B. anthracis*, PlcR is inactivated due to a nonsense mutation in the *plcR* gene (1), and its virulence in mammals is attributed to the expression of specific toxins under the control of the AtxA regulator (2).

The nonsense mutation in the *plcR* gene of *B. anthracis* may represent an evolutionarily stable, species-specific marker. Research by Mignot et al. (8), in which a functional PlcR was expressed in *B. anthracis*, demonstrated that PlcR- and AtxA-controlled regulons were incompatible, as *plcR* expression interfered with sporulation in *B. anthracis*. Since sporulation is a critical component of the ecology of *B. anthracis*, the authors speculated that a functional PlcR is counterselected in this species. Recent sequence comparisons of the *plcR* genes of two phylogenetically distinct *B. anthracis* lineages revealed the same nonsense mutation in the *plcR* gene (9), providing additional evidence to support the species specificity of this mutation.

To initially test the utility of the nonsense mutation in *plcR* as a species-specific marker for *B. anthracis*, we examined the *plcR* gene fragments that surround the nonsense mutation in several *Bacillus* spp. The strains examined included nine genetically diverse *B. anthracis* strains, nine *B. cereus* strains, six *B. thuringiensis* strains, and one unidentified near-neighbor (TET 2b-3) (4). Sequences obtained either from GenBank or from sequencing efforts in our laboratory were compared using MegAlign (Fig. 1). The nonsense mutation was present in all nine of the *B. anthracis* sequences and was absent in the 16 near-neighbor sequences.

Based upon these sequences, we designed a TaqMan-minor groove binding (MGB) allelic discrimination assay around the nonsense mutation. The TaqMan-MGB probes were designed using Primer Express software (Applied Biosystems, Foster City, CA). One probe was designed to specifically hybridize to the *B. anthracis* sequence (5'-VIC-CAAAGCGCTTATTCG TATT-3'-MGB), and the other was designed to hybridize to the alternate allele (5'-FAM-AAAGCGCTTCTTCGTATT-3'-MGB) (Fig. 1 shows probe locations). Real-time PCRs were conducted in 10.0- μ l reaction mixtures that contained 600 nM of both forward (5'-CCAATCAATGTCATACTATTAATTT GACAC-3') and reverse (5'-ATGCAAAAAGCATTATACTT GGACAAT-3') primers (Fig. 1 shows primer locations), 250 nM of each probe, 1 \times Invitrogen Platinum qPCR SuperMix-UDG, and 1.0 μ l of template. Thermal cycling was performed on an ABI 7900 HT sequence detection system (Applied Biosystems) under the following conditions: 50°C for 2 min, 95°C for 2 min, and 40 cycles of 95°C for 15 s and 60°C for 1 min.

To further evaluate the nonsense mutation in *plcR* as a species-specific marker for *B. anthracis*, we used the assay described above to genotype a collection of *B. anthracis* strains representing 89 unique genetic lineages (6). In addition, we genotyped 29 strains that were identified by amplified fragment length polymorphism analysis as genetic near-neighbors of *B. anthracis* (4) (Table 1 shows strain list). All of the *B. anthracis* isolates supported amplification and were shown to have the *plcR* nonsense mutation genotype (T allele). Not surprisingly, genetic near-neighbors that had mutations in the priming site either failed to exhibit amplification or amplified with lower efficiency relative to the four strains that had complete sequence identity to *B. anthracis* except for the nonsense mutation (Table 1; Fig. 1). Of the 29 near-neighbors, 16 failed to exhibit amplification and the remaining 13 exhibited the G allele genotype (Table 1). The presence of the G allele in 5 of the 16 isolates that failed to amplify in the assay was confirmed via sequencing with flanking primers (Fig. 1).

To test the limit of detection of the assay, we utilized a dilution series generated from DNAs from three diverse *B. anthracis* isolates (Ames [A0462], Kruger B1 [A0442], and Voluum [A0488]). DNA was quantified using a Pico Green assay, and template levels ranging from 100 pg to 10.0 fg were used

* Corresponding author. Mailing address: Department of Biological Sciences, Northern Arizona University, Flagstaff, AZ 86011-5640. Phone: (928) 523-1078. Fax: (928) 523-0639. E-mail: Paul.Keim@uau.edu.

† W.R.E. and M.N.V. contributed equally to the research described in this paper.




FIG. 1. Sequence alignment of *plcR* gene fragments from *B. anthracis* and genetic near-neighbors. The numbered lines indicate the sequences of (line 1) the primers and probes used in the assay; (line 2) *B. anthracis* strains Ames, Vollum, A2012, A1055, AUS94, CNEVA9066, Kruger, Sterne, and WNA6153; (line 3) *B. cereus* strains 3A (GenBank accession no. AY785766) and S2-8 and *B. thuringiensis* strain HD1011; (line 4) *B. thuringiensis* strain 97-27 (AY785771); (line 5) near-neighbor strain TET 2b-3; (line 6) *B. cereus* strain AH-527 (AY785767); (line 7) *B. cereus* strain D17 (AY785768) and *B. thuringiensis* strains HD682, HD571, and HD44; (line 8) *B. cereus* strains F3502/72 (AY785769) and R6; (line 9) *B. cereus* strain F2-1 (AY785770); and (line 10) *B. cereus* strains R4 and ATCC 33018 and *B. thuringiensis* strain HD1012 (AY785772). Light shading indicates areas of polymorphism that are detected in the assay. Darker shading indicates nucleotide differences between near-neighbors and *B. anthracis*. * indicates the nonsense mutation.

in the *plcR* TaqMan assay. The assay reliably detected and genotyped *Bacillus anthracis* DNA template at levels as low as 100 fg, with 10-fg samples exhibiting sporadic amplification (Fig. 2).

TABLE 1. List of *Bacillus* sp. strains examined using the assay developed in this study

Species ^{a,b}	Strain ^b	<i>plcR</i> gene fragment sequence ^d	Avg threshold cycle ^e	TaqMan result (allele)
BA	89 diverse strains ^c	2	26.0 ^f	+ (T)
BC	ATCC 4342	NA	No Amp	-
BC	ATCC 14579	NA	No Amp	-
BC	D17	7	No Amp	- (G) ^g
BC	F3-27	NA	No Amp	-
BC	F3502/72	8	27.1	+ (G)
BC	R6	8	27.6	+ (G)
BC	ATCC 33018	10	38.5	+ (G)
BC	D5	NA	28.4	+ (G)
BC	3A	3	24.8	+ (G)
BC	S2-8	3	27.0	+ (G)
BC	F3350/87	NA	No Amp	-
BC	S2-4	NA	34.9	+ (G)
BC	R4	10	34.7	+ (G)
BC	F2-1	9	No Amp	- (G) ^g
BC	AH 527	6	30.0	+ (G)
BT	HD 1015	NA	No Amp	-
BT	HD 681	NA	No Amp	-
BT	HD 288	NA	No Amp	-
BT	HD 526	NA	No Amp	-
BT	97-27	4	33.0	+ (G)
BT	HD 1011	3	26.7	+ (G)
BT	HD 571	7	No Amp	- (G) ^g
BT	HD 682	7	No Amp	- (G) ^g
BT	HD 974	NA	No Amp	-
BT	HD 44	7	No Amp	- (G) ^g
BT	HD 30	NA	No Amp	-
BT	HD 1012	10	33.0	+ (G)
BT	HD 50	NA	No Amp	-
UNK	TET-2B	5	33.3	+ (G)

^a BA, *B. anthracis*; BC, *B. cereus*; BT, *B. thuringiensis*; UNK, unknown *Bacillus* spp.

^b Species and strain designations according to reference 4.

^c The 89 diverse *B. anthracis* strains are described in reference 6.

^d As represented in Fig 1. NA, strain not sequenced.

^e Input, 10 pg, average of triplicate cycle threshold values. No Amp, no amplification.

^f Average of triplicate cycle threshold values from 10-pg input of *B. anthracis* Ames strain.

^g *plcR* genotypes (Fig. 1) were determined via sequencing using flanking primers.

Our data provide further evidence that the nonsense mutation in the *plcR* gene of *B. anthracis* is an evolutionarily stable, species-specific marker. Although additional genetic changes, such as deletions, could produce a nonfunctional PlcR in *B. anthracis* and potentially cause false-negative results in our assay, this was not observed. The presence of this mutation in the 89 genetically diverse *B. anthracis* lineages examined here, as well as the known genetic homogeneity of the species (5), limits the likelihood of alternate genetic mechanisms for *plcR* inactivation in *B. anthracis*. The recent findings of Slamti et al. (9), which demonstrated that this specific *plcR* nonsense




FIG. 2. Results of triplicate analysis of 10-fold serial dilutions of DNA from *B. anthracis* strain A0442. The average cycle threshold values across the three replicates were as follows: 100 pg, 21.4; 10 pg, 24.7; 1 pg, 28.2; 100 fg, 31.2; 10 fg, 34.3. The average cycle threshold values for strains A0488 and A0462 were similar (data not shown), although amplification at the 10-fg level was not consistent.

mutation was not responsible for the nonhemolytic properties of *B. cereus* and *B. thuringiensis* strains, further support the concept that this nonsense mutation is a defining or canonical single nucleotide polymorphism (7) for *B. anthracis*.

The real-time assay presented here represents a potentially valuable diagnostic tool in the event of a future bioterrorist attack. From a biodefense perspective, diagnostic assays allowing rapid and specific identification of *B. anthracis* are critical to initiate appropriate first-response actions, such as remediation measures and prophylactic therapies. As our assay targets a well-characterized, biologically relevant single nucleotide polymorphism, it limits the likelihood of false-negative or -positive results, which can lead to misallocation of resources during an attack scenario. Furthermore, this assay is amenable to high-throughput real-time PCR platforms that are currently used in homeland defense initiatives, such as BioWatch.

In summary, our results indicate that the *plcR* nonsense mutation is ubiquitous in globally and genetically diverse *B. anthracis* isolates and, thereby, represents an excellent target for diagnostic assays. Future studies will involve genotyping more extensive collections of *B. anthracis* and genetic near-neighbors, as well as the optimization and validation of this assay for the specific, low-level detection of *B. anthracis* in complex environmental samples.

This work was supported by grants from the National Institutes of Health, the Department of Homeland Security, and the Cowden Endowment at Northern Arizona University.

We thank Jeff Henrickson for his technical assistance; Jason Farlow for his thoughtful review of the manuscript; and Paul Jackson, Karen

Hill, and their colleagues at Los Alamos National Laboratory for providing the near-neighbor strains.

REFERENCES

1. Agaisse, H., M. Gominet, O. A. Økstad, A. Kolstø, and D. Lereclus. 1999. PlcR is a pleiotropic regulator of extracellular virulence factor gene expression in *Bacillus thuringiensis*. *Mol. Microbiol.* **32**:1043–1053.
2. Guignot, J., M. Mock, and A. Fouet. 1997. AtxA activates the transcription of genes harbored by both *Bacillus anthracis* virulence plasmids. *FEMS Microbiol. Lett.* **147**:203–207.
3. Helgason, E., O. A. Økstad, D. A. Caugant, H. A. Johansen, A. Fouet, M. Mock, I. Hegna, and A. Kolstø. 2000. *Bacillus anthracis*, *Bacillus cereus*, and *Bacillus thuringiensis*—one species on the basis of genetic evidence. *Appl. Environ. Microbiol.* **66**:2627–2630.
4. Hill, K. K., L. O. Ticknor, R. T. Okinaka, M. Asay, H. Blair, K. A. Bliss, M. Laker, P. E. Pardington, A. P. Richardson, M. Tonks, D. J. Beecher, J. D. Kemp, A. Kolstø, A. C. L. Wong, P. Keim, and P. J. Jackson. 2004. Fluorescent amplified fragment length polymorphism analysis of *Bacillus anthracis*, *Bacillus cereus*, and *Bacillus thuringiensis* isolates. *Appl. Environ. Microbiol.* **70**:1068–1080.
5. Keim, P., A. Kalif, J. Schupp, K. Hill, S. E. Travis, K. Richmond, D. M. Adair, M. Hugh-Jones, C. Kuske, and P. Jackson. 1997. Molecular evolution and diversity in *Bacillus anthracis* as detected by amplified fragment length polymorphism markers. *J. Bacteriol.* **179**:818–824.
6. Keim, P., L. B. Price, A. M. Klevytska, K. L. Smith, J. M. Schupp, R. Okinaka, P. J. Jackson, and M. E. Hugh-Jones. 2000. Multiple-locus variable-number tandem repeat analysis reveals genetic relationships within *Bacillus anthracis*. *J. Bacteriol.* **182**:2928–2936.
7. Keim, P., M. N. Van Ert, T. Pearson, A. J. Vogler, L. Y. Huynh, and D. M. Wagner. 2004. Anthrax molecular epidemiology and forensics: using the appropriate marker for different evolutionary scales. *Infect. Genet. Evol.* **4**:205–213.
8. Mignot, T., M. Mock, D. Robichon, A. Landier, D. Lereclus, and A. Fouet. 2001. The incompatibility between the PlcR- and AtxA-controlled regulons may have selected a nonsense mutation in *Bacillus anthracis*. *Mol. Microbiol.* **42**:1189–1198.
9. Slamti, L., S. Perchat, M. Gominet, G. Vilas-Bóas, A. Fouet, M. Mock, V. Sanchis, J. Chaufaux, M. Gohar, and D. Lereclus. 2004. Distinct mutations in PlcR explain why some strains of the *Bacillus cereus* group are nonhemolytic. *J. Bacteriol.* **186**:3531–3538.