

## Predominance of Clones Carrying Panton-Valentine Leukocidin Genes among Methicillin-Resistant *Staphylococcus aureus* Strains Isolated in Japanese Hospitals from 1979 to 1985<sup>∇</sup>

Xiao Xue Ma, Teruyo Ito,\* Piriyaoporn Chongtrakool, and Keiichi Hiramatsu

*Juntendo University, Department of Bacteriology, Tokyo, Japan*

Received 11 May 2006/Returned for modification 3 July 2006/Accepted 5 October 2006

We examined 97 strains of methicillin-resistant *Staphylococcus aureus* (MRSA) isolated between 1979 and 1985, the period of time when the appearance of MRSA strains increased, and we determined that these strains are distinct from the MRSA clones predominating in today's Japanese hospitals. Type IV staphylococcal cassette chromosome *mec* (SCC*mec*) strains were the most frequent, comprising 53.6% of all strains, followed by type I (22.7%) and type II (21.6%) SCC*mec* strains. Among the type IV SCC*mec* strains, the frequencies of two new subtypes, type IV.3 (IVc) and type IV.4 (IVd), were very high, comprising 38.1 and 10.3% of all strains, respectively. Forty-four of the 97 strains (45.3%) were Panton-Valentine leukocidin (PVL) positive. Among the PVL-positive strains, sequence type 30 (ST30)-SCC*mec* type IV strains producing type 4 coagulase were the most frequent. This is in striking contrast to the MRSA strains isolated in the 1990s, most of which were ST5-SCC*mec* type II strains producing type 2 coagulase and positive for the toxic shock syndrome toxin 1 gene. We also identified a new PVL-carrying prophage lysogenized in a type IV.3 SCC*mec* strain, 81/108.  $\phi$ 108PVL was distinct from the three extant PVL-carrying phages and was presumed to be carried by ST30-type IV.3 SCC*mec* strains isolated in Japan. These results provide genetic bases for the transition of MRSA clones in Japan that is commonly considered the transition from coagulase type 4 MRSA strains to coagulase type 2 MRSA strains. The results also suggested that MRSA strains that predominated between 1979 and 1985 were generated from PVL-positive methicillin-susceptible *S. aureus* strains through the integration of SCC*mec* elements.

Methicillin-resistant *Staphylococcus aureus* (MRSA) is one of the major human pathogens and causes a wide range of infections in health care settings and community environments (5, 11, 36). The first MRSA strain was reported in England in 1961, after which it appeared worldwide (25). In Japan, *S. aureus* strains exhibiting low-level resistance to methicillin were reported in the early 1960s, albeit at very low frequencies of less than 3% (20, 35). However, from the late 1970s to the early 1980s, a period when third-generation cephalosporins with a wide spectrum of antimicrobial activity were introduced in the clinical field, MRSA strains began to dominate in Japanese hospitals (8). Since some of the third-generation cephalosporins had weak antibacterial activities toward *S. aureus*, it was suspected that the excessive use of these antibiotics was responsible for the expansion of MRSA strains in hospitals throughout Japan during this time. The frequency of isolation of MRSA strains increased drastically in the early 1980s from 34% (from 1982 to 1983) to 43% (from 1986 to 1987) and from 18.6% (from July to December 1981) to 35.9% (from January to June 1982) (34, 38, 43). From the mid-1980s to the early 1990s,  $\beta$ -lactam antibiotics with improved antibacterial activities towards *S. aureus* were introduced, but by 1990, these antibiotics had become less effective against clinical isolates.

MRSA strains have been characterized by susceptibility testing and coagulase isotyping. Coagulase typing is a method developed in Japan which can classify *S. aureus* strains based on the antigenic difference in coagulase, the protein causing coagulation of plasma (47). In the early 1980s, coagulase type 4 MRSA strains, which showed a heterogeneous profile of oxacillin resistance, disseminated in Japanese hospitals. But from the mid 1980s to 1990s, the majority of hospital MRSA isolates were coagulase type 2 MRSA strains, most of which exhibit high resistance to oxacillin as well as to other many antibiotics. It was suggested that MRSA strains disseminated in Japanese hospitals had undergone a transition (9, 29, 31, 46). We have investigated the genotypes by conducting ribotyping and the carriage of toxin genes and reported that a shift of MRSA strains from coagulase type 4 MRSA strains to coagulase type 2 MRSA strains should be regarded as a shift of MRSA clones. The coagulase type 2 MRSA strains, which were characterized by the production of enterotoxin A and by their specific ribotyping patterns, were prevalent in the early 1980s but declined drastically in the 1990s. On the other hand, coagulase type 4 MRSA strains were characterized by the production of enterotoxin C and toxic shock syndrome toxin 1 (TSST-1) and by having their specific ribotyping patterns (16, 46).

We now know that MRSA strains have evolved from methicillin-susceptible *Staphylococcus aureus* (MSSA) by the acquisition of the staphylococcal cassette chromosome *mec* (SCC*mec*) carrying the *mecA* gene (24, 30). There are several types of SCC*mec* elements that differ in their genomic organi-

\* Corresponding author. Mailing address: Juntendo University, Department of Bacteriology, Hongo 1-1-1 Bunkyo-ku, Tokyo 113-8421, Japan. Phone: 81-3-5802-1041. Fax: 81-3-5684-7830. E-mail: teruybacc@med.juntendo.ac.jp.

<sup>∇</sup> Published ahead of print on 18 October 2006.

TABLE 1. Primers used in this study

Primer purpose	Primer name	Nucleotide sequence (5'-3')	Reference or location on $\phi$ 108PVL
Subtyping of type IV SCC <sub>mec</sub> elements			
IV.1 (IVa)	4a1	TTTGAATGCCCTCCATGAATAAAAT	Okuma et al. (39)
	4a2	AGAAAAGATAGAAGTTCGAAAGA	Okuma et al. (39)
IV.2 (IVb)	4b1	AGTACATTTTATCTTTGCGTA	Okuma et al. (39)
	4b2	AGTCATCTTCAATATCGAGAAAGTA	Okuma et al. (39)
IV.3 (IVc)	4c1	TCTATTCAATCGTTCTCGTATTT	Ma et al. (32)
	4c2	TCGTTGTCATTTAATTCTGAACT	Ma et al. (32)
IV.4 (IVd)	4d1	TTTGAGAGTCCGTCATTATTTCTT	Ma et al. (32)
	4d2	AGAATGTGGTTATAAGATAGCTA	Ma et al. (32)
Identification of virulence factor			
LukS and LukF	PVL-F	ATGTCTGGACATGATCCAA	Ma et al. (32)
	PVL-R	AACTATCTCTGCCATATGGT	Ma et al. (32)
TSST-1	TSST-1A	TGATATGTGGATCCGTCAT	Ma et al. (32)
	TSST-1B	AAACACAGATGGCAGCAT	Ma et al. (32)
CNA	cna-1	ACACCAGACGGTGCAACAATTA	Ma et al. (32)
	cna-2	AGCAATACCGTTTGCATCTGTTA	Ma et al. (32)
SEH	entH-F	ATTCACATCATATGCGAAAGCAG	Ma et al. (32)
	entH-R	ATGTCGAATGAGTAATCTCTAG	Ma et al. (32)
Amplification of the whole genome of $\phi$ 108PVL			
From chromosome to integrase	phiMW-DN	GCAGAAAAAGATGCGATTGAA	
	int-R	CGGATACAAAGACACTACAAA	800–780
From integrase to antirepressor	int-F	TTTGTAGTGTCTTTGTATCCG	780–800
	anti-R	CCGAATGTGTGATTGATGTTT	5229–5209
From antirepressor to terminase large subunit	anti-F	ATTGTATTTGCAGATGCAGTAG	5001–5022
	termi-R	TAAATACATCTTCAATGTCTGTC	19338–19316
From terminase large subunit to portal protein	termi-F	AAACAAGGTAAGTCTCTAATCG	19116–19137
	portal-R	TCTAAATTAGCATCCGTGATAC	21616–21637
From portal protein to tail protein	portal-F	ACACGTGATAAAACAGGAGAA	21069–21089
	tail-R	TTCGACTTTTATCTTCGACTTTTC	27382–27359
From tail protein to LukS-PV	tail-F	TAAAAAGACATGCAAAGAGAGC	27566–27587
	LukS-R	CAGCTTTAGATTCATGAAACGAA	42344–42322
From LukS-PV to chromosome	LukS-F	TGGTCAACTATATCGTGGTTTT	42038–42059
	phiMW-UP	TCGCCACGTTTAGCAATTTTAT	
Amplification of the leftmost region specific to three extant phages in combination with a primer, int-c			
	int-c	TTTGTAGTGTCTTTGTATCCG	
$\phi$ PVL	rec $\phi$ PVL	TCGAAACCTTATCAAAAAGAAACT	
$\phi$ SLT	rec $\phi$ SLT	AAGCTACTGTACAGCGTTATG	
$\phi$ Sa2mw	rec $\phi$ Sa2mw	AATCTACCTTATCCCAGATACGA	
Amplification of the rightmost region specific to three extant phages in combination with LukSR listed above			
$\phi$ PVL	$\phi$ PVL-RE	TCTTAACTTACCTATTAGAACG	
$\phi$ SLT, $\phi$ Sa2mw	$\phi$ SLT-MW-RE	ATTACAAGCGTATGACTTATCGA	
Amplification of the regions located on $\phi$ 108PVL			
A	int-F2	ATGTTTTTCGAGTTTTTGTAGTTAG	393–415
	int-R2	CATTTTAAATTGCCAGCATCTTA	929–950
B	termi-2	GTAGAGGTCAATACAGAATGTT	19066–19087
	portal-R2	AATCTTATGATACGATGTCCC	19743–19763
C	portal-F2	ACACGTGATAAAACAGGAGAA	21069–21089
	portal-R	TCTAAATTAGCATCCGTGATAC	21616–21637
D	tail-F2	CGTTTGTGGGATAAACTACGTAAT	29952–29975
	tail-R2	TAGTAGTACCGAAATGAAATCTC	30386–30408
E	tail-F3	CTTGATTAGACTCAACCAAACCT	31232–31257
	tail-R3	ATAATTGGGATAGCAACGCAA	31704–31725
F	108RE-F	AGGCTTCACCCCTTACTTATTGA	42038–42059
	108RE-R	GATACCTGATAGCGAATTCG	44363–44832




FIG. 1. Structural comparison of different subtypes of type IV SCCmec elements. The structures of four subtypes of type IV SCCmec elements are illustrated based on the nucleotide sequences deposited in the DDBJ, EMBL, and GenBank databases under accession no. AB063172 (type IV.1 SCCmec), AB063173 (type IV.2 SCCmec), AB096217 (type IV.3 SCCmec), and AB097677 (IV.4 SCCmec). The SCCmec element is composed of two essential gene complexes, the *ccr* gene complex (pink) and the *mec* gene complex (light blue). The *ccr* gene complex consists of *ccr* genes which are responsible for the mobility of the SCCmec element and surrounding ORFs. The *mec* gene complex is responsible for methicillin-cephem resistance. Other areas (light gray) of SCCmec are nonessential and are divided into three regions, J1 to J3. Direct repeats containing integration site sequences for SCCmec elements are indicated by red arrowheads. The locations of the primer sets used for PCR amplification of an entire SCCmec element of strain 81/108 and a part of SCCmec element of strain JCSC4469 are indicated by arrows.

zations and structures (22, 23, 33, 40, 44), and it is generally accepted that MRSA clones could be defined by SCCmec type and genotype. The classification of MRSA clones is important for epidemiological study to distinguish MRSA clones that have predominated through a hospital setting and a community setting.

In the current study, we conducted a retrospective analysis of MRSA strains isolated from Japanese hospitals between 1979 and 1985 and during the early 1990s using several different molecular typing methods: SCCmec element typing, virulence gene repertoire, and multilocus sequence typing (MLST) (13). Our results provided genetic proof for the transition of MRSA clones predominating in Japanese hospitals. Furthermore, we found that approximately one-half of MRSA strains disseminated in the early 1980s in Japan were Pantone-Valentine leukocidin (PVL)-positive clones, possibly harboring a novel PVL-carrying phage.

**MATERIALS AND METHODS**

**MRSA strains used in this study.** MRSA strains isolated during three different time periods were tested: 97 MRSA strains isolated from 1979 to 1985 in Japanese hospitals (24 strains isolated at Tokyo University Hospital in 1982, 49 strains isolated at Gunma University Hospital from 1981 to 1985, 22 strains isolated at Jikei University Hospital from 1979 to 1981, and two strains isolated at other hospitals in 1981), 22 strains isolated in 1992 at Tokyo University Hospital, and 138 strains isolated in 1999 in 14 Japanese hospitals (Teikyo University, Fuku-

shima Prefectural Medical College, Iwate Medical College, Kobe University, Akita University, Saga Medical College, Tokyo University, Hiroshima University, Fukuoka University, Mie University, Kurume Medical College, Shiga University, Jikei Medical College, and National Defense Medical College).

**Susceptibility testing.** MICs were determined using the agar dilution method according to the procedure recommended by the Clinical and Laboratory Standards Institute. The antibiotics tested were oxacillin, tetracycline, erythromycin, and gentamicin (Sigma Chemical Co., St. Louis, Mo.), ceftizoxime (Fujiwara Pharmacy Co., Osaka, Japan), imipenem (Banyu Pharmaceutical Co., Tokyo, Japan), and levofloxacin (Daiichi Pharmaceutical Co., Tokyo, Japan).

**SCCmec typing and identification of virulence genes.** SCCmec typing was performed using PCR as described previously (21, 22, 39). The presence of four virulence-related genes, *lukS-PV-lukF-PV*, *tsst-1*, *seh*, and *cna*, was investigated by PCR using the primers listed in Table 1.

**Nucleotide sequencing of type IV.3 and type IV.4 SCCmec elements.** Several DNA fragments spanning the entire nucleotide sequence of the SCCmec element and an SCC-like element of strain 81/108 were amplified by long-range PCR with the five primer sets (Fig. 1): the region spanning *orfX* (an open reading frame [ORF] that works as a part of the gene cassettes SCC; all SCC elements are integrated into the 3' end of *orfX* or into the regions having the specific nucleotide sequence that is very similar to the nucleotide sequence of the 3' end of *orfX*) to *mecA* was amplified using primers cR1 and mA1; the region spanning *mecA* to IS1272 was amplified using primers mA2 and IS4; the region spanning *mecR1* to the *ccr* complex was amplified using primers mcR6 and α6; the region spanning the *ccr* complex to IE25923 was amplified using primers α5 and IE2 (5'-TCCACAAAATTACATATACTCTCT-3'); the region spanning IE25923 to left-flanking chromosomal region flanked to left end of SCCmec was amplified by PCR with primers IE1 (5'-AGAAATTTGTAGCGAATGATGA-3') and cLt2. The DNA fragments encompassing the region from the J1 region to the *mec* gene complex of the SCCmec element of strain JCSC4469 were amplified by




FIG. 2. (A) Essential structural and functional components of  $\phi 108PVL$  element are illustrated. Black arrowheads indicate the locations of primers used to amplify the entire  $\phi 108PVL$  genome. The two red arrowheads flanking the core sequence indicate the *att* sites on the phage element. (B) The locations of the primers used to amplify regions specific to  $\phi PVL$ ,  $\phi SLT$ , and  $\phi Sa2MW$ . (C) The ORFs in and around the  $\phi 108PVL$  element are illustrated as arrows in six possible reading frames. The direction of the arrows indicates the transcriptional direction for each ORF. Color codes are as follows: dark green, ORFs (or the parts of ORFs) that are well conserved among all other three PVL-carrying phages,  $\phi PVL$ ,  $\phi SLT$ , and  $\phi Sa2MW$ ; red, ORFs that are highly homologous to  $\phi PVL$ , which is lysogenized in the *S. aureus* ATCC 49775 strain; yellow, ORFs that are highly homologous to  $\phi SLT$ ; blue, ORFs that are highly homologous to  $\phi Sa2mw$ ; orange, ORFs that are highly homologous to both  $\phi PVL$  and  $\phi SLT$ ; green, ORFs that are highly homologous to both  $\phi SLT$  and  $\phi Sa2mw$ ; white, ORFs that are unique in  $\phi 108PVL$ .

long-range PCR using the following three sets of primers, mA2 and IS4, mCR6 and  $\alpha 6$ , and  $\alpha 5$  and cL1, as indicated in Fig. 1. The PCR products were purified using the QIAquick PCR purification kit (QIAGEN, Hilden, Germany), and their nucleotide sequencing was carried out by fluorescence dideoxy chain termination chemistry using the BigDye Terminator version 1.1 cycle sequencing kit (Applied Biosystems, CA) and an ABI Prism 3100 genetic analyzer (Applied Biosystems, CA). PCR and long-range PCR were performed as described previously (24).

**PCR amplification and nucleotide sequencing of PVL-carrying prophage  $\phi 108PVL$ .** The DNA fragments encompassing the entire phage-specific region in the bacterial chromosome, along with the primers used to amplify them by long-range PCR, were as follows: the phage-flanking chromosomal region to integrase, primers  $\phi MW-DN$  and *intR*; integrase to anti-repressor, primers *int-F* and *anti-R*; anti-repressor to terminase large subunit, primers *anti-F* and *termi-R*; terminase large subunit to portal gene, primers *termi-F* and *portal-R*; portal gene to tail gene, primers, *portal-F* and *tail-R*; tail gene to the *lukS-PV* gene, primers *tail-F* and *LukS-R*; and *lukS-PV* gene to flanking chromosomal region, primers, *LukS-F* and  $\phi MW-UP$ .

The nucleotide sequences of the primers used in these studies are listed in Table 1, and their locations in the phage region of the genome are illustrated in Fig. 2. PCR products were purified using a QIAquick PCR purification kit (QIAGEN, Hilden, Germany), and their nucleotide sequences were determined by primer walking.

**Coagulase isotyping.** Coagulase type was determined by using an inhibition test for the coagulation of plasma with eight specific antisera (Denka Seiken, Niigata, Japan) according to the method of Ushioda et al. (47). Briefly, bacterial strains were grown overnight in brain heart infusion broth and culture supernatants were collected by centrifugation. An appropriately diluted 0.1-ml aliquot of the supernatant was mixed with 0.1 ml of a solution containing each antiserum and incubated at 37°C for 1 h. Diluted rabbit plasma (0.2 ml) was added to each tube, followed by incubation at 37°C for 1 h or more until coagulation of the plasma was observed by visual inspection. The serotype of coagulase produced by

a given strain was determined by the specificity of the antiserum which inhibited coagulation.

**MLST.** Genotypes of representative strains were determined by MLST according to the procedure of Enright et al. (13). Alleles of the seven loci were assigned by a comparison of their sequences to the corresponding loci in the *S. aureus* MLST database ([www.mlst.net](http://www.mlst.net)). Sequence types were determined according to the combined pattern of the seven alleles, and clonal complexes were defined by the BURST (based upon related sequence types) program available on the MLST website.

**Nucleotide sequence accession numbers.** The entire nucleotide sequence of  $\phi 108PVL$  has been deposited in the DDBJ, EMBL, and GenBank databases under accession no. AB243556. The sequences of the type IV.3 (VIc) *SCCmec* of strain 81/108 and the type IV.4 (IVd) *SCCmec* of strain JCSC4469 have been deposited in the DDBJ, EMBL, and GenBank databases under accession no. AB096217 and AB097677, respectively.

## RESULTS

**SCCmec typing of MRSA strains isolated between 1979 and 1985 and the 1990s.** We characterized the *SCCmec* elements carried by 97 MRSA strains isolated between 1979 and 1985 (1979-1985 strains) and 22 MRSA strains isolated in 1992 and compared them to the types of *SCCmec* elements of MRSA strains isolated in 1999, which have previously been reported (6). As shown in Table 2, 95 of the 1979-1985 strains (97.9%) could be classified into one of three types of *SCCmec* elements, judging from the combinations of *mec* gene complex and *ccr* gene complex identified by PCR. Type IV *SCCmec* strains (53.6%) were the most frequent overall (53.6%), followed by



TABLE 2. Presence of virulence determinants in MRSA strains and genotypes of the strains representing each combination<sup>a</sup>

Group (yr of isolation) <sup>b</sup>	Characteristics of MRSA strains isolated at each period						Genotypes of chosen MRSA strains as indicated by:				No. of tested strains	
	SCCmec type	Ratio (%)	Presence of indicated virulent determinant				No. of strains with indicated results	Coagulase isotype	MLST			
			PVL	CNA	TSST-1	SEH			CC	ST		Allelic profile
A (1979-1985)	I (n = 22)	22.7	+	+	-	-	4	4	30	30	2, 2, 2, 2, 6, 3, 2	2
			+	-	-	-	7	4	30	30	2, 2, 2, 2, 6, 3, 2	1
			-	+	-	-	5	4	30	30	2, 2, 2, 2, 6, 3, 2	1
			-	-	-	-	6	4	30	30	2, 2, 2, 2, 6, 3, 2	1
	II.1 (IIa) (n = 21)	21.6	+	-	+	-	1	2	5	5	1, 4, 1, 4, 12, 1, 10	1
	IV.1 (IVa) (n = 1)	1.0	+	+	-	-	1	4	N <sup>c</sup>	N	1, 4, 1, 4, 12, 1, 10	6
	IV.3 (IVc) (n = 37)	38.1	+	+	-	-	30	4	30	30	2, 2, 2, 2, 6, 3, 2	7
			-	-	+	-	1	3	8	8	3, 3, 1, 1, 4, 4, 3	1
	IV.4 (IVd) (n = 10)	10.3	-	-	-	-	10	2	5	5	1, 4, 1, 4, 12, 1, 10	1
	IV.n (IVn) (n = 4)	4.1	+	+	-	-	1	4	30	30	2, 2, 2, 2, 6, 3, 2	1
-			+	-	-	3	4	30	30	2, 2, 2, 2, 6, 3, 2	1	
Nontypeable (n = 2)	1.0	-	-	-	-	1	2	5	5	1, 4, 1, 4, 12, 1, 10	1	
B (1992)	I (n = 1)	4.5	-	-	-	-	1					
	II.1 (IIa) (n = 20)	91.0	-	-	+	-	19	2	5	5	1, 4, 1, 4, 12, 1, 10	3
			-	-	-	-	1					
	IV.3 (IVc) (n = 1)	4.5	-	-	-	-	1					
C (1999)	I (n = 1)	0.7	-	-	-	-	1					
	II.1 (IIa) (n = 120)	91.3	-	-	+	-	114	2	5	5	1, 4, 1, 4, 12, 1, 10	6
			-	-	-	-	6					
	II.n (n = 6)	4.3	-	+	-	-	2					
	IV.3 (IVc) (n = 6)	4.3	-	+	-	+	4	7	1	81	1, 1, 1, 9, 1, 1, 1	2
			-	-	-	-	2					
Nontypeable (n = 5)	3.6	-	-	+	-	5						

<sup>a</sup> CNA, collagen adhesion protein; CC, clonal complex; N, nontypeable; SEH, staphylococcal enterotoxin H; ST, sequence type. +, present; -, absent.

<sup>b</sup> For group A (n = 97), the numbers of PVL-positive and TSST-1-positive strains were 44 (45.3) and 23 (23.7%), respectively, for group B (n = 22), the numbers of PVL-positive and TSST-1-positive strains were 0 and 19 (86.3%), respectively, and for group C (n = 138), the numbers of PVL-positive and TSST-1-positive strains were 0 and 123 (89.1%), respectively.

<sup>c</sup> The sequence type of the strains could not be assigned since the *gfpF* gene could not be amplified by PCR.

type I SCCmec strains (22.7%) and type II SCCmec strains (21.6%). The frequencies of each SCCmec strain, grouped according to hospital and ranked in order from highest to lowest, were as follows: Tokyo University Hospital, type IV, 17/24 (70.8%), type I, 6/24 (25.0%), and type II, 1/24 (4.2%); Gunma University Hospital, type IV, 32/49 (65.3%), type II, 8/49 (16.3%), type I, 7/49 (14.3%), and nontypeable, 2/49 (4.1%); and Jikei University Hospital, type II, 12/22 (54.5%), type I, 9/22 (4.1%), and type IV, 1/22 (4.5%). Although the frequency of each SCCmec strain at the three hospitals was different, it was clear that three types of SCCmec strains were disseminated and over three-quarters of the 1979-1985 strains (76%) carried either a type IV or a type I SCCmec element. In contrast, 126 out of 138 (93%) MRSA strains isolated in 1999 in 14 hospitals carried a type II SCCmec element. When we looked at the MRSA strains isolated in Tokyo University Hospital in particular, 91% of the strains isolated in 1992 and 100% of the strains isolated in 1999 carried a type II SCCmec element, indicating that type II SCCmec strains predominated in the early 1990s, at least at Tokyo University Hospital.

**New subtypes of type IV MRSA strains, IV.3 (IVc) and IV.4 (IVd), predominated in the early 1980s in Japan.** We further

classified the SCCmec type II and type IV elements based on the nucleotide sequences of their J1 regions. The majority of SCCmec type II strains were classified as subtype type II.1 (type IIa), while six strains isolated in 1999 could not be classified into subtype 1 (type IIa) or 2 (type IIb). When we examined the type IV SCCmec elements by PCR experiments using primers that would amplify type IV.1- and type IV.2-specific J1 sequences, we identified one SCCmec type IV.1 strain among 58 SCCmec type IV strains examined. This result suggested that most type IV strains carried SCCmec type IV of unknown subtype. Therefore, we amplified and sequenced several large DNA fragments spanning the entire SCCmec element from MRSA strain 81/108. The nucleotide sequence of the J1 region of the SCCmec element carried by 81/108 was not homologous to any previously reported type IV.1 or type IV.2 SCCmec elements, so we have designated this element type IV.3 SCCmec (Fig. 1). We then designed specific primers to amplify type IV.3-specific J1 region sequences and conducted PCR experiments using chromosomal DNA from strains whose subtypes, based on the sequence of the J1 region, were unclassifiable. According to this type of analysis, 40 of 55 SCCmec type IV strains were type IV.3 SCCmec strains and 15

TABLE 3. Comparison among MICs to 10 antibiotics of MRSA strains isolated in each period

Antibiotic	MIC ( $\mu\text{g/ml}$ ) of MRSA isolated in:			
	1979 to 1985		1999	
	50%	90%	50%	90%
Oxacillin	32	128	256	>512
Ceftizoxime	>512	>512	>512	>512
Imipenem	0.5	64	32	128
Ampicillin	64	256	32	64
Gentamicin	128	512	32	128
Tobramycin	128	512	256	512
Erythromycin	>512	>512	>512	>512
Tetracycline	2	4	64	64
Levofloxacin	0.25	0.5	8	64
Ciprofloxacin	1	2	NT <sup>a</sup>	NT

<sup>a</sup> NT, not tested.

strains still remained unclassifiable. We then amplified and sequenced DNA fragments from MRSA strain JCSC4469 that spanned the region of the SCCmec element from *mecA* to J1. The J1 region of the SCCmec element of strain JCSC4469 was not homologous to that of a type IV.1, IV.2, or IV.3 SCCmec element, so we have designated it type IV.4 (Fig. 1). We then carried out PCR experiments using primers that amplified type IV.4-specific J1 sequences and found that 10 of 15 previously unclassified strains belonged to this type, leaving 5 strains still nontypeable. These results indicated that type IV.3 SCCmec strains, which are infrequent among recent isolates, predominated in Japan between 1979 and 1985 (Table 2).

**Antibiotic susceptibilities.** We determined the MICs of eight antibiotics (oxacillin, imipenem, ampicillin, gentamicin, tobramycin, erythromycin, tetracycline, and levofloxacin) for MRSA strains isolated between 1979 and 1985 and compared them to MICs of MRSA strains isolated in 1992 and 1999 (Table 3). Most of the strains isolated between 1979 and 1985 were susceptible to tetracycline, levofloxacin, and imipenem and showed low-level resistance to oxacillin. In contrast, the majority of MRSA strains isolated in 1999 were highly resistant to all of the antibiotics tested. The MICs of 1992 isolates for nine

antibiotics reported by Tanaka et al. (46) are very similar to those of the 1999 isolates, indicating that MRSA strains became highly resistant to many antibiotics in the early 1990s.

**Distribution of virulence-related genes.** The prevalence of virulence-related genes in all 257 MRSA strains was examined by PCR amplification of four gene loci: *lukS-PV-lukF-PV*, encoding Pantone-Valentine leukocidin; *cna*, encoding collagen adhesion protein; *seh*, encoding staphylococcal enterotoxin H; and *tsst-1*, encoding toxic shock syndrome toxin-1. Three of these genes, *lukS-PV-lukF-PV*, *cna*, and *seh*, were identified in the highly virulent community-acquired MRSA (C-MRSA) strain MW2. The results are summarized in Table 2. Forty-four of 97 MRSA strains (45.3%) isolated between 1979 and 1985 carried *lukS-PV-lukF-PV* genes. In contrast, none of the MRSA strains isolated in 1992 and 1999 carried these genes. We detected *cna* in 44 of 97 MRSA strains isolated between 1979 and 1985, whereas only a few MRSA strains isolated in 1992 and 1999 carried this gene. The prevalence of the *seh* gene was very low, being identified in only four strains isolated in 1999, indicating that this gene was rarely carried by hospital-associated MRSA (H-MRSA) strains in Japan. There was a low prevalence of TSST-1-positive strains among those isolated between 1979 and 1985; however, the majority of the MRSA strains isolated in both the early and late 1990s carried *tsst-1* (86.4% [19 of 22] and 92.0% [127 of 138], respectively).

**Comparison of genotypes of strains chosen from different periods.** To identify the genotypes of MRSA strains isolated at different periods of time, MLST and coagulase typing were conducted on strains representing each combination of SCCmec type and *lukS-PV-lukF-PV*- or *tsst-1*-positive gene profiles (Table 2). Among MRSA strains isolated between 1979 and 1985, ST30-coagulase type 4 strains were the most dominant, followed by ST5-coagulase type 2 strains and ST8-coagulase type 3 strains. All PVL-positive and TSST-1-negative strains were ST30-type 4 coagulase producers and carried a type I or type IV SCCmec element. Curiously, we found a PVL-positive/TSST-1-positive strain that carried a type II SCCmec and was a ST5-coagulase type 2 producer. Among the MRSA strains isolated in the 1990s, ST5-coagulase-type 2 strains were the most dominant, followed by ST81-coagulase type 7 strains. All

TABLE 4. PCR identification for four integrated PVL-carrying genes<sup>a</sup>

Name of tested strain	Coagulase isotype	MLST ST	SCCmec	Regions common to $\phi$ 108PVL						Leftmost regions specific to:			Rightmost regions specific to:	
				A	B	C	D	E	F	$\phi$ PVL	$\phi$ SLT	$\phi$ Sa2mw	$\phi$ PVL	$\phi$ SLT and $\phi$ Sa2mw
JCSC2973	4	30	1B (I)	+	+	+	-	+	-	-	-	-	-	-
JCSC4467	4	30	1B (I)	+	+	+	-	+	-	-	-	-	-	-
JCSC4483	4	30	1B (I)	+	+	+	-	+	-	-	(+)	-	-	-
JCSC2958	2	5	2A.1	+	-	-	-	-	(+)	-	-	-	-	(+)
JCSC4451	4	30	2A.n (II)	+	+	+	+	+	+	-	-	-	-	-
JCSC2913	4	N	2B.1 (IVa)	+	+	+	+	+	+	-	-	-	-	-
81/108	4	30	2B.3 (IVc)	+	+	+	+	+	+	-	-	-	-	-
JCSC4458	4	30	2B.3 (IVc)	+	+	+	+	+	+	-	-	-	-	-
JCSC4461	4	30	2B.3 (IVc)	+	+	+	+	+	+	-	(+)	-	-	(+)
JCSC4485	4	30	2B.3 (IVc)	+	+	+	+	+	+	-	-	-	-	-
155-2	4	30	2B.3 (IVc)	+	+	+	+	+	-	-	-	-	+	-
174-2	4	30	2B.3 (IVc)	+	+	+	(+)	+	(+)	-	(+)	-	-	-
JCSC4465	4	30	2B.n	+	+	+	-	+	-	-	+	-	-	-

<sup>a</sup> N, nontypeable; ST, sequence type. +, present; (+), might be present (the amount of amplified DNA fragment is not much); -, absent.




FIG. 3. Alignment of the four PVL-carrying phages. Structures of  $\phi$ 108PVL,  $\phi$ PVL,  $\phi$ SLT, and  $\phi$ Sa2mw are indicated based on the following nucleotide sequences:  $\phi$ 108PVL (DDBJ, EMBL, and GenBank databases under accession no. AB243556),  $\phi$ PVL (DDBJ, EMBL, and GenBank accession no. AB009866),  $\phi$ SLT (DDBJ, EMBL, and GenBank accession no. NC\_002661), and  $\phi$ Sa2mw (DDBJ, EMBL, and GenBank accession no. BA000033). Genes having sequence identities of more than 90% are linked by light blue shading. Known functions of ORFs are colored as follows: lysogeny, blue; DNA replication, red; recombination, pink; DNA packaging and head, yellow; tail, green; lysis, dark blue; *lukS-PV-lukF-PV*, black.

tested type II.1 SCCmec strains, which were PVL negative/TSST-1 positive, were ST5-coagulase type 2 producers.

**Identification of a novel PVL-carrying phage  $\phi$ 108PVL.** It has been well established that *lukS-PV* and *lukF-PV* are encoded by a prophage that integrated into the *S. aureus* chromosome. To date, the nucleotide sequences of three temperate phages carrying the *lukS-PV-lukF-PV* genes,  $\phi$ PVL,  $\phi$ SLT, and  $\phi$ Sa2mw, have been reported (2, 28, 37). To determine whether MRSA strains isolated in the early 1980s harbored one of the three extant PVL-carrying phages, we designed two primer sets to identify three phages specifically: a primer pair composed of a primer specific for integrases that are common to three phages and three primers specific for the repressor gene of each PVL-carrying phage,  $\phi$ PVL,  $\phi$ SLT, and  $\phi$ Sa2mw; a primer pair composed of a primer specific for the *lukS* gene that is common to three phages and another two primers, one is specific for an ORF, P052, in  $\phi$ PVL and the other is specific for ORFs of unknown function that are conserved in two phages,  $\phi$ SLT and  $\phi$ Sa2mw. PCR was carried out using chromosomal DNAs from 34 of the 44 PVL-positive MRSA strains. The results from selected strains, representing different combinations of SCCmec type and exotoxin repertoire, are presented in Table 4. There were no strains that carry either one of three phages. A strain showed a positive result by PCR for identifying the gene lineage *int* to *rep*, and a strain showed a positive result by PCR for identifying the gene lineage *lukS* to an ORF of unknown function. However, no strain showed a positive result with both primer sets, indicating that these strains did not carry one of three extant PVL-carrying phages.

To determine whether these strains carried a prophage different from those of the three extant phages, DNA fragments

spanning the entire phage genome were amplified by long PCR and sequenced using the chromosomal DNA of MRSA strain 81/108, a ST30-type IV.3 SCCmec strain. The characteristic 25-bp sequences *attP* left and *attP* right, which are located at both ends of the phage genome, were present. Judging from the locations of *attP* left and *attP* right, we estimated that the size of the prophage carried by MRSA strain 81/108, which we termed  $\phi$ 108PVL, was 44,107 bp in length. This was comparable to the sizes of the three extant PVL phages,  $\phi$ PVL,  $\phi$ SLT, and  $\phi$ Sa2mw, which are 41,421, 42,942, and 45,924 bp in size, respectively. Two 29-bp conserved core sequences were identified adjacent to the *attP* left and *attP* right sequences. The core sequences in  $\phi$ 108PVL were identical to the corresponding sequences in  $\phi$ PVL but differed by 2 bp from the conserved core sequences in  $\phi$ SLT and  $\phi$ Sa2mw.

Figure 2A illustrates the genomic organization of  $\phi$ 108PVL. Using BLAST to search for homologies to the three PVL-carrying phages ( $\phi$ PVL, DDBJ/EMBL/GenBank accession no. AB009866;  $\phi$ SLT, DDBJ/EMBL/GenBank accession no. NC\_002661; and  $\phi$ Sa2mw, DDBJ/EMBL/GenBank accession no. BA000033) (Fig. 2C), a total of 59 predicted ORFs larger than 99 bp were identified in and around  $\phi$ 108PVL. The organization of the genome of  $\phi$ 108PVL was very similar to that of the extant PVL-carrying phages and contained regions related to lysogeny, DNA replication/transcriptional regulation, packaging/head, tail, and lysis as well as *lukS-PV-lukF-PV* (Fig. 3; Table 5). Although all four phages were similar in their genomic organizations, not all of the ORFs encoded by the four phages were homologous. There were complexes of ORFs that were highly homologous, and those that appeared to be distantly related.







Among the four phages, *lukS-PV* and *lukF-PV* genes were conserved with a predicted amino acid identity of greater than 99.8%. In addition, two genes encoding holin (*hol*) and amidase (*ami*), which are located upstream of *lukS-PV-lukF-PV*, were also well conserved, with predicted amino acid identities of more than 94.6%. We also identified 49 nucleotides located upstream of *hol* that were conserved among all four phages, with 98.0% nucleotide identity, while 555 nucleotides upstream of *hol* were highly homologous between  $\phi$ PVL and  $\phi$ 108PVL. The 204 nucleotides located between *lukS-PV-lukF-PV* and *attP* right were also highly conserved among the four phages, with nucleotide identities of more than 99.5%.

Integrases from four phages were identical with a ratio of more than 98% amino acid identity, suggesting that all four phages had integrated at the same position on the staphylococcal chromosome. In addition, 58 nucleotides between *attP* left and *int* were highly conserved with 98.3 to 100% nucleotide identities. A region located upstream of *int* and an additional sequence of 886 nucleotides in  $\phi$ 108PVL were identical to the corresponding regions in  $\phi$ Sa2mw, and a 489-nucleotide region in  $\phi$ 108PVL had very high nucleotide identity to  $\phi$ SLT and  $\phi$ PVL (90.9 to 91.1%).

Other highly conserved ORFs included a putative DNA binding protein and a hypothetical protein, P028.

The ORFs that were conserved among at least three of the phages were as follows: P004 and P005, conserved among  $\phi$ 108PVL,  $\phi$ SLT, and  $\phi$ Sa2mw with predicted amino acid identities of more than 99.5% (Fig. 2C and 3), and P027, P030, and P031, conserved among  $\phi$ 108PVL,  $\phi$ PVL and  $\phi$ SLT with predicted amino acid identities of more than 85.7% (Fig. 2C).

Twenty-four ORFs in  $\phi$ 108PVL were highly homologous to only  $\phi$ PVL (Fig. 2C and 3): P006 (cI-like repressor), P008 (antirepressor), P009 (transcriptional regulator), and P033 to P053 (DNA packaging and head and tail formation). The high identities between  $\phi$ 108PVL and  $\phi$ PVL among genes involved in morphogenesis suggested that  $\phi$ 108PVL belonged to *Siphoviridae*, the same group as  $\phi$ PVL, based on the taxonomy criteria based on the genetic organization of the structural gene cluster (4).

Six ORFs in  $\phi$ 108PVL related to DNA replication, P015, P020, P022, P023, P025, and P032, were homologous to only the corresponding ORFs in  $\phi$ SLT (Fig. 2). It is notable that only a small ORF, P003, located downstream of *int*, was homologous to the corresponding ORF in  $\phi$ Sa2mw (Fig. 2). The remaining ORFs were unique to  $\phi$ 108PVL (Fig. 2). Among them, ORF P056, which encodes a putative transposase, was located just upstream of *lukS-PV*. BLAST searches of this ORF revealed that it was identical to transposase encoded by IS1272-SA, which was identified in a strain, MRSA 252, and a phage, PV83.

#### Dissemination of $\phi$ 108PVL among Japanese MRSA strains.

To determine whether PVL-positive MRSA strains isolated between 1979 and 1985 carried  $\phi$ 108PVL, we performed PCR experiments using chromosomal DNA from selected strains and six primer sets designed to amplify the following six genes or gene alleles in  $\phi$ 108PVL: set A, integrase; set B, terminase large subunit; set C, portal protein; set D, tail-length tape measure protein; set E, tail-length tape measure protein; and set F, transposase and the region at the rightmost end of

$\phi$ 108PVL (Fig. 2A and Table 1). All strains were positive in PCR with primer set A, indicating that they shared the same integrase and that the integration site was located at the same position on the bacterial chromosome. We found a strain (ST5-SCCmec type II.1) negative by PCR with the remaining five primer sets, indicating that it might carry a PVL-carrying phage which was distantly related to  $\phi$ 108PVL. Three ST30-SCCmec type I strains and an ST30-type IV.n SCCmec (a type IV SCCmec strain of which the J1 region was not classified into extant J1 regions of type IV SCCmec elements) were positive by PCR with primer sets B, C, and E and negative by PCR with primer sets D and F. A SCCmec type IV.1 strain and five SCCmec type IV.3 strains, all of which belonged to ST30, gave positive results with all other five sets of primers, indicating that they carried  $\phi$ 108PVL. A SCCmec type IV.3 strain was positive using four sets of primers, indicating that it carried a phage similar to  $\phi$ 108PVL, differing only in the presence of transposase encoded by IS1272-SA, which is found in MRSA strain 252 in its intact form.

## DISCUSSION

**The changing epidemiology of MRSA strains disseminated in Japanese hospitals.** We conducted a retrospective study of MRSA strains disseminated in Japanese hospitals between 1979 and 1980 using molecular epidemiological methods. We showed that MRSA clones predominating in Japanese hospitals have changed drastically in 20 years. In contrast to MRSA clones predominated in 1999, most of which were ST5-SCCmec type II strains, the majority of MRSA strains isolated between 1979 and 1980 belonged to ST30 and carried either a type IV or a type I SCCmec element. In contrast to MRSA strains isolated in 1999, which were highly resistant against all tested antibiotics, MRSA strains isolated between 1979 and 1980 showed heterogeneous resistance to oxacillin and were susceptible to carbapenems, new quinolones, and tetracycline. They were different in toxin repertoire, too. The *lukS-PV-lukF-PV* genes, which are identified in the majority of C-MRSA strains, were identified in only MRSA strains isolated between 1979 and 1985. Since 22 MRSA strains isolated in Tokyo University Hospital in 1992 showed characteristics similar to those of MRSA strains isolated in 1999, we suppose that the shift of MRSA clones might have occurred in the early 1990s, at least in the case of Tokyo University Hospital. The type of coagulase correlated very well to the MLST genotype.

The change of MRSA clinical isolates, noticed in many facilities as the change of antibiotic susceptibility patterns and coagulase types, can be regarded as the change of MRSA clones from ST30-type I SCCmec or ST30-type IV SCCmec to ST5-type II SCCmec. In Japan, carbapenems, new quinolones, and minocyclines have been used for the treatment of MRSA infections since the mid-1980s to the late 1980s. Extensive use of antibiotics might exert selective pressures on bacteria, and only those strains that carry or acquire resistance genes or that acquire resistance through mutation are able to adapt and survive. Under the selective pressure caused by the extensive use of antibiotics, MRSA clones carrying type II SCCmec, which carries several resistance genes, might have replaced to the MRSA strains predominating in early 1980s, which could to adapt to the environmental change.

The historical shift of MRSA clones is not limited to Japanese hospitals. In 1960s, MRSA clones represented by strain COL or NCTC10442 (ST250-type 1 SCCmec) predominated in the United Kingdom. But, other clones, e.g., epidemic MRSA 16, represented by MRSA 252 (ST36-type II SCCmec), began to predominate in the United Kingdom in 1990s (15, 17, 26). In a Greece hospital, a change from ST30-type IV SCCmec strains to ST239-type III or type IIIA SCCmec IV had occurred (1). The ancient MRSA clones might have been replaced by the highly resistant MRSA clone, which has a strong capacity to spread or to survive under the selective pressure of antibiotics.

**Characteristics of MRSA strains isolated between 1979 and 1985.** It is interesting that approximately three-fourths of MRSA strains isolated between 1979 and 1985 carried either a type IV SCCmec element, which was identified primarily in C-MRSA strains, or a type I SCCmec element, which was identified in the first reported MRSA strains in England. Since we used strains that were stocked 20 years ago, it was very difficult to classify them into hospital-associated MRSA strains or community-associated MRSA strains due to the lack of detailed information that is required for the definition, e.g., the date of hospitalization or previous association to medical facilities. Using the records from two hospitals, we could classify the MRSA strains into two groups, strains obtained from outpatients and strains obtained from inpatients. They were as follows (the type is followed by the number obtained from inpatients and outpatients, respectively): isolates from Gunma Hospital, type I, 2 and 5; type II.1, 3 and 5; type IV.1, 0 and 1; type IV.3, 14 and 5; type IV.4, 8 and 2; type IV.n, 2 and 0; and nontypeable, 1 and 1; isolates from Tokyo University Hospital, type I, 2 and 4; type II.1, 0 and 1; type IV.3, 11 and 4; and type IV.n, 2 and 0. This analysis indicated that there was no significant difference among SCCmec types carried by outpatients and those carried by inpatients. It could be presumed that MRSA strains, which were similar to hospital-associated MRSA strains, predominated in the community as well. In the case of a large outbreak of *S. aureus* in Uruguay, which was caused by a highly virulent Uruguay clone represented by UR6 (ST30-type IV.3 SCCmec), the Uruguay clones were identified not only from community isolates but also from hospital isolates (32). These data suggested that both C-MRSA strains and H-MRSA strains could not be defined by their genotypes and SCCmec types, although representative C-MRSA clones and H-MRSA clones have been identified.

MRSA strains isolated between 1979 and 1985 carried *lukS-PV-lukF-PV* genes in a ratio of 45.3%. Since none of the tested MRSA strains isolated in 1992 and 1999 carried *lukS-PV-lukF-PV* genes, it is a remarkable characteristic of MRSA strains isolated between 1979 and 1985. Oka et al. reported that mortality associated with bacteremia was very high at the ratio of 47.8% (93 cases total) at Tokyo Metropolitan Geriatric Hospital from 1973 through 1984 (38). Although *lukS-PV-lukF-PV* genes were identified in highly virulent C-MRSA strains, e.g., MW2 that caused the death of healthy infants, we do not have the data to conclude whether the cause of high mortality is due to the presence of PVL-positive MRSA strains. Further retrospective studies of MRSA strains isolated from patients with well-documented clinical histories will help answer that question.

**Evolution of PVL-carrying phages.** It is well known that PVL-positive *S. aureus* strains harbor a bacteriophage carrying *lukS-PV-lukF-PV*. All extant PVL-carrying phages belonged to the class of Shi21-like *Siphoviridae*. Canchaya et al. classified the *Staphylococcus* prophages into five groups based on the similarities in genomic structures, predominantly of the genes related to morphogenesis, such as those specifying the phage head and tail (4).  $\phi$ SLT and  $\phi$ Sa2mw were determined to be part of one group based on nucleotide sequence identity in the region encoding the head and tail genes.  $\phi$ SLT has an elongated shape, and  $\phi$ Sa2mw is presumed to have a similar morphology. Similarly,  $\phi$ PV83 and  $\phi$ PVL were determined to belong to a second group and exhibit an icosahedral head morphology (27). The novel phage that we identified in the current study, which we designated  $\phi$ 108PVL, shared basic structural components with all three extant PVL-carrying phages and showed the highest similarity to  $\phi$ PVL.

The four PVL-carrying phages had two highly homologous regions, the regions in and around the gene encoding integrase and the regions in and around the lysis-related genes and *lukS-PV-lukF-PV*. These observations strongly suggested that PVL-carrying phages evolved from non-PVL-carrying phage by acquiring the region containing *lukS-PV-lukF-PV*, although it is not certain whether the acquisition of the region containing *lukS-PV-lukF-PV* was earlier than the acquisition of the region carrying the gene for integrase. Discrepancies in the lengths of the homologous regions among these phages suggested that the acquisition of these two regions might have occurred independently by one or more illegitimate recombination events.

**Characteristics of  $\phi$ 108PVL.** The  $\phi$ 108PVL carried a transposase which is not present in the three extant PVL phages. The noncoding region between the genes for amidase and transposase had very low similarity to other phages, but some noncoding regions between transposase and *lukS-lukF* showed rather high similarity, although the length of this region differed for each, i.e., the lengths were 339 bp ( $\phi$ PVL, 98.2%), 340 bp ( $\phi$ SLT, 97.4%), and 362 bp ( $\phi$ Sa2mw, 99.5%). These results suggested that  $\phi$ 108PVL acquired transposase, along with the noncoding region between the genes for amidase and transposase, as encoded by IS1272-SA. It seems that IS1272 integrated into the phage, as is the case for  $\phi$ PV83.

Kaneko and Kamio found no evidence for a tail structure on the  $\phi$ PVL phage particle using electron microscopy, although the nucleotide sequence of this phage suggests that it carries tail-related genes and that  $\phi$ PVL was defective in its ability to infect any *S. aureus* strains experimentally (27). Since most of the ST30 MRSA strains examined to date carry  $\phi$ 108PVL by PCR, it can be presumed that the phage was responsible for disseminating the PVL gene among MRSA strains isolated in the early 1980s in Japan. However, when we examined the genes encoding tail-length tape measure protein in  $\phi$ 108PVL, we found that the coding region was split into two ORFs, P049 and P050, similar to what is observed in  $\phi$ PVL. In contrast,  $\phi$ SLT, which is able to infect *S. aureus* strains experimentally, encodes a large tail-length tape measure protein of 2,067 amino acids. It therefore seemed that  $\phi$ 108PVL and  $\phi$ PVL are both inactive and that the spread of phages among the *S. aureus* strains occurred in an ancient time, and since which time, lysogenized phages have become inactive. Our data re-



vealed that the genome of  $\phi$ 108PVL is a mosaic, suggesting that  $\phi$ 108PVL was produced as a result of lateral gene transfer and illegitimate recombination events. Thus, it seemed reasonable to assume that  $\phi$ 108PVL and  $\phi$ PVL originated from a relatively close ancestor during their revolution.

**The origin of PVL-positive ST30 MRSA strains isolated in Japan.** Infections caused by C-MRSA have been reported worldwide, and their genetic background and the carriage of virulence factors have been examined. Vandenesch et al. reported that all tested C-MRSA strains shared type IV *SCCmec* and *lukS-PV-lukF-PV* genes (48). Although it has been pointed out that the presence of *lukS-PV-lukF-PV* genes could not be used for defining C-MRSA strains, *lukS-PV-lukF-PV* genes have been identified in many C-MRSA strains. They reported that the genotypes of PVL-carrying C-MRSA strains were classified into six strains, ST1, -8, -30, -59, -80, and -93. Enright et al. reported eight major international epidemic MRSA strains, ST5, -8, -22, -36, -45, -289, -247, and -250 (12). Pandemic clones were also reported as New York/Japan clone (ST5-type II *SCCmec*), pediatric clone (ST5-type IV *SCCmec*), archaic clone (ST250-type I *SCCmec*), Iberian clone (ST247-*SCCmec* type 1A), Hungarian clone (ST239-type III *SCCmec*), and Brazilian clone (ST239-*SCCmec* type IIIA) by Oliveira et al. (40). The genotypes of PVL-carrying C-MRSA strains were different from those of pandemic clones other than ST8, which was identified mostly in MSSA strains (12). Subsequent study revealed that most PVL-carrying MRSA strains isolated worldwide belonged to one of six genotypes reported by Vandenesch et al. (48): ST1, from the United States; ST30, from the United States, Australia, England, Singapore, Belgium, Uruguay, and Japan; ST80, from England, The Netherlands, Denmark, Greece, Belgium, France, Switzerland, and Germany; ST8, from Belgium and the United States; and ST59, from the United States and Taiwan (3, 7, 10, 14, 18, 19, 32, 41, 45, 49, 50).

It is noteworthy that Japanese PVL-positive MRSA strains are mostly ST30, known as the southwest Pacific clone after the region where it was first described (48).

But when we investigated the subtypes of type IV *SCCmec* elements carried by ST30 strains, we found that different subtypes of *SCCmec* elements were carried by these strains. Australian strains carried type IV.1 *SCCmec* elements, whereas the Uruguay and Japanese PVL-positive ST30 MRSA clones carried a type IV.3 *SCCmec* similar to those of Japanese MRSA strains. However, their banding patterns under pulsed-field gel electrophoresis were closely related (32).

Robinson et al. reported that phage type 80/81 MSSA strains, which have been spread throughout the world, were PVL positive and belonged to ST30 and suggested that these MSSA strains might have changed to MRSA strains through the acquisition of different types of *SCCmec* elements independently (42). It seems likely that ST30 MSSA strains carrying *lukS-PV-lukF-PV* existed in the Japanese community as well and evolved into MRSA strains through the acquisition of a type of *SCCmec* element. This is partly supported by the observation that 38% (15 of 39) of MSSA strains isolated in the 1960s carried *lukS-PV-lukF-PV* (X. X. Ma et al., unpublished data). The diversity in the types of *SCCmec* elements identified in these isolates indicates that integration of *SCCmec* elements occurred independently in several different time periods.

In this paper, we introduced a method to distinguish PVL-carrying phage by PCR. By adopting the method, we found that most Japanese MRSA strains disseminated in 1979 to 1985 carried  $\phi$ 108PVL and that a ST5-type II *SCCmec* MRSA strain carried a phage which was different from three extant PVL phages as well as  $\phi$ 108PVL (X. X. Ma et al., unpublished). By using our methods, we will be able to classify PVL phages to know which phage was integrated in MSSA strains and whether ST30 MRSA strains isolated in Australia and Uruguay carried a phage homologous to  $\phi$ 108PVL. Further studies are awaited to learn the origin or spread of PVL-positive MRSA strains.

#### ACKNOWLEDGMENTS

We thank Toyoji Okubo and Shizuko Iyobe of Gunma University for providing us with MRSA strains.

This work was supported by a grant-in-aid for 21st Century COE Research and a grant-in-aid for Scientific Research on Priority Areas from the Ministry of Education, Science, Sports, Culture and Technology of Japan.

#### REFERENCES

- Aires de Sousa, M., C. Bartzavali, I. Spiliopoulou, I. S. Sanches, M. I. Crisostomo, and H. de Lencastre. 2003. Two international methicillin-resistant *Staphylococcus aureus* clones endemic in a university hospital in Patras, Greece. *J. Clin. Microbiol.* **41**:2027–2032.
- Baba, T., F. Takeuchi, M. Kuroda, H. Yuzawa, K. Aoki, A. Oguchi, Y. Nagai, N. Iwama, K. Asano, T. Naimi, H. Kuroda, L. Cui, K. Yamamoto, and K. Hiramatsu. 2002. Genome and virulence determinants of high virulence community-acquired MRSA. *Lancet* **359**:1819–1827.
- Boyle-Vavra, S., B. Ereshesky, C. C. Wang, and R. S. Daum. 2005. Successful multiresistant community-associated methicillin-resistant *Staphylococcus aureus* lineage from Taipei, Taiwan, that carries either the novel staphylococcal chromosome cassette *mec* (*SCCmec*) type V<sub>T</sub> or *SCCmec* type IV. *J. Clin. Microbiol.* **43**:4719–4730.
- Canchaya, C., C. Proux, G. Fournous, A. Bruttin, and H. Brussow. 2003. Prophage genomics. *Microbiol. Mol. Biol. Rev.* **67**:238–276.
- Chambers, H. F. 2001. The changing epidemiology of *Staphylococcus aureus*. *Emerg. Infect. Dis.* **7**:178–182.
- Chongtrakool, P., T. Ito, X. X. Ma, Y. Kondo, S. Trakulsomboon, C. Tien-sasitorn, M. Jamklang, T. Chavalit, J. H. Song, and K. Hiramatsu. 2006. Staphylococcal cassette chromosome *mec* (*SCCmec*) typing of methicillin-resistant *Staphylococcus aureus* strains isolated in 11 Asian countries: a proposal for a new nomenclature for *SCCmec* elements. *Antimicrob. Agents Chemother.* **50**:1–12.
- Coombs, G. W., G. R. Nimmo, J. M. Bell, F. Huygens, F. G. O'Brien, M. J. Malkowski, J. C. Pearson, A. J. Stephens, and P. M. Giffard. 2004. Genetic diversity among community methicillin-resistant *Staphylococcus aureus* strains causing outpatient infections in Australia. *J. Clin. Microbiol.* **42**:4735–4743.
- Deguchi, K. 1982. Sensitivity of *Staphylococcus aureus* isolated clinically to CEPs. Results obtained with strains originating in superficial abscess in infants. *Jpn. J. Antibiot.* **35**:807–811. (In Japanese.)
- Deguchi, K., N. Yokota, M. Koguchi, Y. Nakane, Y. Suzuki, S. Fukuyama, and R. Ishihara. 1991. Biological types and drug-sensitivities of multiple strains of freshly isolated methicillin-resistant *Staphylococcus aureus*. *Jpn. J. Antibiot.* **44**:1376–1385. (In Japanese.)
- Denis, O., A. Deplano, H. De Beenhouwer, M. Hallin, G. Huysmans, M. G. Garrino, Y. Glupczynski, X. Malaviole, A. Vergison, and M. J. Struelens. 2005. Polyclonal emergence and importation of community-acquired methicillin-resistant *Staphylococcus aureus* strains harbouring Panton-Valentine leucocidin genes in Belgium. *J. Antimicrob. Chemother.* **56**:1103–1106.
- Eady, E. A., and J. H. Cove. 2003. Staphylococcal resistance revisited: community-acquired methicillin resistant *Staphylococcus aureus*—an emerging problem for the management of skin and soft tissue infections. *Curr. Opin. Infect. Dis.* **16**:103–124.
- Enright, M. C., D. A. Robinson, G. Randle, E. J. Feil, H. Grundmann, and B. G. Spratt. 2002. The evolutionary history of methicillin-resistant *Staphylococcus aureus* (MRSA). *Proc. Natl. Acad. Sci. USA* **99**:7687–7692.
- Enright, M. C., N. P. Day, C. E. Davies, S. J. Peacock, and B. G. Spratt. 2000. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J. Clin. Microbiol.* **38**:1008–1015.
- Faria, N. A., D. C. Oliveira, H. Westh, D. L. Monnet, A. R. Larsen, R. Skov, and H. de Lencastre. 2005. Epidemiology of emerging methicillin-resistant



- Staphylococcus aureus* (MRSA) in Denmark: a nationwide study in a country with low prevalence of MRSA infection. *J. Clin. Microbiol.* **43**:1836–1842.
15. Gill, S. R., D. E. Fouts, G. L. Archer, E. F. Mongodin, R. T. Deboy, J. Ravel, I. T. Paulsen, J. F. Kolonay, L. Brinkac, M. Beanan, R. J. Dodson, S. C. Daugherty, R. Madupu, S. V. Angiuoli, A. S. Durkin, D. H. Haft, J. Vamathevan, H. Khouri, T. Utterback, C. Lee, G. Dimitrov, L. Jiang, H. Qin, J. Weidman, K. Tran, K. Kang, I. R. Hance, K. E. Nelson, and C. M. Fraser. 2005. Insights on evolution of virulence and resistance from the complete genome analysis of an early methicillin-resistant *Staphylococcus aureus* strain and a biofilm-producing methicillin-resistant *Staphylococcus epidermidis* strain. *J. Bacteriol.* **187**:2426–2438.
  16. Hiramatsu, K. 1995. Molecular evolution of MRSA. *Microbiol. Immunol.* **39**:531–543.
  17. Holden, M. T., E. J. Feil, J. A. Lindsay, S. J. Peacock, N. P. Day, M. C. Enright, T. J. Foster, C. E. Moore, L. Hurst, R. Atkin, A. Barron, N. Bason, S. D. Bentley, C. Chillingworth, T. Chillingworth, C. Churcher, L. Clark, C. Corton, A. Cronin, J. Doggett, L. Dowd, T. Feltwell, Z. Hance, B. Harris, H. Hauser, S. Holroyd, K. Jagels, K. D. James, N. Lennard, A. Line, R. Mayes, S. Moule, K. Mungall, D. Ormond, M. A. Quail, E. Rabinowitz, K. Rutherford, M. Sanders, S. Sharp, M. Simmonds, K. Stevens, S. Whitehead, B. G. Barrell, B. G. Spratt, and J. Parkhill. 2004. Complete genomes of two clinical *Staphylococcus aureus* strains: evidence for the rapid evolution of virulence and drug resistance. *Proc. Natl. Acad. Sci. USA* **101**:9786–9791.
  18. Holmes, A., M. Ganner, S. McGuane, T. L. Pitt, B. D. Cookson, and A. M. Kearns. 2005. *Staphylococcus aureus* isolates carrying Panton-Valentine leukocidin genes in England and Wales: frequency, characterization, and association with clinical disease. *J. Clin. Microbiol.* **43**:2384–2390.
  19. Hsu, L. Y., T. H. Koh, T. Y. Tan, T. Ito, X. X. Ma, R. T. Lin, and B. H. Tan. 2006. Emergence of community-associated methicillin-resistant *Staphylococcus aureus* in Singapore: a further six cases. *Singap. Med. J.* **47**:20–26.
  20. Ichikawa, T., et al. 1966. Drug resistance of staphylococci. *Chemotherapy* **14**:1–8. (In Japanese.)
  21. Ito, T., K. Kuwahara, and K. Hiramatsu. Staphylococcal cassette chromosome *mec* (SCC*mec*) analysis of MRSA. *Methods Mol. Med.*, in press.
  22. Ito, T., Y. Katayama, K. Asada, N. Mori, K. Tsutsumimoto, C. Tiensasitorn, and K. Hiramatsu. 2001. Structural comparison of three types of staphylococcal cassette chromosome *mec* integrated in the chromosome in methicillin-resistant *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **45**:1323–1336.
  23. Ito, T., X. X. Ma, F. Takeuchi, K. Okuma, H. Yuzawa, and K. Hiramatsu. 2004. Novel type V staphylococcal cassette chromosome *mec* driven by a novel cassette chromosome recombinase, *ccrC*. *Antimicrob. Agents Chemother.* **48**:2637–2651.
  24. Ito, T., Y. Katayama, and K. Hiramatsu. 1999. Cloning and nucleotide sequence determination of the entire *mec* DNA of pre-methicillin-resistant *Staphylococcus aureus* N315. *Antimicrob. Agents Chemother.* **43**:1449–1458.
  25. Jevons, M. P. 1961. "Celbenin"-resistant staphylococci. *Br. Med. J.* **124**:124–125.
  26. Johnson, A. P., H. M. Aucken, S. Cavendish, M. Ganner, M. C. Wale, M. Warner, D. M. Livermore, and B. D. Cookson. 2001. Dominance of EMRSA-15 and -16 among MRSA causing nosocomial bacteraemia in the UK: analysis of isolates from the European Antimicrobial Resistance Surveillance System (EARSS). *J. Antimicrob. Chemother.* **48**:143–144.
  27. Kaneko, J., and Y. Kamio. 2004. Bacterial two-component and hetero-heptameric pore-forming cytolytic toxins: structures, pore-forming mechanism, and organization of the genes. *Biosci. Biotechnol. Biochem.* **68**:981–1003.
  28. Kaneko, J., T. Kimura, S. Narita, T. Tomita, and Y. Kamio. 1998. Complete nucleotide sequence and molecular characterization of the temperate staphylococcal bacteriophage  $\phi$ PVL carrying Panton-Valentine leukocidin genes. *Gene* **215**:57–67.
  29. Kanno, H. 1989. Treatment for patients of MRSA infection. *Clin. Dr.* **15**:1950–1952. (In Japanese.)
  30. Katayama, Y., T. Ito, and K. Hiramatsu. 2000. A new class of genetic element, staphylococcal cassette chromosome *mec*, encodes methicillin resistance in *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **44**:1549–1555.
  31. Konno, M. 1989. Etiology and preventive measures for methicillin-resistant *Staphylococcus aureus* (MRSA) infection. *Modern Med.* **44**:2544–2553. (In Japanese.)
  32. Ma, X. X., A. Galiana, W. Pedreira, M. Mowszowicz, I. Christophersen, S. Machiavello, L. Lope, S. Benaderet, F. Buela, W. Vincentino, M. Albini, O. Bertaux, I. Constenla, H. Bagnulo, L. Llosa, T. Ito, and K. Hiramatsu. 2005. Community-acquired methicillin-resistant *Staphylococcus aureus*, Uruguay. *Emerg. Infect. Dis.* **11**:973–976.
  33. Ma, X. X., T. Ito, C. Tiensasitorn, M. Jamklang, P. Chongtrakool, S. Boyle Vavra, R. S. Daum, and K. Hiramatsu. 2002. Novel type of staphylococcal cassette chromosome *mec* identified in community-acquired methicillin-resistant *Staphylococcus aureus* strains. *Antimicrob. Agents Chemother.* **46**:1147–1152.
  34. Matsumoto, K., A. Takahashi, H. Shishido, K. Watanabe, N. Riktomi, H. Suzuki, and T. Yoshida. 1989. A comparison of the changes in *Staphylococcus aureus* during two periods. *Chemotherapy* **37**:549–562. (In Japanese.)
  35. Mitsuhashi, S., H. Hashimoto, T. Kasuga, and T. Sawai. 1965. Relationship between penicillin G resistance and susceptibility to synthetic penicillins in *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **5**:374–379.
  36. Naimi, T. S., K. H. LeDell, K. Como-Sabetti, S. M. Borchardt, D. J. Boxrud, J. Etienne, S. K. Johnson, F. Vandenesch, S. Fridkin, C. O'Boyle, R. N. Danila, and R. Lynfield. 2003. Comparison of community- and health care-associated methicillin-resistant *Staphylococcus aureus* infection. *JAMA* **290**:2976–2984.
  37. Narita, S., J. Kaneko, J. Chiba, Y. Piemont, S. Jarraud, J. Etienne, and Y. Kamio. 2001. Phage conversion of Panton-Valentine leukocidin in *Staphylococcus aureus*: molecular analysis of a PVL-converting phage,  $\phi$ SLT. *Gene* **268**:195–206.
  38. Oka, S., K. Urayama, T. Inamatsu, and K. Shimada. 1986. Staphylococcal bacteremia. 2. Review of 93 cases with *Staphylococcus aureus* bacteremia of the aged. *Kansenshogaku Zasshi* **60**:602–607. (In Japanese.)
  39. Okuma, K., K. Iwakawa, J. D. Turnidge, W. B. Grubb, J. M. Bell, F. G. O'Brien, G. W. Coombs, J. W. Pearson, F. C. Tenover, M. Kapi, C. Tiensasitorn, T. Ito, and K. Hiramatsu. 2002. Dissemination of new methicillin-resistant *Staphylococcus aureus* clones in the community. *J. Clin. Microbiol.* **40**:4289–4294.
  40. Oliveira, D. C., A. Tomasz, and H. Lencastre. 2001. The evolution of pandemic clones of methicillin-resistant *Staphylococcus aureus*: identification of two ancestral genetic backgrounds and the associated *mec* elements. *Microb. Drug Resist.* **7**:349–361.
  41. Pan, E. S., B. A. Diep, H. A. Carleton, E. D. Charlebois, G. F. Sensabaugh, B. L. Haller, and F. Perdreau-Remington. 2003. Increasing prevalence of methicillin-resistant *Staphylococcus aureus* infection in California jails. *Clin. Infect. Dis.* **37**:1384–1388.
  42. Robinson, D. A., A. M. Kearns, A. Holmes, D. Morrison, H. Grundmann, G. Edwards, F. G. O'Brien, F. C. Tenover, L. K. McDougal, A. B. Monk, and M. C. Enright. 2005. Re-emergence of early pandemic *Staphylococcus aureus* as a community-acquired methicillin-resistant clone. *Lancet* **365**:1256–1258.
  43. Shimada, K., S. Oka, H. Suzuki, T. Inamatsu, and K. Urayama. 1985. Staphylococcal bacteremia. 1. Bacteremia due to methicillin-cephem resistant *Staphylococcus aureus*. *Kansenshogaku Zasshi* **59**:459–463. (In Japanese.)
  44. Shore, A., A. S. Rossney, C. T. Keane, M. C. Enright, and D. C. Coleman. 2005. Seven novel variants of the staphylococcal chromosomal cassette *mec* in methicillin-resistant *Staphylococcus aureus* isolates from Ireland. *Antimicrob. Agents Chemother.* **49**:2070–2083.
  45. Takizawa, Y., I. Taneike, S. Nakagawa, T. Oishi, Y. Nitahara, N. Iwakura, K. Ozaki, M. Takano, T. Nakayama, and T. Yamamoto. 2005. A Panton-Valentine leukocidin (PVL)-positive community-acquired methicillin-resistant *Staphylococcus aureus* (MRSA) strain, another such strain carrying a multiple-drug resistance plasmid, and other more-typical PVL-negative MRSA strains found in Japan. *J. Clin. Microbiol.* **43**:3356–3363.
  46. Tanaka, T., K. Okuzumi, A. Iwamoto, and K. Hiramatsu. 1995. A retrospective study on methicillin-resistant *Staphylococcus aureus* clinical strains in Tokyo University Hospital. *J. Infect. Chemother.* **1**:40–49.
  47. Ushioda, H., T. T. Sakai, S. Zen-Yoji, H. Nishiwaki, and A. Hidano. 1981. Coagulase typing of *Staphylococcus aureus* and its application in routine work, p. 77–83. *In* J. J. Gustav (ed.), *Staphylococci and staphylococcus infections*. Fischer Verlag, Stuttgart, Germany.
  48. Vandenesch, F., T. Naimi, M. C. Enright, G. Lina, G. R. Nimmo, H. Heffernan, N. Liassine, M. Bes, T. Greenland, M. E. Reverdy, and J. Etienne. 2003. Community-acquired methicillin-resistant *Staphylococcus aureus* carrying Panton-Valentine leukocidin genes: worldwide emergence. *Emerg. Infect. Dis.* **9**:978–984.
  49. Wannet, W. J., E. Spalburg, M. E. Heck, G. N. Pluister, E. Tiemersma, R. J. Willems, X. W. Huijsdens, A. J. de Neeling, and J. Etienne. 2005. Emergence of virulent methicillin-resistant *Staphylococcus aureus* strains carrying Panton-Valentine leukocidin genes in The Netherlands. *J. Clin. Microbiol.* **43**:3341–3345.
  50. Witte, W., C. Bralke, C. Cuny, B. Strommenger, G. Werner, D. Heuck, U. Jappe, C. Wendt, H. J. Linde, and D. Harmsen. 2005. Emergence of methicillin-resistant *Staphylococcus aureus* with Panton-Valentine leukocidin genes in central Europe. *Eur. J. Clin. Microbiol. Infect. Dis.* **24**:1–5.