

## Two Different Pantone-Valentine Leukocidin Phage Lineages Predominate in Japan<sup>∇</sup>

Xiao Xue Ma,<sup>1,7</sup> Teruyo Ito,<sup>1,2\*</sup> Yoko Kondo,<sup>2</sup> Moe Cho,<sup>2</sup> Yukio Yoshizawa,<sup>3</sup> Jun Kaneko,<sup>4</sup> Atsuo Katai,<sup>5</sup> Masato Higashiide,<sup>6</sup> Shanshuang Li,<sup>2</sup> and Keiichi Hiramatsu<sup>1,2</sup>

Department of Bacteriology, Juntendo University, Tokyo, Japan<sup>1</sup>; Department of Infection Control Science, Graduate School of Juntendo University, Tokyo, Japan<sup>2</sup>; Radioisotope Research Facility, The Jikei University School of Medicine, Tokyo, Japan<sup>3</sup>; Laboratory of Applied Microbiology, Department of Microbial Biotechnology, Graduate School of Agricultural Science, Tohoku University, Sendai, Japan<sup>4</sup>; Division of Laboratory Medicine, Kinan General Hospital, Wakayama, Japan<sup>5</sup>; Kotobiken Medical Laboratories, Inc., Ibaragi, Japan<sup>6</sup>; and Department of Medical Microbiology and Parasitology, China Medical University, Shenyang, China<sup>7</sup>

Received 24 January 2008/Returned for modification 27 April 2008/Accepted 14 July 2008

We determined the entire nucleotide sequence of  $\phi$ Sa2958-carrying Pantone-Valentine leukocidin (PVL) gene, which was lysogenized in a sequence type 5 staphylococcal cassette chromosome *mec* (SCC*mec*) type II strain of methicillin-resistant *Staphylococcus aureus* (MRSA). Based on the nucleotide sequences of PVL phages, we developed PCRs to discriminate among five PVL phages, with a preliminary classification into two morphological groups (elongated-head type and icosahedral-head type) with four PCRs, including two PCRs for identifying the gene lineage between *lukS-PV* and the tail gene. The phages were then classified into five types by four PCRs identifying each phage-specific structure. With these PCRs, we examined the PVL phage types of 67 MRSA strains isolated in Japan from 1979 through 1985 and since 2000 and found that two morphologically distinct phages were predominant in Japan. The icosahedral-head-type phage, represented by the  $\phi$ 108PVL type, was identified for 39 of 53 strains isolated from 1979 through 1985. Of 26 other Japanese isolates, 25 belonged either definitively or presumably to elongated-head types as follows: 3 belonged to the  $\phi$ Sa2958 type; 8 were determined to belong to an elongated-head type, but a determination of greater specificity was not made; and 14 belonged to a  $\phi$ Sa2958-like phage of unknown type. We induced prophages by treatment with mitomycin C from six strains of the  $\phi$ Sa2958 type or of  $\phi$ Sa2958-like unknown-type phages; five of six strains carried intact PVL-carrying phages, which can infect other *S. aureus* strains and might generate novel PVL-positive strains of *S. aureus*. That various SCC*mec* elements were carried by different strains of the same phage type suggests that *S. aureus* strains might independently acquire PVL phages before they acquire various SCC*mec* elements.

Since the early 1990s, the number of strains of methicillin-resistant *Staphylococcus aureus* (MRSA) isolated from community-acquired infections (CA-MRSA) has increased (5, 11, 31). Characteristics of CA-MRSA strains have been investigated and compared with those of health care-associated MRSA strains. The genotypes and staphylococcal cassette chromosome *mec* (SCC*mec*) types of CA-MRSA strains differed greatly from those of health care-associated MRSA strains. In addition, most of CA-MRSA strains are reported to carry the Pantone-Valentine leukocidin (PVL) gene at a high level of incidence (11).

PVL was first reported in 1932 (35). It is a two-component toxin composed of LukF-PV, with a molecular mass of 34 kDa, and LukS-PV, with a molecular mass of 32 kDa (36). These two proteins, LukF-PV and LukS-PV, act synergistically and cause damage on cell membranes by forming pores, resulting in the lysis of polymorphonuclear leukocytes and macrophages (6, 13).

The genes for these two proteins, *lukS-PV* and *lukF-PV*, respectively, are carried by prophages that are integrated into the *S. aureus* chromosome. To our knowledge, five PVL phages,  $\phi$ PVL,  $\phi$ SLT,  $\phi$ Sa2mw,  $\phi$ 108PVL, and  $\phi$ Sa2usa, have

been reported to date (1, 9, 20, 21, 28, 32). These phages carry the *lukS-PV* and *lukF-PV* genes, which show more than 99% nucleotide identity and are integrated at the same position in the *S. aureus* chromosome. However, the morphologies of the phage particles are not identical. The phage morphologies can be classified into two types, namely, the icosahedral-head type and the elongated-head type (4). We have examined PVL-positive MRSA strains isolated from 1979 through 1985 and have determined the nucleotide sequence of  $\phi$ 108PVL, carried by SCC*mec* type IV.3 (SCC*mec* types are designated hereafter by a capital roman numeral [e.g., type IV] or by a capital roman numeral, a decimal point, and an Arabic numeral [e.g., type IV.3]) sequence type 30 (ST30) MRSA strain 81/108 (28). When we examined the carriage of  $\phi$ 108PVL phage in representative MRSA strains isolated from 1979 through 1985, phage  $\phi$ 108PVL was most often identified in ST30 MRSA strains, whereas the SCC*mec* type II.1 ST5 MRSA strain JCSC2958 was negative by five of six PCRs using different sets of primers to identify  $\phi$ 108PVL, indicating that the strain might carry another new PVL phage. Both PVL-positive methicillin-sensitive *S. aureus* (MSSA) strains and MRSA strains have been isolated worldwide, including in the United States (2, 3, 10, 37), France (39), Australia (7), England (15), Canada (30), Singapore (16), Belgium (8), and Uruguay (27). We wondered how these PVL-positive strains evolved by acquiring each PVL phage and how each PVL phage evolved by acquir-

\* Corresponding author. Mailing address: Department of Bacteriology, Juntendo University, Hongo 2-1-1, Bunkyo-ku, Tokyo 113-8421, Japan. Phone: 011-3-5802-1041. Fax: 011-3-5684-7830. E-mail: teruybac@med.juntendo.ac.jp.

<sup>∇</sup> Published ahead of print on 6 August 2008.

ing the *lukS-PV* and *lukF-PV* genes. In an attempt to answer these questions, we determined the nucleotide sequence of a new PVL phage carried by JCSC2958 and developed a PCR system to identify the lysogenized PVL phages. By applying the newly developed PCRs, we have found that Japanese PVL-positive *S. aureus* strains carry two morphologically distinct PVL phages, namely, the icosahedral-head type, represented by  $\phi$ 108PVL phage, and the elongated-head type, represented by  $\phi$ Sa2958, and that these types differ from those for strains disseminated in the United States and France.

#### MATERIALS AND METHODS

**MRSA and MSSA strains.** Sixty-five PVL-positive MRSA strains isolated in Japan were tested. Fifty-three strains were isolated from 1979 through 1985: 12 were isolated at Tokyo University Hospital, 23 at Gunma University Hospital, 10 at Tokyo Geriatric Hospital, and 9 at The Jikei University Hospital. Twelve PVL-positive MRSA strains were isolated over the past 8 years as follows: 1 strain was isolated from pus from the skin of an outpatient at Juntendo University Hospital in 2002; 1 strain was isolated from an inpatient with pneumonia at Moji Rosai Hospital (identified by the surveillance of PVL-positive strains conducted by the SRL Laboratory and kindly provided by Hiroshi Kuramoto); 1 strain was isolated in 2007 by Masato Higashiide from a 7-year-old boy with an abdominal wound; 2 strains were kindly provided by the LVFX Surveillance Group, headed by Keizo Yamaguchi, and consisted of 1 isolated from the pus of a 27-year-old man in 2002 (strain EB00449) and 1 isolated from the centesis of a 61-year-old man in 2000 (strain DB00319) (40); and 7 strains were isolated by Atsuo Katai at Kinan General Hospital between 2005 and 2007. Strains MW2 and 81/108 were used as representative PVL-positive MRSA strains, strain ATCC 49775 was used as a representative PVL-positive MSSA strain, and strain RN4220/ $\phi$ SLT was also used as a control for the induction experiment for PVL-positive phages.

To identify phages induced from lysogenized bacteria, two MSSA strains, RN4220 and 1039, were used as indicator strains. These mutant strains, which lack prophages and restriction systems, were obtained from the NCTC8325 and Terashima strains, respectively (41).

**SCCmec typing and identification of virulence genes.** SCCmec typing and identification of virulence genes were performed with PCRs as described previously (22, 27, 34).

**Determination of the entire nucleotide sequence of  $\phi$ Sa2958.** The fosmid library from the genomic DNA of JCSC2958 was constructed by using the CopyControl fosmid library production kit (Epicentre Biotechnologies, Madison, WI) according to the protocols recommended by manufacturer. Briefly, genomic DNA was prepared from cells of JCSC2958 cells with Isoplant (Nippon Gene Co., Tokyo, Japan). Approximately 400  $\mu$ g of the genomic DNAs was sheared into small fragments by aspiration and expulsion through an injection needle. Subsequent size selection of 40-kb DNA fragments was performed by running samples in agarose gel (SeaKem gold agarose; Cambrex Bio Science Rockland, Inc., Rockland, ME). To recover the size-fractionated DNA after electrophoresis, agar slices containing the corresponding sizes of DNA were cut and then melted in GElase solution. The recovered DNA was purified and treated with the end repair enzyme mix that came with the kit. The purified insert DNA was ligated into CopyControl pCCIFOS cloning-ready vector with DNA ligase and then packaged according to the manufacturer's instructions. Each 10  $\mu$ l of the fosmid library was mixed with 100  $\mu$ l of EPI300-T1R cells and incubated for 20 min to promote phage infection. Each reaction mixture was spread on an L-agar plate containing 12.5  $\mu$ g/ml of chloramphenicol and incubated at 37°C overnight. Five hundred colonies were screened with two primer pairs for the PVL gene and the integrase gene (Table 1), which are conserved in nearly all extant PVL phages. Plasmids were extracted from positive clones and used as templates for nucleotide sequencing. In some cases, DNA fragments amplified by long-range PCR were used as the templates. Primers used for long-range PCR are listed in Table 1. Nucleotide sequences were subsequently determined with the primer-walking method.

**MLST and coagulase typing.** The genotypes of representative strains were determined by means of multilocus sequence typing (MLST) by the method of Enright et al. (12). The coagulase types of all tested strains were determined with the serological method established by Ushioda et al. (38).

**Induction of prophages from *S. aureus* cells.** A 0.3-ml portion of overnight culture was inoculated to 3 ml of BHI broth. After the culture had been shaken for 2 h at 37°C, mitomycin C (Kyowa Hakko Kogyo Co., Ltd., Tokyo, Japan) was

added to a final concentration of 1  $\mu$ g/ml, and the culture was cultivated at 37°C with shaking for 50 min. Cells were then precipitated by centrifugation and resuspended with 300  $\mu$ l of L broth. A 100- $\mu$ l portion was added to 3 ml of L broth and incubated further at 37°C until the cell suspension became transparent. The supernatant was sterilized by means of a membrane filter with 0.45- $\mu$ m-diameter pores (Whatman Corp., Clifton, NJ).

**Identification of PVL-carrying phages by plaque hybridization.** Phage solutions were diluted 10-fold serially, and a 0.1-ml portion of each solution was mixed with a 0.3-ml overnight culture of the RN4220 or 1039 strain and kept at room temperature for 15 min. Three milliliters of L broth containing 0.6% agar was added to each mixture, which was then poured onto heart infusion agar plates. The plates were incubated overnight at 30°C to form plaques on the lawns. Plates with an appropriate numbers of plaques were selected, and the plaques were transferred onto a piece of Biodyne membrane (Pall Biodyne A, pore size, 1.2  $\mu$ m; Pall Life Sciences, Ann Arbor, MI). The filters were denatured by submersion for 5 min in 0.5 M NaOH and 1.5 M NaCl and then neutralized by submersion twice for 5 min in a solution of 0.5 M Tris-HCl (pH 7.3), 1.5 M NaCl, and 1 mM EDTA.

The DNAs were cross-linked to the nylon membrane by use of the Stratilinker UV cross-linker (Stratagene Japan K. K., Tokyo, Japan). Probes to identify PVL-carrying phages were prepared by labeling DNA fragments with digoxigenin by use of the digoxigenin DNA labeling and detection kit (Roche Applied Science, Penzberg, Germany). The DNA fragments used for plaque hybridization were amplified with PCR and genomic DNA of the JCSC2958 strain as a template with a pair of primers to identify the *lukS-PV* and *lukF-PV* genes (Table 1). The subsequent experiment was performed as described by the manufacturer.

**PCRs to classify PVL phages.** We developed eight PCRs to classify the PVL-encoding prophages by use of chromosomal DNA preparations from PVL-positive *S. aureus* strains as templates. Primers used for identifying PVL-encoding phages are listed in Table 1. These primers were designed on the bases of nucleotide sequences of five PVL phages, namely,  $\phi$ 108PVL,  $\phi$ Sa2mw,  $\phi$ PVL,  $\phi$ SLT, and  $\phi$ Sa2958, which are deposited in DDBJ/EMBL/GenBank databases under accession nos. AB009866, BA000033, AB009866, NC\_002661, and AP009363, respectively. The locations of the primers are shown below (see Fig. 3).

Two PCRs were designed to identify the carriage of two morphologically distinct phages. PCR-1 was designed to identify the carriage of phages with isometric hexagonal heads by amplifying the portal gene and the head gene, which are conserved in  $\phi$ PVL and  $\phi$ 108PVL. PCR-2 was designed to identify the carriage of phages with elongated heads by amplifying the portal gene and the head gene, which are conserved in  $\phi$ Sa2mw,  $\phi$ Sa2958, and  $\phi$ SLT. If two DNA fragments could be amplified with PCR-1, we then proceeded to PCR-3, which was designed to identify the gene lineage between the tail gene and the *AP009363* gene and to verify that the tail gene identified with PCR-1 belonged to a PVL-carrying phage with a hexagonal head.

If two DNA fragments were amplified with PCR-2, we proceed to PCR-4, which was designed to identify the gene lineage between the tail gene and the *lukS* gene to verify that the tail gene of the elongated shape is located in relation to the *lukS* gene with a primer pair commonly conserved among  $\phi$ Sa2958,  $\phi$ Sa2mw, and  $\phi$ SLT.

PCR-5 to -8 were designed to identify five PVL phages by amplifying the gene lineage between the integrase gene, which is commonly carried by all reported PVL-carrying phages, and genes located on the region related to the lysogeny or recombination of each PVL phage.

PCR-5 was designed to identify  $\phi$ PVL and  $\phi$ 108PVL by integrase and two open reading frames (ORFs), the repressor for  $\phi$ PVL and the antirepressor for  $\phi$ 108PVL. PCR-6, -7, and -8 were designed to identify  $\phi$ Sa2958,  $\phi$ Sa2mw, and  $\phi$ SLT by detecting the gene lineage between the integrase gene and the genes located downstream of the gene: those for a hypothetical protein (JCP004) for  $\phi$ Sa2958, *cro* repressor protein for  $\phi$ Sa2mw, and a single-stranded binding protein for  $\phi$ SLT.

Chromosomal DNAs were prepared with the small-scale phenol extraction method and used as PCR templates. The PCRs were performed using a Gene Amp 9600 thermal cycler (Perkin-Elmer Cetus Instruments, Emeryville, CA). The reaction mixtures for PCR-1, -2, and -5 to -8 contained 50 ng of template DNA, each oligonucleotide primer (0.2 mM), 400 mM of each deoxynucleoside triphosphate, 1 $\times$  Ex *Taq* buffer with magnesium, and 4 U of Ex *Taq* polymerase (Takara Shuzo Co. Ltd., Kyoto, Japan) in final volumes of 50  $\mu$ l. PCRs for five PCRs (PCR-1, -2, -5, -6, and -7) consisted of 30 cycles of denaturation (95°C, 60 s), annealing (50°C, 60 s), and extension (72°C, 2 min).

PCR-3, -4, and -8 were carried out with long-range PCR using an Expand high-fidelity PCR kit adhering to the protocols recommended by manufacturer. The 6- $\mu$ l PCR mixture was subjected to agarose gel electrophoresis to detect amplified DNA fragments.

TABLE 1. List of primers used in this study

Objective of primer construction	Primer name	Nucleotide sequence (5'-3')	Size (bp) of PCR product	Gene or structure on which primer was designed	PVL phage(s)	Location(s) of primer(s)	
For screening of fosmid library	int-R2	CATTTTAATTGCCAGCATCTTA	558	<i>int</i>	φSa2958	1547–1525	
	int-F2	ATGTTTTTCGAGTTTTTGAGTTAG		<i>int</i>		989–1011	
	PVL-F	ATGTCTGGACATGATCCAA	970	<i>lukS-PV</i>		45386–45404	
	PVL-R	AACTATCTCTGCCATATGGT		<i>lukF-PV</i>		46356–46337	
For long-range PCR to amplify entire structure of φSa2958	From chromosome to integrase	phiMW2-DN1	GCAGAAAAAGATGCGATTGAA	(2.5 kb)	Chromosome of JCSC2958	Upstream of φ2958PVL 1655–1634	
		intR	AGGATATCGAAAAAGATGAATC		<i>int</i>		
	From integrase to DNA polymerase	intc	TTGTAGTGTCTTTGTATCCG	9,214	<i>int</i>	1376–1396 10589–10569	
		MW1425-R	TTGTTGCCATTTTTCAAGATC		<i>pol</i>		
	From DNA polymerase to ORF JP030	MW1425-F	CTAAAGTAGATAATGAGCCTT	8,060	<i>pol</i>	10237–10257 18296–18276	
		2958–1405R1	TCCCTTTTCTTGCTTCATTTT		JP030		
	From virulence-associated protein E to portal protein	CF35	ACGAAGACGATTTTATCAAGG	6,097	<i>por</i>	17509–17529 23605–23585	
		2958-portalR1	CACTATATCTTCAGAGACATA		<i>virE</i>		
	From terminase large subunit to tail length tape measure protein	CF47	AAAGTTATCTAATTCGATGGC	10,480	<i>terL</i>	22990–23010 33469–33450	
		CF67	GGGCTCTTGAATACATATCT		<i>mtp</i>		
From tail length tape measure protein to ORF JP052	2958–1392-F10	ATACTGAAAAGTGGTGAATG	8,741	<i>mtp</i>	32807–32827 41547–41528		
	phi-M-TR1	GACTTCCTAAGTCGAAATAG		JP052			
From ORF JP052 to <i>lukF-PV</i> gene	phi-M-T1	TGGATTAACATAATCTAGTCG	4,877	JP052	41480–41500 46356–46337		
	PVL-R	AACTATCTCTGCCATATGGT		<i>lukF-PV</i>			
From ORF JP058 to chromosome	LukS-RR	TGGTCAACTATATCGTGGTTTT	(2 kb)	JP058	44574–44595 Downstream of φSa2958		
	phiMW-UP	TCGCCACGTTTAGCAATTTTAT		Chromosome of JCSC2958			
For classifying phages	PCR-1 (for identifying φ108PVL and φPVL)	portal-1F	ACACGTGATAAAACAGGAGAA	569	<i>por</i>	φ108PVL	
		portal-1R	TCTAAATTAGCATCCGTGATAC		<i>por</i>		
		tail-1R	ATAATTGGGATAGCAACGCAA	489	<i>mtp</i>		
	PCR-2 (for identifying φ2958PVL, φSLT, and φSa2mw)	portal-2F	GATGGCTAGTTTGCCCTTGA	656	<i>por</i>	φSa2958	
		portal-2R	CTGAGGGCAATTGAAAAACG		<i>por</i>		
		tail-2F	CATAGCGCTAATGTCGCAAA	468	<i>mtp</i>		
	PCR-3 (for identifying the gene lineage between genes of φ108PVL and φPVL, and <i>lukS-PV</i> )	tail-2R	AGCCTCCATTGTTTGTGTTGG		<i>mtp</i>	23005–23024 23660–23641 30040–30059 30507–30488	
		lukSR1	ACGAAGTAGCAATAGGAGTGA	10,497	<i>lukS-PV</i>	φ108PVL	
	teil-ico-F	AGATTTAGAAGAGGAGGCACGA		<i>mtp</i>	42326–42306 31830–31851		
	PCR-4 (for identifying the gene lineage between genes of φSa2958, φSa2mw, and φSLT, and <i>lukS-PV</i> )	lukSR1	ACGAAGTAGCAATAGGAGTGA	9,483	<i>lukS-PV</i>	φSa2958	44861–44841
For identifying each PVL phage	PCR-5 (for identifying φPVL and φ108PVL)	intF2	ATGTTTTTCGAGTTTTTGAGTTAG	4,340	<i>int</i>	φ108PVL, φPVL	393–415, 24310–24332
		108-aR	TCAAATCCGTAATCACTCATTCT		<i>ant</i>	φ108PVL	4732–4710
	PCR-6 (for identifying φSa2958)	PVL-aR	TTCACAACTAAACCTATCATTGT	1,411	JP030	φPVL	25720–25697
		int-F2	ATGTTTTTCGAGTTTTTGAGTTAG	2,238	<i>int</i>	φSa2958	989–1011 3226–3203
	PCR-7 (for identifying φSa2mw)	2958-aR	TGGTAATCAACCATTCACTTATGA		JP004		
		int-F2	ATGTTTTTCGAGTTTTTGAGTTAG	4,065	<i>int</i>	φSa2mw	1574920–1574898 1570856–1570879
	PCR-8 (for identifying φSLT)	MW2-aR	TAAGTTCTGGTGTCACTTCTAAT		<i>cro</i>		
		int-F2	ATGTTTTTCGAGTTTTTGAGTTAG	8,770	<i>int</i>	φSLT	123–145 8892–8868
	SLT-aR	TCTTACCAAATGCAACACAACGAAT		<i>ssb</i>			




FIG. 1. (A) The structure of  $\phi$ Sa2958. Black arrowheads indicate the locations of primers used to amplify the entire  $\phi$ Sa2958 genome. The two red arrowheads flanking the core sequence indicate the *att* sites on the phage element. (B) The ORFs in  $\phi$ Sa2958. The ORFs are shown as squares in six possible reading frames. The direction of the arrow indicates the transcriptional direction for each ORF. Color codes are as follows: dark green, ORFs (or the parts of ORFs) that are well conserved among four other PVL-carrying phages, namely,  $\phi$ Sa2mw,  $\phi$ SLT,  $\phi$ PVL, and  $\phi$ 108PVL; light green, ORFs that are highly homologous to  $\phi$ Sa2mw and  $\phi$ SLT; light purple, ORFs that are highly homologous to  $\phi$ Sa2mw; orange, an ORF that is homologous to  $\phi$ SLT and  $\phi$ PVL; yellow, an ORF that is homologous to  $\phi$ SLT and  $\phi$ 108PVL; white, ORFs that are unique to  $\phi$ Sa2958; light blue, ORFs that are homologous to  $\phi$ Sa2mw; dark blue, ORFs that are homologous to  $\phi$ SLT.

**Nucleotide sequence accession number.** The entire nucleotide sequence of  $\phi$ Sa2958 has been deposited in the DDBJ/EMBL/GenBank database under accession no. AP009363.

## RESULTS

**The structure of a novel PVL phage,  $\phi$ Sa2958.** The  $\phi$ Sa2958 phage is 46,046 bp in length, which is a size comparable to those of other PVL phages identified to date. A total of 59 predicted ORFs of larger than 99 bp were identified in  $\phi$ Sa2958 (Fig. 1 and Table 2). The whole G+C content of the novel PVL phage was 33.1%, which is comparable to those of *S. aureus* genomes. The gene coding potential for  $\phi$ Sa2958 was 90.6%, with approximately 1.28 genes/kbp of nucleotide sequence.

The 29-bp core sequence in  $\phi$ Sa2958 exactly matched the corresponding core sequence conserved in  $\phi$ Sa2mw but differed from those of  $\phi$ SLT,  $\phi$ PVL, and  $\phi$ 108PVL by 1 bp. The 25-bp sequences of the *attB* site located on the chromosome side were identified flanking both ends of  $\phi$ Sa2958. The rightmost (*attB-R*) sequence was identical with that of  $\phi$ Sa2mw, whereas the leftmost (*attB-L*) sequence differed by 2 bp from that of  $\phi$ Sa2mw and differed significantly from those of  $\phi$ PVL and  $\phi$ 108PVL. The 25-bp attachment sites (*attP*) at both ends of the phage were well conserved among all five PVL phages.

The organization of the  $\phi$ Sa2958 genome was similar to those of other extant PVL phages, indicating that the following order of modules on the phage genome is well conserved: regions related to lysogeny, DNA replication/transcriptional regulation regions, the packaging/head, the tail, and the lysis

module as well as *lukS-PV* and *lukF-PV*. Fifty-nine ORFs were roughly classified into five groups on the basis of similarities with extant PVL phages.

Group 1 contained 10 genes (JP013, JP014, JP023, JP024, JP027, JP056, JP057, JP058, JP059, and JP060) conserved among five extant PVL phages, with nucleotide identities of more than 90% (Fig. 1). The integrase gene (*int*) was located at the leftmost side of  $\phi$ Sa2958. It showed high similarity to *int* genes carried by extant PVL phages, with more than 98% identity in amino acids. ORFs JP056, JP057, JP059, and JP060 encoded holin, amidase, and LukS-PV and LukF-PV proteins, respectively. Both the *lukF-PV* and *lukS-PV* genes showed nucleotide identities of more than 99% with those of other PVL phages. In particular, the *lukF-PV* gene in  $\phi$ Sa2958 exhibited 100% homology with that of  $\phi$ PVL. ORF JP058 encodes 48 amino acids with no assigned function identified by means of a BLASTP search. This ORF was also identified in  $\phi$ PVL, although it was disrupted and its function could not be assigned because of the insertion of a "T" nucleotide.

Group 2 contained 27 ORFs (JP011, JP012, JP028, and JP032 to -055) conserved among  $\phi$ Sa2958,  $\phi$ Sa2mw, and  $\phi$ SLT (Fig. 1). Because eight ORFs that might be associated with phage DNA packaging and head and tail formation are involved (JP034 [terminase small subunit], JP035 [terminase large subunit], JP036 [portal protein], JP037 [prohead protease], JP038 [capsid protein], JP043 and JP044 [major tail protein], and JP047 [phage tail tape measure protein]),  $\phi$ Sa2958 could be determined on the basis of taxonomy criteria as belonging to the same subgroup as  $\phi$ Sa2mw and  $\phi$ SLT.



TABLE 2. ORFs in and around  $\phi$ 2958PVL and their similarities to  $\phi$ Sa2mw and  $\phi$ 108PVL<sup>a</sup>

ORF and special structure	Start (bp)	End (bp)	Size (bp)	aa <sup>a</sup>	Gene	Function	$\phi$ Sa2mw			$\phi$ 108PVL		
							% Identity to $\phi$ Sa2mw genome	Corresponding ORF (size in bp)	% aa identity	% Identity to $\phi$ 108PVL genome	Corresponding ORF (size in bp)	% aa identity
(JP001) <sup>b</sup>	295	936	642	214		Hypothetical protein						
<i>attB1</i>	842	866	25			Attachment site on chromosome						
Core	867	895	29			Core sequence						
<i>attP</i> -left	896	920	25			Attachment site on phage						
JP002	2184	979	1,206	401	<i>int</i>	Phage integrase	98.7	Integrase (1,206)	98.6	Integrase (1,206)	99.8	
JP003	2310	2924	615	204		Na/K-ATPase	47.4	Putative membrane protein	52.1		26.7	
JP004	3285	3103	183	60		Hypothetical protein	52.2	MW1441 (624)	52.2		45	
JP005	4301	3381	921	306		Hypothetical protein	45.9		45			
JP006	4931	4317	615	204	<i>rep</i>	Putative transcriptional repressor	45.2	Repressor (324)	49.1	cf-like repressor (717)	34.3	24
JP007	5103	5330	228	75		Hypothetical protein	54.4		54.7	Putative cro-like repressor (243)	23.1	
JP008	5356	5520	165	54	<i>ant</i>	Putative antirepressor	52.1		50.3	Putative antirepressor (753)	31.4	
JP009	5927	6145	219	72		Hypothetical protein	51.8		52			
JP010	6195	6440	246	81		Hypothetical protein	100	MW1434 (246)	47.9		100	
JP011	6774	6409	366	121		Hypothetical protein	100	MW1433 (366)	46.1		100	
JP012	6829	7044	216	71		Hypothetical protein	100	MW1432 (216)	55.6		66.2	
JP013	7069	7332	264	87		Hypothetical protein	82.6	MW1431 (264)	99.6		78.8	
JP014	7347	7505	159	52		Hypothetical protein	93.7	MW1430 (162)	98.7		96.2	
JP015	7584	7907	324	107		Hypothetical protein	92.9	MW1429 (324)	91.6		51	
JP016	7922	8284	363	120		Hypothetical protein	97.8	MW1428 (363)	48.9		97.5	
JP017	8281	9447	1,167	388		Hypothetical protein	97.4	MW1427 (1,167)	46.5		97.9	
JP018	9473	10030	558	185		Hypothetical protein	98.7	MW1426 (558)	49.7		99.5	
JP019	10098	12050	1,953	650	<i>pol</i>	DNA polymerase	99.8	DNA polymerase (1,962)	99.7		99.7	
JP020	12063	12248	186	61		Hypothetical protein	100	MW1424 (186)	69.9		100	
JP021	12245	12649	405	134		Putative DNA-binding protein	100	MW1423 (402)	57		100	
JP022	12649	12906	258	85		Hypothetical protein	97.7	MW1422 (258)	52.8		97.6	
JP023	13118	13366	249	82		Hypothetical protein	92.4	MW1420 (243)	91.6		93.8	
JP024	13407	13661	255	84		Hypothetical protein	89.8	MW1419 (255)	96.8	P028 (249)	89	
JP025	13811	14347	537	178		Hypothetical protein	95.2	MW1415 (534)	49.1	P030 (372)	98.8	
JP026	14645	14881	237	78		Hypothetical protein	54		100	P032 (237)	100	
JP027	14865	15026	162	53		Hypothetical protein	91.7	MW1411 (156)	92.1		86.3	
JP028	15094	15294	201	66		Hypothetical protein	98.5	MW1410 (201)	61.6		97	
JP029	15347	17794	2,448	815	<i>virE</i>	Virulence-associated protein E	98.9	Virulence-associated protein E (2,448)	46.7		99.6	
JP030	18135	18425	291	96		Hypothetical protein	95.2	MW1406 (198)	50		98.5	
JP031	18502	19773	1,272	423		Phage helicase	99.4	Phage helicase (1,359)	47.8		100	
JP032	19786	20223	438	145		Phage regulatory protein	89.7	MW1404 (438)	51.4		93.8	
JP033	20380	20694	315	105		Hypothetical protein	97.5	MW1403 (315)	51.5		95.2	
JP034	20805	21128	324	107	<i>terS</i>	Terminase small subunit	97.5	Terminase small subunit (306)	49.4	Terminase small subunit (468)	28.6	
JP035	21448	22809	1,362	453	<i>terL</i>	Terminase large subunit	91.6	Terminase large subunit (1,692)	46.1	Terminase large subunit (1,695)	24.8	
JP036	23006	24052	1,047	348	<i>por</i>	Portal protein	98.9	Portal protein (1,239)	48.7	Portal protein (1,326)	21.9	
JP037	24036	24809	774	257	<i>pro</i>	Prohead protease	99.7	Protease (774)	46.7	Prohead protease (585)	24.6	
JP038	24776	25984	1,209	402		Putative capsid protein	99.9	Capsid (1,164)	49.3	Capsid (1,248)	19.2	
JP039	26053	26331	279	92		Hypothetical protein	100	MW1397 (279)	49.4		100	
JP040	26343	26675	333	110		Hypothetical protein	96.4	MW1396 (333)	49.5		97.3	
JP041	26672	27073	402	133		Hypothetical protein	99.5	MW1395 (402)	48.5		100	
JP042	27074	27469	396	131		Hypothetical protein	100	MW1394 (396)	50.4		100	
JP043	27504	28145	642	213		Major tail protein	99.8	Major tail protein (642)	47.8		100	
JP044	28237	28692	456	151		Major tail protein	99.8	Major tail protein (456)	49.3		99.3	
JP045	28750	29100	351	116		Hypothetical protein	99.7	MW1391 (342)	50.4		100	
JP046	29142	29300	159	52		Hypothetical protein	100	No corresponding gene	56.8		56.8	
JP047	29314	35514	6,201	2066	<i>mip</i>	Tail length tape measure protein	98.1	Tail length tape measure protein (6,201)	46.4	Tail length tape measure protein (3,333)	25.3	
JP048	35514	36338	825	274		Hypothetical protein	100	MW1389 (825)	48.5		100	
JP049	36347	37930	1,584	527		Hypothetical protein	100	MW1388 (1,584)	45.7		100	
JP050	37930	38220	291	96		Hypothetical protein	100	MW1387 (291)	49.5		100	

JP051	38236	40146	1,911	636	Hypothetical protein	99.6	MW1386 (1,911)	99.8	45.4
JP052	40113	41612	1,500	499	Hypothetical protein	99.7	MW1385 (1,467)	99.8	47.5
JP053	41612	42001	390	129	Hypothetical protein	100	MW1384 (390)	100	48.1
JP054	41994	42158	165	54	Hypothetical protein	100	MW1383 (165)	100	54
JP055	42204	42503	300	99	Hypothetical protein	100	MW1382 (300)	100	51.1
JP056	42639	42941	303	100	Holin	100	Holin (303)	100	100
JP057	42952	44406	1,455	484	Amidase (peptidoglycan hydrolase)	98.3	Amidase (1,455)	99.2	99.2
JP058	44556	44687	132	43	Hypothetical protein	98.5	No corresponding gene	98.5	
JP059	44795	45733	939	315	<i>lukS-PV</i>	99.8	LukS-PV (939)	100	100
JP060	45735	46712	978	325	<i>lukF-PV</i>	99.9	LukF-PV (978)	99.9	100
<i>attP-right</i>	46917	46941	25		Attachment site on phage				
Core	46942	46970	29		Core sequence				
<i>attB2</i>	46971	46995	25		Attachment site on chromosome				

<sup>a</sup> aa, amino acid.

<sup>b</sup> ORF in parentheses was located outside of the  $\phi$ 2958PVL. Two sets of core sequences and attachment site sequences were also identified flanking the  $\phi$ 2958PVL.




FIG. 2. Representative results of seven multiplex PCRs (M-PCRs) for identifying PVL phages. A 1-kb molecular weight marker (MWM) was run on both sides of the gel. Lanes: 1, PCR-1 using chromosomal DNA of 81/108; 2, PCR-2 using chromosomal DNA of JCSC2958; 3, PCR-3 using chromosomal DNA of 81/108; 4, PCR-4 using chromosomal DNA of JCSC2958; 5, PCR-5 using chromosomal DNA of 81/108; 6, PCR-5 using chromosomal DNA of ATCC 49775; 7, PCR-6 using chromosomal DNA of JCSC2958; 8, PCR-7 using chromosomal DNA of MW2; and 9, PCR-8 using chromosomal DNA of RN4220 ( $\phi$ SLT).

Group 3 contained 12 ORFs (JP015 to JP022, JP025, JP029, JP030, and JP031) that showed high similarity only with  $\phi$ Sa2mw (Fig. 1). Three genes were associated with phage replication during the phage lytic process. Representative genes in group 3 encode DNA polymerase (JP019), DNA-binding protein (JP021), helices (JP031), and virulence-associated protein E (JP029).

Group 4 contained three ORFs that were homologous only to one or two other phages. JP026, JP003, and JP010 (Fig. 1) were homologous only to  $\phi$ 108PVL and  $\phi$ SLT, only to  $\phi$ PVL and  $\phi$ SLT, and only to  $\phi$ 108PVL, respectively.

Group 5 contained six ORFs (JP004 to -009) that were uniquely conserved in  $\phi$ Sa2958 (Fig. 1). Interestingly, JP006, containing the transcription repressor gene, was not homologous to the corresponding genes in other extant PVL phages. A search with the BLASTP program revealed that JP006 showed 100% identity with the corresponding gene conserved in lytic phage 47 and showed 65% identity with that in lytic phage 29, whose entire genomic sequences have been determined by Kwan et al. (23).

The JP006 protein contains a helix-turn-helix DNA-binding motif that is present in a large family of transcriptional regulators. Another unique ORF, JP008, which is conserved in  $\phi$ Sa2958 and was identified as an antirepressor gene, was shown to have 93% similarity with the corresponding gene in phage 47. The remaining four unique ORFs in  $\phi$ Sa2958 could not be assigned any functions through searches with the BLASTP program.

**Characterization of PVL-carrying prophages in Japanese MRSA strains.** We conducted PCR experiments to identify PVL phages carried by 53 MRSA strains isolated in from 1979 through 1985 and 12 MRSA strains isolated in the 2000s with PCRs as described in Materials and Methods. The representative results of these PCRs are shown in Fig. 2 and the results of PCR experimentation are summarized in Table 3. When PCR-1 was performed with the chromosomal DNAs of these same five strains carrying each PVL phage, DNA fragments of

TABLE 3. Results of PCR experimentation<sup>a</sup>

Yr of isolation (no. of strains)	Identification of (no. of strains):					
	Two kinds of phages by:		Gene lineage of <i>lukS-PV</i> to tail gene by:		Phage-specific modules related to lysogeny and recombination by PCR-5, -6, and -7	PVL-carrying phage type (no. of strains)
	PCR-1 (icosahedrally shaped head)	PCR-2 (elongated head)	PCR-3 (tail gene common to $\phi$ 108PVL and $\phi$ PVL)	PCR-4 (tail gene common to $\phi$ Sa2mw, $\phi$ SLT, and $\phi$ Sa2958)		
1979–1985 (53)	+ (39)	+ (39)	+ (35)	ND	$\phi$ 108PVL (33) NT (2)	$\phi$ 108PVL type (33) Icosahedral-head type (2)
	– (14)	+ (14)	– (4) ND	ND + (1)	$\phi$ 108PVL (4) $\phi$ Sa 2958 (1)	$\phi$ 108PVL-like type (4) $\phi$ Sa2958 type (1)
2000s (12)	– (12)	+ (11)	ND	– (13) + (10)	$\phi$ Sa 2958 (13) $\phi$ Sa 2958 (2) NT (8)	$\phi$ Sa2958-like phage of unknown type (13) $\phi$ Sa2958 type (2) Elongated-head type (8)
		– (1)	ND	– (1) – (1)	$\phi$ Sa 2958 (1) $\phi$ Sa 2958 (1)	$\phi$ Sa2958-like phage of unknown type (1) NT (1)

<sup>a</sup> ND, not determined; NT, nontypeable.

two expected sizes, 489 bp for the tail gene and 569 bp for the head gene, were amplified with chromosomal DNAs of 81/108 ( $\phi$ 108PVL), ATCC 49775 ( $\phi$ PVL), and MW2 ( $\phi$ Sa2mw). When PCR-2 was performed with chromosomal DNAs of these same five strains, two DNA fragments of expected sizes, 466 bp for the tail gene and 655 bp for the head gene, were amplified with chromosomal DNAs of JCSC2958 ( $\phi$ Sa2958), RN4220 ( $\phi$ SLT), MW2 ( $\phi$ Sa2mw), and 81/108 ( $\phi$ 108PVL).

MW2 was positive by PCR-1 and PCR-2, because this *S. aureus* strain carries two additional different phages in its chromosomal DNA, and one of them,  $\phi$ Sa3mw, has the same morphogenesis region as does  $\phi$ PVL (1, 19). Strain 81/108 was also positive with PCR-2 and seemed to carry other phages that did not carry *lukS-PV* and *lukF-PV* genes or have the structure to react with primers for PCR-2. Most of the PCR-1-positive strains we had tested up to this point in our study were also PCR-2 positive.

Therefore, to confirm whether the structures related to morphology belonged to PVL-carrying phages, we performed PCR-3 and PCR-4 to verify whether the PVL-carrying prophage belonged to either the icosahedral-head-type phages or the elongated-head-type phages by amplifying gene lineages between *lukS* and *mtp* that belong to either the icosahedral-head type or the elongated-head type.

DNA fragments of the expected sizes were amplified using PCR-3 and chromosomal DNAs of 81/108 ( $\phi$ 108PVL) and ATCC 49775 ( $\phi$ PVL) and using PCR-4 and chromosomal DNAs of MW2 ( $\phi$ Sa2mw), JCSC2958 ( $\phi$ Sa2958), and RN4220 ( $\phi$ SLT).

Up to that point, 38 of 39 PCR-1-positive strains were positive by PCR-2. However, PCR-3 was performed with chromosomal DNAs of 39 strains, and a DNA fragment of 10.5 kb was successfully amplified from 35 of 39 PCR-1-positive strains, indicating that these strains carried PVL phages of the icosahedral-head type. Sixty-six of 67 strains were positive by PCR-2, whereas DNA fragments of different sizes were amplified with chromosomal DNA of JCSC4274. When we performed PCR-4 with chromosomal DNAs with 26 strains for which no or only one DNA fragment was amplified by PCR-1, DNA fragments were amplified with chromosomal DNAs of 11 of these 26 strains.

Furthermore, we performed PCR-5 to -8 to classify these PVL phages into five different types. We conducted PCR-5 to determine whether these phages belonged to either the  $\phi$ PVL type or the  $\phi$ 108PVL type. A DNA fragment of 4,340 bp, indicating the carriage of a  $\phi$ 108PVL-specific gene (fragment PCR-1 A), was amplified with the DNAs of 33 of 35 strains, whereas a DNA fragment of 1,411 bp, indicating the carriage of  $\phi$ PVL (fragment PCR-2 A), was amplified in two cases, together with a DNA fragment of 4,340 bp. Because two PCR experiments to identify the other component of  $\phi$ PVL yielded negative results and four PCR experiments to identify the  $\phi$ 108PVL-specific regions yielded positive results (data not shown), we conclude that these 33 strains that were positive by PCR-1 amplified fragment PCR-1 A, which carried the phage belonging  $\phi$ 108PVL type. Four strains that were negative by PCR-4 were confirmed to belong to the  $\phi$ 108PVL type, with four PCR experiments to identify  $\phi$ 108PVL-specific regions and two long-range PCRs to amplify the DNA fragments covering the region from *mtp* to *lukS*. Therefore we regarded these four strains as being of the  $\phi$ 108PVL-like type. Two strains, with the gene lineage between *lukS* and *mtp* common to  $\phi$ 108PVL and  $\phi$ PVL, could not be classified into one of the five extant PVL phage types tested for up to that point.

In contrast, of the 26 strains for which no or only one DNA fragment was amplified by PCR-1, 25 were positive by PCR-2. In the case of strain JCSC7247, only one DNA fragment was amplified, suggesting the carriage of *mtp* in common with the elongated-head type.

PCR-4 was performed with DNAs of 26 strains to verify the gene lineage between the tail gene and the *lukS* gene by use of a primer pair designed for the tail gene that is commonly conserved in  $\phi$ Sa2958,  $\phi$ Sa2mw, and  $\phi$ SLT and for *lukS*. Of the 14 strains isolated from 1979 through 1985, only 1 was positive by PCR; in contrast, 10 of 20 strains isolated in the 2000s were positive by PCR. In addition, we performed PCR-6, -7, and -8 to classify elongated-head-type phages as of either the  $\phi$ Sa2958 type, the  $\phi$ Sa2mw type, or the  $\phi$ SLT type. Interestingly, neither the  $\phi$ Sa2mw type nor the  $\phi$ SLT type, both of which are carried by *S. aureus* strains isolated in the United States or France, was identified, whereas three strains carrying  $\phi$ Sa2958-type phage and eight strains carrying the elongated-

TABLE 4. Characterization of PVL-positive MRSA strains isolated in 1979 to 1985, 2002, and 2004

Yr of isolation (no. of strains)	Phage type <sup>a</sup> (no. of strains)	Coagulase isotype	SCCmec type (no. of strains)	Strain genotype by MLST				
				No. of tested strains	ST	Allelic profile		
1979–1985 (53)	φ108PVL (33)	4	I (1)	1	30	2, 2, 2, 2, 6, 3, 2		
			II (2)	1	30	2, 2, 2, 2, 6, 3, 2		
			IV.1 (8)	3	30	2, 2, 2, 2, 6, 3, 2		
				2	N <sup>b</sup>	2, 2, N, 2, 6, 3, 2		
			IV.3 (21)	3	30	2, 2, 2, 2, 6, 3, 2		
			IV.5 (1)	1	30	2, 2, 2, 2, 6, 3, 2		
			φ108PVL-like (4)	4	IV.1 (1)	1	30	2, 2, 2, 2, 6, 3, 2
					IV.3 (3)	1	30	2, 2, 2, 2, 6, 3, 2
			Icosahedral-head type (2)	4	IV.3 (1)	1	30	2, 2, 2, 2, 6, 3, 2
					type II.n (1)	1	30	2, 2, 2, 2, 6, 3, 2
			φSa2958 type (1)	2	II (1)	1	5	1, 4, 1, 4, 12, 1, 10
			Sa2958-like phage of unknown type (13)	4	I (10)	2	30	2, 2, 2, 2, 6, 3, 2
					II (1)	1	30	2, 2, 2, 2, 6, 3, 2
IV.3 (1)	1	30			2, 2, 2, 2, 6, 3, 2			
IV.n (1)	1	30			2, 2, 2, 2, 6, 3, 2			
IV.3 (2)	2	30			2, 2, 2, 2, 6, 3, 2			
2000s (12)	φSa2958 (2)	4	IV.3 (1)	1	30	2, 2, 2, 2, 6, 3, 2		
			NT <sup>c</sup> (6)	1	30	2, 2, 2, 2, 6, 3, 2		
Elongated-head type (8)	4	I (1)	1	765	2, 2, 2, 2, 6, 104, 2			
		V (1)	1	59	19, 23, 15, 2, 19, 20, 15			
		Nontypeable (1)	7					

<sup>a</sup> Phage type judged with M-PCRs.

<sup>b</sup> N, the ST of this strain could not be assigned, since the *glpF* gene could not be amplified by PCR. The allele numbers other than *glpF* were the same as those for ST30.

<sup>c</sup> NT, nontypeable.

head-type phage were identified. We determined the nucleotide sequence of a prophage lysogenized in MRSA strain JCSC2958, which was isolated in Japan in 1981. However, 13 of 14 strains isolated from 1979 through 1985 were definitively determined to be of the φSa2958 type, because we could not amplify the region between *mtp* and *lukS* with long-range PCR. Because these 13 strains as well as a strain isolated in the 2000s were positive by PCR-6, identifying a specific ORF in φSa2958, and by several types of long-range PCR, amplifying the gene lineage between *mtp* to some ORFs of φSa2958 located between the tail gene and *lukS*, e.g., JP052 or JP053, we consider them to be φSa2958-like phage of an unknown type. One strain carried a prophage that could not be classified into either one of the two types; this strain was exceptional, because it was negative by the seven PCRs other than PCR-6.

**Characterization of PVL-positive MRSA clones.** We next characterized these strains by determining the coagulase isotypes and SCCmec types of all 65 strains and the MLST types of selected strains (Table 4). Fifty-two of the 53 MRSA strains isolated from 1979 through 1985 and 11 of the 12 MRSA strains isolated in the 2000s carried type 4 coagulase. Only JCSC2958, which we used for sequencing the φ2958PVL genome, carried type 2 coagulase, and only JCSC7247 carried type 7 coagulase. When we performed MLST of the chosen isolates, we found that all coagulase type 4 strains belonged to ST30, whereas a coagulase type 2 strain belonged to ST5 and a coagulase type 7 strain belonged to ST59. The data showed that the genotypes of MRSA strains carrying PVL phages isolated in Japan were extremely homogeneous and were represented by ST30 coagulase type 4 strains.

However, the SCCmec elements carried by these strains were extremely diverse. Among the 33 MRSA strains identified as of the φ108PVL type, SCCmec type IV.3 strains (21 of

33) were the most frequent and were followed by SCCmec type IV.1 strains (8 of 33), SCCmec type II strains (2 of 33), and SCCmec type I strains (1 of 33). Three strains identified as being of the φSa2958 type carried type II SCCmec and type IV.1 SCCmec elements. Eight strains classified as the elongated-head type carried type IV.3 SCCmec and nontypeable SCCmec elements, which were similar to those of type IV.3 because they carried type 2 *ccr* and the J1 region of type IV.3 SCCmec and *IS1272*. Since we could not amplify the gene lineage between *mecA* and *IS1272*, we tentatively considered them as untypeable. Other strains also carried several SCCmec elements.

Interestingly, no strains with icosahedral-head-type phages were identified in isolates from the 2000s. The prophages of three strains were judged to be of the φSa2958 type. In contrast to JCSC2958, two other strains isolated in the 2000s belonged to the coagulase type 4 ST30 genotype and carried a type IV.3 SCCmec element. A φSa2958-like phage of unknown type was identified in the 2000s as belonging to the coagulase type 4 ST765 (CC30) group, carried type I SCCmec, and was similar to the isolates from 1979 through 1985. A strain carrying an untypeable PVL-carrying phage belonged to coagulase type 7 ST59 and carried the type V SCCmec element.

**Induction of infective PVL phages.** To date, the φSLT is the only PVL phage that has been induced from cells and found capable of infecting other cells. Through a structural comparison of PVL phages, we found that both φSa2958 and φSa2mw carried possibly intact genes encoding head and tail proteins, and we inferred that these PVL phages might also be infective. To confirm this hypothesis, we induced prophages by treating these phages with mitomycin C and mixed them with cells of restriction-negative strains RN4220 and 1039. Because plaque formation does not indicate whether these plaques were generated by in-



TABLE 5. Identification of PVL phages induced by mitomycin C treatment

Strain	Coagulase isotype	MLST	SCC <i>mec</i> type	Phage type <sup>a</sup>	Indicator strain	No. of:		
						Phages induced by mitomycin C (PFU/ml)	Tested plaques	PVL-positive plaques (%)
MW2	7	1	IV.1	φSa2mw (S)	RN4220	2.34 × 10 <sup>6</sup>	234	234 (100)
					1039	3.86 × 10 <sup>6</sup>	386	386 (100)
JCSC2958	2	5	II	φSa2958 (S)	RN4220	2.26 × 10 <sup>9</sup>	226	226 (100)
					1039	1.05 × 10 <sup>10</sup>	316	316 (100)
RN4220/φSLT	3	8	— <sup>b</sup>	φSLT (S)	RN4220	1.11 × 10 <sup>6</sup>	111	111 (100)
					1039	2.23 × 10 <sup>5</sup>	223	223 (100)
81/108	4	30	IV.3	φ108PVL (S)	RN4220	4.2 × 10 <sup>2</sup>	42	0
					1039	1.82 × 10 <sup>5</sup>	182	0
JCSC2973	4	30	I	φSa2958-like (P)	RN4220	2.32 × 10 <sup>5</sup>	232	0
					1039	4.08 × 10 <sup>6</sup>	408	2 (0.49)
M11	4	30	IV.3	φSa2958-like (P)	RN4220	2.05 × 10 <sup>8</sup>	205	205 (100)
					1039	2.33 × 10 <sup>8</sup>	233	233 (100)
JCSC4465	4	30	IV.n	φSa2958-like (P)	RN4220	1.34 × 10 <sup>8</sup>	134	0
					1039	1.48 × 10 <sup>8</sup>	148	4 (2.7)
JCSC4625	4	30	IV.3	φSa2958 type (P)	RN4220	2.61 × 10 <sup>5</sup>	261	261 (100)
					1039	3.97 × 10 <sup>6</sup>	397	397 (100)
JCSC6055	4	30	IV.3	φSa2958 type (P)	RN4220	No plaque	NT <sup>c</sup>	NT
					1039	No plaque	NT	NT

<sup>a</sup> Phage types assigned with PCR are indicated by (P), and those judged by sequences of phage genomes are indicated by (S).

<sup>b</sup> —, none.

<sup>c</sup> NT, nontypeable.

duced PVL phages, we conducted plaque hybridization experiments with a probe for the *lukS-PV* and *lukF-PV* genes to examine whether these plaques were of PVL-carrying phages.

The results are summarized in Table 5. We tested four MRSA strains (JCSC2973, JCSC4465, M11, and JCSC2958) isolated from 1979 through 1985, along with MRSA strains isolated in 2002 (JCSC4625) and 2004 (JCSC6605). Three strains carried φSa2958-type PVL phages, and three strains carried φSa2958-like PVL phages of unknown type. In addition, prophages φSa2mw and φ108PVL, whose genomes have been sequenced, were induced from strains MW2 and 81/108, as was lysogenized φSLT from strain RN4220 (φSLT). The cell suspensions of eight of nine tested strains became transparent after the addition of mitomycin C, indicating that phages might be generated upon the stress of DNA synthesis inhibition by mitomycin C. When these phage lysates were diluted and mixed with cells of indicator strains RN4220 and 1039, we found that most of the induced phages could propagate in 1039, whereas only some of them could propagate in RN4220.

As expected, phage lysates from MW2 and JCSC2958 generated plaques on the 1039 and RN4220 lawns, which hybridized with the probe for the *lukS-PV* and *lukF-PV* genes, indicating that φSa2mw and φSa2958 can infect other *S. aureus* strains, as can the previously reported φSLT. The phage lysate from 81/108 generated plaques, but no PVL-positive plaque was observed, indicating that the φ108PVL phage did not induce or cannot infect the strains tested so far. This result was consistent with our speculation that φ108PVL might be defective, because the tail gene of φ108PVL was disrupted in a manner similar to that seen for φPVL (21).

Phage lysates of five of six MRSA strains carrying either the φSa2958-type PVL phage or the φSa2958-like PVL phage of unknown type generated PVL-positive plaques on indicator strains; however, the ratios of PVL-positive plaques ranged from 2% to 100%. In the case of JCSC6055, whose cell sus-

pension did not become transparent after the addition of mitomycin C, no plaques were formed on the lawns of 1039 and RN4220 after the addition of the filtrate.

## DISCUSSION

**Characteristics of φSa2958.** Canchaya et al. have classified the *Staphylococcal* phages into five groups based on similarities in genes related to morphogenesis (4). All extant PVL phages are Sfi21-like *cos*-site phages of the *Siphoviridae* family, which can be further subdivided into two groups based on morphology: those with isomeric hexagonal heads and those with elongated heads. The PVL phages φSa2mw and φSa2usa, which are carried by MRSA strains isolated in the United States, have elongated heads. The φSLT phage was isolated from a French strain of *S. aureus* and has been shown by electron microscopy to have an elongated head measuring 100 by 50 nm and a flexible tail that is 400 nm long (32). Interestingly, the φSa2958 phage we described here belongs to the elongated-head group, whereas φ108PVL, identified in Japan, has an isometric hexagonal head, as does φPVL (20).

When we performed a detailed comparison of the genomic structure of φSa2958 with those of other extant PVL phages, represented by φSa2mw and φ108PVL (Fig. 3 and Table 2), we found that most of the region between φSa2958 and φSa2mw is conserved. Similarly, this region is also conserved among φSa2usa and φSLT. In contrast, the regions containing ORFs related to lysogeny (ORFs JP004 to -009) were unique to each phage. The average G+C content of the region from JP004 to JP009 was approximately 29.2%, which was lower than the average of the whole φSa2958 genome (33.1%). These data indicated that these genes might be acquired from organisms with lower G+C contents.

The theory of modular exchange in phages, which is known as the modular theory, has become the basis of a popular




FIG. 3. Structural comparison of 5 PVL phages. Structures of  $\phi$ 108PVL,  $\phi$ PVL,  $\phi$ SLT,  $\phi$ Sa2mw, and  $\phi$ Sa2958 are illustrated on the bases of the nucleotide sequences as follows:  $\phi$ 108PVL (DDBJ/EMBL/GenBank databases under accession no. AB243556),  $\phi$ PVL (DDBJ/EMBL/GenBank accession no. AB009866),  $\phi$ SLT (DDBJ/EMBL/GenBank accession no. NC\_002661),  $\phi$ Sa2mw (DDBJ/EMBL/GenBank accession no. BA000033), and  $\phi$ Sa2958 (DDBJ/EMBL/GenBank accession no. AP009363). Genes having nucleotide identities of more than 90% are linked by blocks filled with colors as follows: pink, genes conserved among five PVL phages; light purple, genes conserved among  $\phi$ Sa2958,  $\phi$ Sa2mw, and  $\phi$ SLT; blue, genes conserved between  $\phi$ PVL and  $\phi$ 108PVL. The positions of primers for M-PCR 1 to 7 are indicated by green arrowheads. Blue arrows indicate ORFs well conserved in all PVL phages other than *lukS* and *lukF* genes; black arrows indicate *lukS* and *lukF* genes; light green and dark green arrows indicate genes encoding major tail proteins of the elongated-head type and the icosahedral-head type, respectively; and yellow arrows and other arrows indicate genes encoding terminase subunits (large and small), portal proteins, prohead proteases, and capsid proteins of the elongated-head type and the icosahedral-head type, respectively.

hypothesis for phage evolution (26, 33). According to this theory,  $\phi$ Sa2958 and  $\phi$ Sa2mw must share a common ancestor and must have evolved as close relatives, while the module exchange was likely a recent event. We note here that functionally important structures of the repressor and anti-repressor genes are located in this module (25). We now know that in most known prophages, immunity is elicited by a repressor protein that prevents transcription initiation at promoters controlling the expression of lytic function (14, 24). Furthermore, the immunity of the phage is sometimes complicated by the presence of an anti-repressor gene that can prevent the expression of the repressor activity. The nucleotide sequences of the repressor genes conserved in  $\phi$ Sa2958 and  $\phi$ Sa2mw exhibited differences from each other at the DNA level; however, when we investigated the protein structure encoded by these repressor genes, we found that the essential helix-turn-helix domain was well conserved, indicating that both repressor genes execute functional activities. This finding suggests that  $\phi$ Sa2958 and  $\phi$ Sa2mw might utilize different mechanisms for maintain-

ing their immunity when they are integrated into bacterial genomes; to achieve this goal, these phages had to adopt the most beneficial module during the process when they interacted with host bacteria or with other phages.

**Two lineages of PVL phage are predominant in MRSA and MSSA strains in Japan.** We designed several PCRs to identify PVL phages with the notion that the type of PVL-carrying phage should be determined on the bases of the gene lineage between *lukS-PV* and *lukF-PV* and other phage components. We first developed PCR-1 and -2 to identify portal or head genes, but we soon noticed that these PCRs might also be used to identify components of phages other than PVL-carrying phages. Therefore, we developed PCRs to identify the gene lineage between *lukS* and *mtp*, which is located rather far from the *lukS-PV* and *lukF-PV* genes. In addition, we developed PCRs to classify individual phages by identifying the genes in lysogeny-related regions or recombination-related regions in combination with the integrase gene, which is carried by all reported PVL-carrying phages. During this step, we noticed

that some modules are commonly shared among phages. As shown in Fig. 3,  $\phi$ Sa2958 was similar to  $\phi$ Sa2mw, and  $\phi$ Sa2958 was similar to  $\phi$ SLT (data not shown). The primer pairs we reported here are only those selected pairs that react only with one of five phages.

We have found that two PVL lineages, the icosahedral-head type and the elongated-head type, are present in Japan. Phages of the  $\phi$ 108PVL type were identified mostly in MRSA strains isolated from 1979 through 1985. Of the 53 MRSA strains examined, 33 isolates were identified as the carrier of PVL-carrying phage of the  $\phi$ 108PVL type, and 4 isolates were identified as carriers of the  $\phi$ 108PVL-like type; however, no phages of this type were identified in the isolates from the 2000s. Furthermore, no  $\phi$ PVL-type strain was identified among the MRSA strains tested. However, when 13 PVL-positive MSSA strains isolated in the 2000s were examined, 2 were found to belong to the icosahedral-head type and were likely to be of the  $\phi$ PVL type. Because the tail gene of  $\phi$ 108PVL was truncated, we assume that the phage had already lost its infectivity but carried an integrated truncated prophage, which might explain why  $\phi$ 108PVL-type strains have not been identified recently.

In contrast, phages with an elongated head, e.g.,  $\phi$ Sa2958 as well as  $\phi$ Sa2mw and  $\phi$ SLT, carried intact tail and head genes.  $\phi$ Sa2958 was identified from an ST5 *SCCmec* type II MRSA strain identified in 1981, whereas two  $\phi$ Sa2958 type strains of ST30 *SCCmec* type IV.3 strains were identified in the 2000s, suggesting that the phage might infect other strains and become integrated into their chromosomes as a prophage. We considered the prophages carried by 14 strains (13 isolates from 1975 to 1985 and 1 isolate from the 2000s) as  $\phi$ Sa2958-like phages of unknown type, because we could not clarify the gene lineage between *mtp* and *lukS-PV*. The strains carrying the prophage might be derived from the same clone because they belonged to ST30 and carry a type I *SCCmec* element and because the prophages of all three tested strains could be induced, proving their infectivity; these findings suggest that the recent isolate might be a descendant or that the phage might infect other strains. Interestingly, eight strains with elongated-head-type PVL-carrying prophages were identified. Because these strains were identified not only from seven isolates in Wakayama prefecture but also from an isolate from a 27-year-old man in Kanagawa prefecture, which is far from Wakayama, in 2002 (strain EB00449), we speculate that the prophage might have been induced from the cells of *S. aureus* and then infected other *S. aureus* cells to generate the novel PVL phage-carrying strain. A coagulase type 7 ST59 strain carried an as-yet-untypeable PVL phage. Vandenesch et al. reported that five major STs, namely, ST1, -8, -30, -59, and -80, carry PVL phages and that ST59 PVL-positive strains have been isolated in the United States (39). Because the ST59 strain is not common in Japan, the strains of other countries might have been introduced into Japan. Whether the strain carries a novel PVL phage is the next problem that should be clarified. It might carry a PVL phage that does not belong to the two extant groups.

When we characterized these PVL-positive strains by means of *SCCmec* typing and genotyping, we found that the most of them produced type 4 coagulase and belonged to ST30 or CC30 and carried *SCCmec* elements of various types, including type I, type II, type IV.1, and type IV.3 (17, 18, 29). We

speculate that MSSA strains of coagulase type 4 ST30 independently acquired two phages,  $\phi$ 108PVL and  $\phi$ Sa2958, or other elongated-head phages before they acquired *SCCmec* elements. The characterization of PVL phages carried by MSSA strains isolated in the 1960s has shown that MSSA strains carrying either of the PVL phages were present in the Japanese community. Therefore, we presume that the MSSA strains evolved into PVL-positive MRSA strains by acquiring *SCCmec* elements of various types. When we examined the types of *SCCmec* elements carried by PVL-positive MRSA strains, we found that 20 of 33 MRSA strains of the  $\phi$ 108PVL type carried type IV.3 *SCCmec* and that 8 of 33 carried type IV.1 *SCCmec*. It was noteworthy that most of the PVL-positive *SCCmec* type IV.1 strains had been isolated at the Tokyo Geriatric Hospital, whereas, and most of the PVL-positive *SCCmec* type IV.3 strains had been isolated at Gunma University Hospital. The data indicated that MRSA clones disseminating in different Japanese hospitals were not identical and that the spread of MRSA strains is regional.

**$\phi$ Sa2958 and  $\phi$ Sa2958-like phages of unknown type are intact prophages.** When we studied the structure of  $\phi$ Sa2958, identified in this study, we realized that the amino acid sequence of the tail-length tape measure protein designated JP047 showed 98% similarity in with that of  $\phi$ SLT, which has been shown to be the first PVL phage that could infect *S. aureus* strains experimentally (32). Because the tail-length tape measure protein of  $\phi$ Sa2mw carried by MW2, a CA-MRSA strain isolated in North Dakota, was similar to that of  $\phi$ SLT, we presumed that the newly identified  $\phi$ Sa2958 as well as  $\phi$ Sa2mw might also be able to infect to *S. aureus* strains.

To determine whether these two phages as well as phages of the  $\phi$ Sa2958 type were intact and could infect *S. aureus* strains, we tested MW2 and six MRSA strains carrying  $\phi$ Sa2958-type phage or  $\phi$ Sa2958-like PVL phage of unknown type. The PVL phages carried by five of six MRSA strains could infect indicator cells, although the ratios of PVL phages in the phage lysates were not similar. In addition, the sizes of plaques generated on the lawn were larger with 1039 than with RN4220. With three strains, namely, M11, JCSC2958, and JCSC4625, all phages induced by treatment with mitomycin C were PVL positive. In contrast, with two strains, JCSC2973 and JCSC4625, the ratios of PVL-positive phages were very low. In the case of JCSC6055, which did not generate any plaques, the apparent reason that phages were not induced was that the cell suspension did not turn transparent after mitomycin C was added. These discrepancies may be due to many host cell factors, e.g., the presence of other prophages and differences in gene expression that confer the SOS response. Although we did not test a large number of strains, we chose  $\phi$ Sa2958-type MRSA strains of a representative genotype identified in this study. Except for JCSC6055, the five strains (derived from four different clones as indicated), namely, JCSC2973 (ST30 *SCCmec* type I), M11 and JCSC4625 (ST30 *SCCmec* type IV.3), JCSC4465 (ST30 *SCCmec* type IV.n), and JCSC2958 (ST5 *SCCmec* type II), generated intact PVL phages. This result indicates that  $\phi$ Sa2958-type and  $\phi$ Sa2958-like phages of unknown type can be frequently induced from many *S. aureus* strains with diverse genetic backgrounds. These PVL phages might be induced with or without any stress from *S. aureus* and infect other appropriate recipient *S. aureus* strains, resulting in



the phage conversion of nonvirulent bacteria to virulent bacteria. The facts that two recently isolated strains carried  $\phi$ Sa2958 and that one of them generated infective PVL phages suggest that the  $\phi$ Sa2958 type might be the cause of the recent emergence of PVL-positive MRSA strains in Japan. As shown in Table 5,  $\phi$ Sa2mw carried by MW2 was infectious for two indicator strains, RN4220 and 1039. The USA300 strain has recently been the predominant CA-MRSA clone in the United States.  $\phi$ Sa2usa carries PVL and belongs to the same family as  $\phi$ Sa2mw and  $\phi$ Sa2958, which have elongated heads. Although we could not identify any plaques of  $\phi$ Sa2usa, we have tested FPR3757, whose genome has been sequenced; these  $\phi$ Sa2mw-type or  $\phi$ Sa2usa-type PVL phages might be the cause of the recent appearance of PVL-positive MRSA strains in the United States. Our data also indicate that ST30 MRSA strains carry two different PVL phages. Further study will clarify whether ST30 PVL-positive MRSA clones can be grouped with Western Samoan clones.

In conclusion, we have developed a method of combining several PCRs to identify PVL phage carried by an organism. Our results indicate that two lineages of PVL phages have existed in Japan for many years and that intact phages with elongated heads, such as  $\phi$ Sa2958 and  $\phi$ Sa2958-like phage of unknown type or other phages known simply as elongated-head types, might disseminate among MRSA strains in Japan and confer greater virulence.

#### ACKNOWLEDGMENTS

We thank Yukio Utsui (Daiichi-Sankyo Pharmaceutical Co. Ltd.), Toyoji Okubo (Gunma University), and Takashi Inamatsu (Tokyo Geriatric Hospital) for providing MRSA strains isolated from 1979 through 1985. We thank Toyoko Oguri (Juntendo University Hospital) for providing MRSA strain JCSC4625 and Takashi Kuramoto (Mojiri Rosai Hospital) for providing MRSA strain JCSC6055. We thank the staff of SLR Laboratory for conducting PCR experiments to identify PVL-positive *S. aureus* strains. We thank Keizo Yamaguchi (Toho University) for providing MRSA strains collected by the surveillance study for the susceptibility testing for quinolone antibiotics.

This work was supported by a grant-in-aid for 21st-century COE research, a grant-in-aid for scientific research on priority areas, and a grant-in-aid for scientific research (C19590456) from the Ministry of Education, Science, Sports, Culture and Technology of Japan.

#### REFERENCES

- Baba, T., F. Takeuchi, M. Kuroda, H. Yuzawa, K. Aoki, A. Oguchi, Y. Nagai, N. Iwama, K. Asano, T. Naimi, H. Kuroda, L. Cui, K. Yamamoto, and K. Hiramatsu. 2002. Genome and virulence determinants of high virulence community-acquired MRSA. *Lancet* **359**:1819–1827.
- Baggett, H. C., T. W. Hennessy, K. Rudolph, D. Bruden, A. Reasonover, A. Parkinson, R. Sparks, R. M. Donlan, P. Martinez, K. Mongkolrattanothai, and J. C. Butler. 2004. Community-onset methicillin-resistant *Staphylococcus aureus* associated with antibiotic use and the cytotoxin Pantone-Valentine leukocidin during a furunculosis outbreak in rural Alaska. *J. Infect. Dis.* **189**:1565–1573.
- Begier, E. M., K. Frenette, N. L. Barrett, P. Mshar, S. Petit, D. J. Boxrud, K. Watkins-Colwell, S. Wheeler, E. A. Cebelinski, A. Glennen, D. Nguyen, and J. L. Hadler. 2004. A high-morbidity outbreak of methicillin-resistant *Staphylococcus aureus* among players on a college football team, facilitated by cosmetic body shaving and turf burns. *Clin. Infect. Dis.* **39**:1446–1453.
- Canchaya, C., C. Proux, G. Fournous, A. Bruttin, and H. Brussow. 2003. Prophage genomics. *Microbiol. Mol. Biol. Rev.* **67**:238–276.
- Chambers, H. F. 2001. The changing epidemiology of *Staphylococcus aureus*. *Emerg. Infect. Dis.* **7**:178–182.
- Colin, D. A., I. Mazurier, S. Sire, and V. Finck-Barbancon. 1994. Interaction of the two components of leukocidin from *Staphylococcus aureus* with human polymorphonuclear leukocyte membranes: sequential binding and subsequent activation. *Infect. Immun.* **62**:3184–3188.
- Coombs, G. W., J. C. Pearson, F. G. O'Brien, R. J. Murray, W. B. Grubb, and K. J. Christiansen. 2006. Methicillin-resistant *Staphylococcus aureus* clones, Western Australia. *Emerg. Infect. Dis.* **12**:241–247.
- Denis, O., A. Deplano, H. De Beenhouwer, M. Hallin, G. Huysmans, M. G. Garrino, Y. Glupczynski, X. Malaviole, A. Vergison, and M. J. Struelens. 2005. Polyclonal emergence and importation of community-acquired methicillin-resistant *Staphylococcus aureus* strains harbouring Pantone-Valentine leukocidin genes in Belgium. *J. Antimicrob. Chemother.* **56**:1103–1106.
- Diep, B. A., S. R. Gill, R. F. Chang, T. H. Phan, J. H. Chen, M. G. Davidson, F. Lin, J. Lin, H. A. Carleton, E. F. Mongodin, G. F. Sensabaugh, and F. Perdreau-Remington. 2006. Complete genome sequence of USA300, an epidemic clone of community-acquired methicillin-resistant *Staphylococcus aureus*. *Lancet* **367**:731–739.
- Diep, B. A., G. F. Sensabaugh, N. S. Somboona, H. A. Carleton, and F. Perdreau-Remington. 2004. Widespread skin and soft-tissue infections due to two methicillin-resistant *Staphylococcus aureus* strains harboring the genes for Pantone-Valentine leukocidin. *J. Clin. Microbiol.* **42**:2080–2084.
- Eady, E. A., and J. H. Cove. 2003. Staphylococcal resistance revisited: community-acquired methicillin resistant *Staphylococcus aureus*—an emerging problem for the management of skin and soft tissue infections. *Curr. Opin. Infect. Dis.* **16**:103–124.
- Enright, M. C., N. P. Day, C. E. Davies, S. J. Peacock, and B. G. Spratt. 2000. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J. Clin. Microbiol.* **38**:1008–1015.
- Finck-Barbancon, V., G. Duportal, O. Meunier, and D. A. Colin. 1993. Pore formation by a two-component leukocidin from *Staphylococcus aureus* within the membrane of human polymorphonuclear leukocytes. *Biochim. Biophys. Acta* **1182**:275–282.
- Heinrich, J., M. Velleman, and H. Schuster. 1995. The tripartite immunity system of phages P1 and P7. *FEMS Microbiol. Rev.* **17**:121–126.
- Holmes, A., M. Ganner, S. McGuane, T. L. Pitt, B. D. Cookson, and A. M. Kearns. 2005. *Staphylococcus aureus* isolates carrying Pantone-Valentine leukocidin genes in England and Wales: frequency, characterization, and association with clinical disease. *J. Clin. Microbiol.* **43**:2384–2390.
- Hsu, L. Y., T. H. Koh, T. Y. Tan, T. Ito, X. X. Ma, R. T. Lin, and B. H. Tan. 2006. Emergence of community-associated methicillin-resistant *Staphylococcus aureus* in Singapore: a further six cases. *Singapore Med. J.* **47**:20–26.
- Ito, T., K. Okuma, X. X. Ma, H. Yuzawa, and K. Hiramatsu. 2003. Insights on antibiotic resistance of *Staphylococcus aureus* from its whole genome: genomic island SCC. *Drug Resist. Update* **6**:41–52.
- Ito, T., Y. Katayama, K. Asada, N. Mori, K. Tsutsumimoto, C. Tiensasitorn, and K. Hiramatsu. 2001. Structural comparison of three types of staphylococcal cassette chromosome *mec* integrated in the chromosome in methicillin-resistant *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **45**:1323–1336.
- Kaneko, J., and Y. Kamio. 2004. Bacterial two-component and hetero-heptameric pore-forming cytolysin: structures, pore-forming mechanism, and organization of the genes. *Biosci. Biotechnol. Biochem.* **68**:981–1003.
- Kaneko, J., T. Kimura, Y. Kawakami, T. Tomita, and Y. Kamio. 1997. Pantone-Valentine leukocidin genes in a phage-like particle isolated from mitomycin C-treated *Staphylococcus aureus* V8 (ATCC 49775). *Biosci. Biotechnol. Biochem.* **61**:1960–1962.
- Kaneko, J., T. Kimura, S. Narita, T. Tomita, and Y. Kamio. 1998. Complete nucleotide sequence and molecular characterization of the temperate staphylococcal bacteriophage  $\phi$ PVL carrying Pantone-Valentine leukocidin genes. *Gene* **215**:57–67.
- Kondo, Y., T. Ito, X. X. Ma, S. Watanabe, B. N. Kreiswirth, J. Etienne, and K. Hiramatsu. 2007. Combination of multiplex PCRs for staphylococcal cassette chromosome *mec* type assignment: rapid identification system for *mec*, *cr*, and major differences in junkyard regions. *Antimicrob. Agents Chemother.* **51**:264–274.
- Kwan, T., J. Liu, M. DuBow, P. Gros, and J. Pelletier. 2005. The complete genomes and proteomes of 27 *Staphylococcus aureus* bacteriophages. *Proc. Natl. Acad. Sci. USA* **102**:5174–5179.
- Lindqvist, B. H., G. Deho, and R. Calendar. 1993. Mechanisms of genome propagation and helper exploitation by satellite phage P4. *Microbiol. Rev.* **57**:683–702.
- Ljungquist, E., K. Kockum, and L. E. Bertani. 1984. DNA sequences of the repressor gene and operator region of bacteriophage P2. *Proc. Natl. Acad. Sci. USA* **81**:3988–3992.
- Lucchini, S., F. Desiere, and H. Brussow. 1999. Comparative genomics of *Streptococcus thermophilus* phage species supports a modular evolution theory. *J. Virol.* **73**:8647–8656.
- Ma, X. X., A. Galiana, W. Pedreira, M. Mowszowicz, I. Christophersen, S. Machiavello, L. Lope, S. Benaderet, F. Buela, W. Vincentino, M. Albini, O. Bertaux, I. Constenla, H. Bagnulo, L. Llosa, T. Ito, and K. Hiramatsu. 2005. Community-acquired methicillin-resistant *Staphylococcus aureus*, Uruguay. *Emerg. Infect. Dis.* **11**:973–976.
- Ma, X. X., T. Ito, P. Chongtrakool, and K. Hiramatsu. 2006. Predominance of clones carrying Pantone-Valentine leukocidin genes among methicillin-resistant *Staphylococcus aureus* strains isolated in Japanese hospitals from 1979 to 1985. *J. Clin. Microbiol.* **44**:4515–4527.
- Ma, X. X., T. Ito, C. Tiensasitorn, M. Jamklang, P. Chongtrakool, S. Boyle-Vavra, R. S. Daum, and K. Hiramatsu. 2002. Novel type of staphylococcal cassette chromosome *mec* identified in community-acquired methicillin-resis-



- tant *Staphylococcus aureus* strains. *Antimicrob. Agents Chemother.* **46**:1147–1152.
30. Mulvey, M. R., L. MacDougall, B. Cholin, G. Horsman, M. Fidyk, and S. Woods. 2005. Community-associated methicillin-resistant *Staphylococcus aureus*, Canada. *Emerg. Infect. Dis.* **11**:844–850.
  31. Naimi, T. S., K. H. LeDell, K. Como-Sabetti, S. M. Borchardt, D. J. Boxrud, J. Etienne, S. K. Johnson, F. Vandenesch, S. Fridkin, C. O'Boyle, R. N. Danila, and R. Lynfield. 2003. Comparison of community- and health care-associated methicillin-resistant *Staphylococcus aureus* infection. *JAMA* **290**:2976–2984.
  32. Narita, S., J. Kaneko, J. Chiba, Y. Piemont, S. Jarraud, J. Etienne, and Y. Kamio. 2001. Phage conversion of Pantone-Valentine leukocidin in *Staphylococcus aureus*: molecular analysis of a PVL-converting phage, phiSLT. *Gene* **268**:195–206.
  33. Neve, H., K. I. Zenz, F. Desiere, A. Koch, K. J. Heller, and H. Brussow. 1998. Comparison of the lysogeny modules from the temperate *Streptococcus thermophilus* bacteriophages TP-J34 and Sfi21: implications for the modular theory of phage evolution. *Virology* **241**:61–72.
  34. Okuma, K., K. Iwakawa, J. D. Turnidge, W. B. Grubb, J. M. Bell, F. G. O'Brien, G. W. Coombs, J. W. Pearman, F. C. Tenover, M. Kapi, C. Tien-sasitorn, T. Ito, and K. Hiramatsu. 2002. Dissemination of new methicillin-resistant *Staphylococcus aureus* clones in the community. *J. Clin. Microbiol.* **40**:4289–4294.
  35. Pantone, P. N., M. C. Camb, F. C. O. Valentine, and M. R. C. P. Lond. 1932. Staphylococcal toxin. *Lancet* **i**:506–508.
  36. Prevost, G., B. Cribier, P. Couppie, P. Petiau, G. Supersac, V. Finckel, H. Barbancon, H. Monteil, and Y. Piemont. 1995. Pantone-Valentine leukocidin and gamma-hemolysin from *Staphylococcus aureus* ATCC 49775 are encoded by distinct genetic loci and have different biological activities. *Infect. Immun.* **63**:4121–4129.
  37. Tenover, F. C., L. K. McDougal, R. V. Goering, G. Killgore, S. J. Projan, J. B. Patel, and P. M. Dunman. 2006. Characterization of a strain of community-associated methicillin-resistant *Staphylococcus aureus* widely disseminated in the United States. *J. Clin. Microbiol.* **44**:108–118.
  38. Ushioda, H., T. Terayama, S. Sakai, H. Zen-Yoji, M. Nishiwaki, and A. Hidano. 1981. Coagulase typing of *Staphylococcus aureus* and its application in routine work, p. 77–83. *In* J. Jaljaszewicz (ed.), *Staphylococci and staphylococcus infections*. Zentbl. Bakteriologie, suppl. 10. Gustav Fischer Verlag, Stuttgart, Germany.
  39. Vandenesch, F., T. Naimi, M. C. Enright, G. Lina, G. R. Nimmo, H. Heffernan, N. Liassine, M. Bes, T. Greenland, M. E. Reverdy, and J. Etienne. 2003. Community-acquired methicillin-resistant *Staphylococcus aureus* carrying Pantone-Valentine leukocidin genes: worldwide emergence. *Emerg. Infect. Dis.* **9**:978–984.
  40. Yamaguchi, K., and A. Ohno. 2005. Investigation of the susceptibility trends in Japan to fluoroquinolones and other antimicrobial agents in a nationwide collection of clinical isolates: a longitudinal analysis from 1994 to 2002. *Diagn. Microbiol. Infect. Dis.* **52**:135–143.
  41. Yoshizawa, Y. 1985. Isolation and characterization of restriction negative mutants of *Staphylococcus aureus*. *Jikeikai Med. J.* **32**:415–421.