

## Genetic Heterogeneity among Strains of *Treponema phagedenis*-Like Spirochetes Isolated from Dairy Cattle with Papillomatous Digital Dermatitis in Japan<sup>∇</sup>

Takahisa Yano,<sup>1</sup> Ryoko Yamagami,<sup>1</sup> Kazuhiro Misumi,<sup>2</sup> Chikara Kubota,<sup>3</sup> Kyaw Kyaw Moe,<sup>1</sup> Tetsuya Hayashi,<sup>4,5,8</sup> Kazunori Yoshitani,<sup>6</sup> Osamu Ohtake,<sup>7</sup> and Naoaki Misawa<sup>1,8\*</sup>

Laboratory of Veterinary Public Health, Department of Veterinary Science, Faculty of Agriculture, University of Miyazaki, 1-1 Gakuenkibanadai-nishi, Miyazaki 889-2192, Japan<sup>1</sup>; Laboratory of Veterinary Surgery, Department of Veterinary Science, Faculty of Agriculture, Kagoshima University, 1-21-24 Kourimoto, Kagoshima 890-0065, Japan<sup>2</sup>; Laboratory of Veterinary Reproduction, Department of Veterinary Science, Faculty of Agriculture, Kagoshima University, 1-21-24 Kourimoto, Kagoshima 890-0065, Japan<sup>3</sup>; Division of Microbiology, Department of Infectious Diseases, Faculty of Medicine, University of Miyazaki, 5200 Kiyotake, Miyazaki 889-1692, Japan<sup>4</sup>; Division of Bioenvironmental Science, Frontier Science Research Center, University of Miyazaki, 5200 Kiyotake, Miyazaki 889-1692, Japan<sup>5</sup>; Hokubu Veterinary Clinical Center, NOSAI Chiba, 99-1 Katori, Chiba 289-0407, Japan<sup>6</sup>; Tsuyama Veterinary Clinical Center, NOSAI Okayama, 8-9 Kokubunji, Tsuyama, Okayama 708-0843, Japan<sup>7</sup>; and Project for Zoonoses Education and Research, University of Miyazaki, 1-1 Gakuenkibanadai-nishi, Miyazaki 889-2192, Japan<sup>8</sup>

Received 14 August 2008/Returned for modification 11 December 2008/Accepted 3 January 2009

**Papillomatous digital dermatitis (PDD) is an infectious foot disease of cattle that is prevalent throughout the world. Although it has been prevalent in Japan since the first case was reported in 1992, full epidemiological and bacteriological examinations have not been conducted. We collected 91 lesions of PDD from 80 dairy cattle on 12 farms in eight regions of Japan to isolate the spirochetes that are frequently detected in lesions. We isolated 40 strains of spirochetes from 24 cattle (30.0%) by a simple two-step culture technique, in which the biopsy samples were incubated at 4°C for 48 to 72 h in an enrichment broth supplemented with antibiotics, which improved the rate of isolation, and then inoculated on selective agar plates. All spirochetes examined were catalase positive and oxidase negative and showed weak beta-hemolytic activity. Enzyme activities were identical to those of *Treponema phagedenis* ATCC 27087. Sequencing of the 16S rRNA gene showed that all strains isolated had >99% identity to those of the *T. phagedenis* type strain and of *T. phagedenis*-like strains isolated from PDD lesions in the United States and Europe. Pulsed-field gel electrophoresis and PCR-based random amplified polymorphism DNA methods revealed considerable diversity among strains isolated not only from different cattle but also from the same individuals. These findings may provide further evidence for the role of these treponemes in the pathogenesis of persistent PDD.**

Papillomatous digital dermatitis (PDD) is an infectious foot disease in cattle that was first reported in Italy in 1974 (4) and is now recognized throughout the world (3, 8, 31). In Japan, PDD in dairy cattle is now found in most regions (24, 26, 37) since the first case was reported in 1992 (20). PDD begins as a superficial dermatitis with an erosive lesion and later forms a hyperkeratotic papillomatous lesion with long hairlike projections (12). These lesions are usually located on the rear of the foot between the bulbs of the heel (3). PDD often leads to lameness and decreases in body weight and milk production, causing economic loss and animal welfare problems (30).

It is suspected that PDD is caused by bacteria, since treatment with antibiotics such as oxytetracycline and penicillin G results in rapid resolution of the lesions (19, 29, 46). Microscopic examinations of biopsy samples or direct stamp preparations of the lesions show various bacterial morphotypes, including long filaments, rods, coccoids, and helices. Although

several culturable bacteria have been isolated from PDD lesions (3, 7, 22, 30, 36), it remains unclear whether all are involved in the etiology.

Among the bacteria detected in PDD lesions, a large number of spirochetes are observed (6, 10, 25, 32, 39). Choi et al. detected five spirochetal phylotypes using 16S rRNA gene sequence analysis (6). Similarly, Moter et al. detected four spirochetal phylotypes by using in situ hybridization with oligonucleotide probes of the 16S rRNA gene from treponemes (23). Indeed, several *Treponema* species have been isolated from dairy cattle with PDD, and are closely related to oral *Treponema* species in humans, including *Treponema phagedenis*, *T. denticola*, *T. vincentii*, and *T. medium* (6, 16, 21). Furthermore, Schrank et al. isolated and characterized the newly proposed species *Treponema brennaborensis* (35). These reports suggest that multiple *Treponema* species are present in PDD lesions. Their presence in both superficial lesions and deeper layers of the dermis (2, 8, 12, 25) suggests that they play an important role in the infection. However, it has not been proved that these organisms satisfy Koch's postulates.

One possible reason why the etiology of PDD is not yet fully understood is the difficulty of isolating and culturing *Treponema* species present in PDD lesions. Several research groups

\* Corresponding author. Mailing address: Laboratory of Veterinary Public Health, Department of Veterinary Science, Faculty of Agriculture, University of Miyazaki, 1-1 Gakuenkibanadai-nishi, Miyazaki 889-2192, Japan. Phone and fax: 81-985-58-7284. E-mail: a0d901u@cc.miyazaki-u.ac.jp.

<sup>∇</sup> Published ahead of print on 14 January 2009.

TABLE 1. Isolation of treponemes from cattle with PDD

Region	Farm	No. of cattle examined	No. of lesions sampled <sup>a</sup>	No. of cattle with isolated treponemes	Isolation rate in cattle (%)	No. of treponeme strains isolated <sup>b</sup>
Hokkaido	A	6	8	5	83.3	17
Hokkaido	B	8	12	6	75.0	10
Yamagata	C	2	3	2	100	2
Chiba	D	4	4	1	25.0	1
Chiba	E	2	2	1	50.0	1
Hyougo	F	22	22	2	9.1	2
Hiroshima	G	6	6	1	16.7	1
Kumamoto	H	2	2	1	50.0	1
Miyazaki	I	11	13	0	0	0
Kagoshima	J	5	6	0	0	0
Kagoshima	K	9	9	5	55.6	5
Kagoshima	L	3	4	0	0	0
Total	12	80	91	24	30.0 (mean)	40

<sup>a</sup> PDD lesions from 12 cattle in farms A, B, C, I, J, and L were taken from both rear feet.

<sup>b</sup> The number of strains includes different colonies isolated from the same lesion and different foot lesions from identical cattle in farms A and B.

have isolated *Treponema* species by the use of methods for the isolation of oral treponemes in humans (9, 16, 21, 35, 43). An immunomagnetic bead method using polyclonal antibodies against *T. denticola* and *T. vincentii* was used to separate *Treponema* from other bacteria (9). The growth rate of many *Treponema* species is poor, and the presence of other fast-growing bacteria in PDD lesions often prevents their isolation (43). No selective media supplemented with suitable antimicrobial agents for the primary isolation of treponemes from PDD lesions have been developed. Here, we sought to isolate treponemes from cattle with PDD lesions for the first time in Japan. We developed a simple two-step culture method which improved the rate of isolation of spirochetes and examined the biochemical characteristics and 16S rRNA gene sequences for phylogenetic analysis. Finally, we examined the genotypic heterogeneity among the strains by molecular typing methods.

#### MATERIALS AND METHODS

**Bacterial strains.** *T. phagedenis* ATCC 27087 and *T. denticola* JCM8225 were used as type strains for comparing characteristics of treponemes from PDD lesions. These strains were cultured on the anaerobic medium described below. Isolated strains were suspended in brucella broth (BBL Becton and Dickinson, Sparks, MD) supplemented with 10% glycerol and were kept at  $-80^{\circ}\text{C}$  until testing.

**Sampling of PDD lesions.** We sampled 91 PDD lesions from 80 dairy cattle (Holstein) on 12 farms (A to L) in eight prefectures of Japan from October 2004 to January 2007 (Table 1). Lesions from 12 cattle on farms A, B, C, I, J, and L were taken from both rear feet. Cattle were restrained in a frame, and the affected foot was restrained with a rope. Before sampling, the lesions were washed with water using a soft brush to remove fecal material. Biopsy samples were obtained by using a sterile dermal biopsy punch with a 6-mm diameter (Kai Industries Co., Ltd., Gifu, Japan) and placed in the enrichment medium for anaerobes described below. The biopsy samples were kept on ice or at  $4^{\circ}\text{C}$  and transported to the laboratory within 48 h after sampling.

**Media for transport and isolation.** To isolate treponemes from PDD lesions, we had to kill other bacteria present in the specimens with selective agents. We used rifampin and enrofloxacin, which have been used by others for the isolation of *Treponema* species from PDD lesions (9, 43). We modified an anaerobic medium, named PDDTp agar, to isolate the treponemes. PDDTp consists of GAM agar (Nissui Pharmaceutical Co., Tokyo, Japan) as a base medium for anaerobes supplemented with 0.8% brucella broth (BBL), 0.8% heart infusion broth (Nissui Pharmaceutical), 10% heat-inactivated fetal bovine serum (Invitrogen, San Diego, CA), 10% defibrinated horse blood, rifampin (1  $\mu\text{g}/\text{ml}$ ), and enrofloxacin (1  $\mu\text{g}/\text{ml}$ ). As a transport and enrichment medium, we used PDDTp

broth: GAM broth (Nissui Pharmaceutical) for anaerobes supplemented with the same concentrations of brucella broth (BBL), heart infusion broth (Nissui), fetal bovine serum (Invitrogen), and antibiotics without horse blood.

**Primary microbial isolation.** Biopsy specimens were kept in PDDTp broth at  $4^{\circ}\text{C}$  for 48 to 72 h in air after receipt at our laboratory. The tissue was cut with a sterile scalpel and was stamped onto PDDTp agar plates. Then, several stamped spots were streaked with a loop. The plates were incubated at  $35^{\circ}\text{C}$  for 14 days without oxygen in an AnaeroPack (Mitsubishi Gas Chemical Co., Tokyo, Japan). Film-like or swarming colonies were stained with crystal violet solution (used for Gram staining) for morphological observations. Spiral-shaped rods were analyzed for their biochemical characteristics and 16S rRNA gene sequences as described below. The isolates were subcultured on fresh PDDTp agar plates, and then the bacteria were suspended in brucella broth supplemented with 10% glycerol. These isolates were kept at  $-80^{\circ}\text{C}$  until testing.

**Electron microscopic observations.** Bacteria grown on PDDTp agar plates at  $35^{\circ}\text{C}$  for 14 days were suspended in distilled water: 10  $\mu\text{l}$  of bacterial suspension was mixed with an equal volume of 1% phosphotungstate for negative staining. One drop was placed on a collodion-carbon-coated grid and examined with a Hitachi H-800MU transmission electron microscope (Hitachi Seisakusho, Tokyo, Japan). Three spirochetes were examined for determination of the size and the numbers of periplasmic flagella per cell.

**Enzymatic activities of the isolates.** Bacteria grown on PDDTp agar plates at  $35^{\circ}\text{C}$  for 14 days were examined for hemolytic activity and for catalase and oxidase production. Other enzymatic activities were tested by using an API-ZYM strip system (Biomérieux Japan, Tokyo, Japan). Bacteria were suspended in saline and washed twice with phosphate-buffered saline (PBS) by centrifugation of  $6,000 \times g$  for 20 min at  $4^{\circ}\text{C}$ . After a washing step, cells were resuspended with PBS and adjusted to a density equivalent to McFarland turbidity standard 5. The suspensions were then inoculated into wells of the API-ZYM strip system, which contains substrates for detecting 19 enzymes. The strips were incubated and interpreted according to the manufacturer's instructions.

**Amplification of 16S rRNA gene.** Forty isolates were grown on PDDTp agar plates at  $35^{\circ}\text{C}$  for 10 to 14 days, and chromosomal DNA was prepared with CTAB (cetyltrimethyl ammonium bromide), as described elsewhere (44). The near complete 16S rRNA gene ( $\sim 1,500$  bp) was PCR amplified with universal primers 8F (5'-AGAGTTTGATCMTGGCTCAG-3') and 15R (5'-AAGGAGG TGATCCARCCGCA-3'), which were based on the *Escherichia coli* 16S rRNA numbering system (14). PCR was performed in a final volume of 20  $\mu\text{l}$  containing 20 pM concentrations of forward and reverse primers, a 200  $\mu\text{M}$  concentration of each deoxynucleoside triphosphate, 0.5 U of *Taq* DNA polymerase (Qiagen, Inc., Tokyo, Japan),  $1 \times$  PCR buffer, and 20 ng of DNA template. Thermal cycle conditions were 30 cycles of denaturation at  $94^{\circ}\text{C}$  for 1 min, annealing at  $55^{\circ}\text{C}$  for 1 min, and extension at  $72^{\circ}\text{C}$ , followed by a final  $72^{\circ}\text{C}$  for 10 min. After amplification, the resulting amplicons were electrophoresed in 1% agarose gel.

**16S rRNA sequence data analysis.** The fragments for sequencing were amplified by PCR as described above, and the products were purified by using a QIAquick PCR purification kit (Qiagen). The nucleotide sequences of 40 isolates were determined directly from the PCR fragment in a PCR-based reaction by using an ABI Prism BigDye terminator cycle sequencing ready reaction kit (Applied Biosystems, Tokyo, Japan) and analyzed with a 3130 DNA sequencer (Applied Biosystems). An  $\sim 800$ -bp fragment was sequenced from each 5' and 3' end of the amplicon, and then they were assembled to determine the nucleotide sequence of the nearly complete 16S rRNA gene ( $\sim 1,500$  bp) using GENETYX-WIN 5.4 software. Database similarity searches were performed through the National Center for Biotechnology Information by use of the BLASTN algorithm. To align the 16S rRNA gene sequences, CLUSTAL W software was used (5). Phylogenetic trees were drawn by using TreeView software (28).

**Molecular typing of isolates by PFGE.** *Treponema* strains were grown on PDDTp agar plates without antibiotics for 2 weeks. Bacteria were suspended in 10 mM PBS (pH 7.2), washed twice, and then used to construct plugs with a CHEF Genomic DNA plug kit (Japan Bio-Rad Laboratories, Tokyo, Japan) according to the manufacturer's instructions. Encapsulated DNA was digested at  $37^{\circ}\text{C}$  with 20 U of XbaI (New England Biolabs, Beverly, MA) for 16 h and then separated in 1.2% agarose gel by pulsed-field gel electrophoresis (PFGE) in 1:5-diluted Tris-borate-EDTA (45 mM Tris, 45 mM borate, 1 mM EDTA) buffer (pH 8.3).

The running conditions were as follows: initial switch time, 5 s; final switch time, 50 s; field strength, 6 V/cm with a  $120^{\circ}$  field angle; and 22 h at  $14^{\circ}\text{C}$ . Restriction fragments were stained with ethidium bromide and visualized under UV light.

**RAPD DNA analysis.** For RAPD [random (ly) amplified polymorphic DNA] analysis of *Treponema* strains, reaction mixtures were prepared in a total volume of 25  $\mu\text{l}$ , containing 20 ng of DNA, 2.5  $\mu\text{l}$  of  $10 \times$  PCR buffer, 3 mM  $\text{MgCl}_2$ , 0.625




FIG. 1. Transmission electron microscopy of *T. phagedenis*-like spirochete strain HT201 isolated from a PDD lesion. The spirochete has eight axial flagella at each end.

U of *Taq* DNA polymerase (Qiagen), 250  $\mu$ M concentrations of each deoxynucleoside triphosphate (Invitrogen), and a 160 pM concentration of random primer 1254 (5'-CCGCAGCCAA-3') (1). The PCR program comprised 4 cycles of 94°C for 5 min, 36°C for 5 min, and 72°C for 5 min; followed by 30 cycles of 94°C for 1 min, 36°C for 1 min, and 72°C for 2 min; with a final extension at 72°C for 10 min. The resulting amplicons were separated by electrophoresis in 1% agarose gel and visualized by ethidium bromide. This analysis was repeated at least twice to confirm its reproducibility.

**Dendrogram analysis of PFGE results and 16S rRNA gene sequences.** To examine the relationships between strains, we calculated banding patterns in PFGE by UPGMA (unweighted pair-group method with arithmetic averages) analysis (38) using the Statistica clustering program (Design Technologies, Tokyo, Japan). 16S rRNA gene sequences among strains isolated in the present study and spirochetes submitted to GenBank were compared, and phylogenetic trees were constructed by using the neighbor-joining method of Saitou and Nei (33).

**Nucleotide sequence accession number.** The partial 16S rRNA gene sequence of strain YG3903R was submitted to GenBank and assigned accession number FJ004921.

## RESULTS

**Isolation of helical bacteria from PDD biopsy samples.** The PDDTp culture plates were examined every 7 days after inoculation. Treponemes formed diffuse film-like colonies at the stamped spots after 14 days at 35°C. Although bacterial colonies other than treponemes also grew on the selective agar plates, treponeme-like colonies were easily differentiated from them. Suspected colonies were stained with crystal violet for morphological observations. Helix-shaped bacteria were taken from the edge of their colonies and subcultured on fresh PDDTp agar plates to obtain pure cultures. A total of 40 spirochetes were isolated from 24 of 80 cattle with PDD (30.0%) and from 25 of 91 PDD lesions (27.5%) from nine farms in seven prefectures (Table 1). Multiple strains were isolated from a single lesion or from both feet of six cattle on farms A and B. The isolation rate ranged from 0 to 100% of cattle (Table 1).

**Electron microscopic observation.** Electron microscopy of strain HT201 showed helical shapes 6 to 15  $\mu$ m long and 0.2 to 0.3  $\mu$ m in diameter (Fig. 1). These bacteria had eight axial flagella attached to each end.

**Biochemical characteristics.** We examined 12 better-growing strains among the 40 isolates since enough amount of bacterial cells were necessary for the tests. The strains which did not recover one loopful of bacteria from one PDDTp agar plate after 14 days of incubation were not subjected to the tests. They showed weak beta-hemolytic activity on PDDTp agar plates supplemented with horse blood. *T. phagedenis* ATCC 27087 and all PDD isolates were catalase positive and oxidase negative. The enzyme activities are shown in Table 2. PDD isolates showed results similar to those of *T. phagedenis* ATCC 27087 but not to those of *T. denticola* (Table 2).

**16S rRNA gene sequence analysis.** A fragment of the 16S rRNA gene (~1,500 bp) of each 40 strains was amplified by using universal primers. BLAST searches of the sequence data showed 99 to 100% identity of all strains examined with each other, as well as with PDD strains isolated in the United States, the United Kingdom, and Denmark (accession numbers L78126, EF645248, and AM942450, respectively). The phylogenetic tree (Fig. 2) placed our PDD isolates (only strain YG3903R is shown in the figure) in a single cluster closely related to *T. phagedenis* ATCC 27087.

**PFGE.** We compared 26 strains (Table 3) and type strains of *T. denticola* and *T. phagedenis*. They included multiple colonies isolated from the same lesion or from different lesions on the same cattle on farms A and B. The results revealed considerable diversity among strains isolated not only from different cattle but also from the same individuals (Fig. 3).

**RAPD analysis.** The same strains used for PFGE (Table 3) were tested by RAPD analysis. The RAPD profiles of the isolates appeared to be divergent, as seen in the PFGE anal-

TABLE 2. Enzyme activities of *T. phagedenis*-like spirochetes isolated from PDD lesions

Strain	API-ZYM result <sup>b</sup>																			Other analyses <sup>c</sup>		
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	H	C	O
<i>T. denticola</i> JCM8152	-	+	+	-	+	-	-	+	-	+	-	+	+	-	-	-	-	-	-	+	+	-
<i>T. phagedenis</i> ATCC 27087	+	+	+	-	-	-	-	-	-	+	-	-	+	+	-	-	+	-	+	+	+	-
<i>T. phagedenis</i> -like spirochetes from PDD <sup>a</sup>	+	+	+	-	-	-	-	-	-	+	-	-	+	+	-	-	+	-	+	+	+	-

<sup>a</sup> Twelve strains of *T. phagedenis*-like spirochetes isolated from PDD were examined for the biochemical tests.

<sup>b</sup> Enzymes are represented by numbers above each column as follows: 1, alkaline phosphatase; 2, C4 esterase; 3, C8 esterase lipase; 4, C14 lipase; 5, leucine arylamidase; 6, valine arylamidase; 7, cystine arylamidase; 8, trypsin; 9, chymotrypsin; 10, acid phosphatase; 11, naphtholphosphohydrolase; 12,  $\alpha$ -galactosidase; 13,  $\beta$ -galactosidase; 14,  $\beta$ -glucuronidase; 15,  $\alpha$ -glucosidase; 16,  $\beta$ -glucosidase; 17, *N*-acetyl- $\beta$ -glucosaminidase; 18,  $\alpha$ -mannosidase; 19,  $\alpha$ -fucosidase.

<sup>c</sup> H, hemolytic activity; C, catalase; O, oxidase.

ysis. Even the colonies from the same host revealed different banding patterns, confirming that the result corresponded to that of the PFGE analysis (Fig. 4).

**Dendrograms by UPGMA.** A total of 36 banding patterns were detected in PFGE of the 26 strains examined. Cluster analysis indicated that they formed three major clusters, and each strain showed multiple branching (Fig. 5). Although strains HD21R-R7 and HD21-66 and strains HD26-67, HD26-43, HD26-R4, and HD26-R5 were isolated from two individual

cattle on Farm A, they were represented in all three clusters (Fig. 5).

## DISCUSSION

Culture methods appropriate for isolating treponemes effectively from PDD lesions have not previously been developed. A number of researchers have shown the difficulty of culturing and isolating *Treponema* species from PDD lesions, since the lesions are usually contaminated with feces and soil. Since other coinfecting bacteria (7, 22, 27, 34, 45) may have different growth characteristics and sensitivities to selective agents, as the growth rate of many of *Treponema* species is poor, and as the presence of other bacteria in PDD lesions often prevents the isolation of treponemes because of their overgrowth (43),




FIG. 2. Phylogenetic tree based on nearly complete 16S rRNA gene sequences of strain YG3903R isolated from PDD in the present study and from other *Treponema* species. *Brachyspira hyodysenteriae* was added as an outgroup.

TABLE 3. Strains from 19 cattle used for PFGE and RAPD types

Cattle no. <sup>a</sup>	Strain	Farm	Lesion source <sup>b</sup>	Yr of isolation	Region
1	HT201	H	R-L	2004	Kumamoto
2	YG3093R	C	R-R	2005	Yamagata
3	YG5618	C	R-R	2005	Yamagata
4	HG42	F	NA	2005	Hyogo
5	IZ6-2	K	NA	2005	Kagoshima
6	IZ7-2	K	R-L	2005	Kagoshima
7	CH6	D	R-R	2005	Chiba
8	CH9	E	R-L	2006	Chiba
9*	HD21-66	A	R-L	2007	Hokkaido
	HD21R-R7	A	R-R	2007	Hokkaido
10*	HD22-63	A	R-L	2007	Hokkaido
	HD22R-16	A	R-R	2007	Hokkaido
11	HD23-1	A	R-L	2007	Hokkaido
12	HD24-2	A	R-R	2007	Hokkaido
13†	HD26-43	A	R-R	2007	Hokkaido
	HD26-67	A	R-R	2007	Hokkaido
	HD26-R4	A	R-R	2007	Hokkaido
	HD26-R5	A	R-R	2007	Hokkaido
14†	HD27-4	B	R-L	2007	Hokkaido
	HD27-24	B	R-L	2007	Hokkaido
	HD27-R6	B	R-L	2007	Hokkaido
15	HD30R-16	B	R-R	2007	Hokkaido
16	HD31-84	B	R-L	2007	Hokkaido
17	HD32-R10	B	R-L	2007	Hokkaido
18	HD33-86	B	R-R	2007	Hokkaido
19	HD34-27	B	R-L	2007	Hokkaido

<sup>a</sup> \*, Strains isolated from different lesions on the same cow; †, strains isolated from the same lesion on the same cow.

<sup>b</sup> R-R, rear right foot; R-L, rear left foot; NA, not available.




FIG. 3. PFGE analysis of PDD isolates. We examined 26 strains isolated from PDD lesions and two type strains of *Treponema* species. Multiple colonies from individual PDD lesions on the same host are included. Lanes and strains are as follows: 1, *T. denticola* JCM8225; 2, *T. phagedenis* ATCC 27087; 3, HT201; 4, YG3903R; 5, YG5618; 6, HG42; 7, IZ6-2; 8, IZ7-2; 9, CH6; 10, CH9; 11, HD21-66; 12, HD21R-R7; 13, HD22-63; 14, HD22R-16; 15, HD23-1; 16, HD24-2; 17, HD26-43; 18, HD26-67; 19, HD26-R4; 20, HD26-R5; 21, HD27-4; 22, HD27-24; 23, HD27-R6; 24, HD30-R16; 25, HD31-84; 26, HD32-R10; 27, HD33-86; and 28, HD34-27. Lanes M contain the ladder markers for PFGE.

varying the culture media, temperature, and time could improve isolation.

In the present study, we developed a simple two-step culture method that improved the rate of isolation. Initially, we used enrichment culture with selective antibiotics at 35°C, followed by direct plating on anaerobic agar plates. However, the enrichment culture allowed other bacteria to proliferate. Next, we inoculated biopsy specimens on PDDTp agar plates immediately after sampling, but we isolated only one spirochete from about 60 PDD lesions (data not shown). The rate of the isolation of treponemes from the samples in PDDTp broth, in which they were transported over 2 days, was improved over previous methods. After incubation at 4°C for 48 to 72 h in the PDDTp broth, nontreponeme bacteria inoculated on PDDTp agar plates grew slowly, and it was easier to separate the treponemes from them. Although why the other bacteria did not grow well remains unclear, many of them may have been killed by the antibiotics in the broth during incubation at low temperature, conditions which the isolated treponemes may be better able to survive. All treponemes isolated by this method were identified as *T. phagedenis*-like spirochetes.

Walker et al. isolated *T. phagedenis*-like, *T. denticola*-like, and *T. vincentii*/*T. medium*-like spirochetes in the United States (43). Recently, Evans et al. isolated these three groups from PDD lesions in United Kingdom cattle (16). Further-




FIG. 4. RAPD fingerprints of 26 PDD isolates and 2 type strains. Lanes and strains are as follows: 1, *T. denticola* JCM8225; 2, *T. phagedenis* ATCC 27087; 3, HT201; 4, YG3903R; 5, YG5618; 6, HG42; 7, IZ6-2; 8, IZ7-2; 9, CH6; 10, CH9; 11, HD21-66; 12, HD21R-R7; 13, HD22-63; 14, HD22R-16; 15, HD23-1; 16, HD24-2; 17, HD26-43; 18, HD26-67; 19, HD26-R4; 20, HD26-R5; 21, HD27-4; 22, HD27-24; 23, HD27-R6; 24, HD30-R16; 25, HD31-84; 26, HD32-R10; 27, HD33-86; and 28, HD34-27. Lanes M contain 1.0-kb (left) and 0.1-kb (right) ladder markers.

more, Schrank et al. isolated and characterized the newly proposed species *T. brennaborensis* (35). Since it is still unclear which *Treponema* is the most predominant species in PDD lesions in Japan, we cannot conclude that our culture method is suitable for the isolation of treponemes other than *T. phagedenis*-like spirochetes. However, *T. denticola* and *T. phagedenis* isolated from humans were able to grow on the PDDTp medium used in the present study (data not shown). Thus, the existence of treponemes other than *T. phagedenis*-like spirochetes in PDD lesions of cattle in Japan should be determined by molecular biological methods, such as PCR and in situ hybridization.

All 16S rRNA gene sequences of the isolates showed 99 to 100% identity with each other and with some isolates from the




FIG. 5. Dendrogram of cluster analysis based on PFGE banding patterns. We examined 26 strains isolated from PDD lesions and 2 type strains of *Treponema* species.

United States (43) and the United Kingdom (9, 16). Phylogenetic analysis showed a close relationship between our PDD isolates and *T. phagedenis* ATCC 27087. Our isolates were catalase positive and oxidase negative and showed weak beta-hemolytic activity. Enzyme activities of all isolates examined were identical, and the enzyme profile was more closely related to that of *T. phagedenis* ATCC 27087 but not that of *T. denticola* JCM8225. However, the activities were not identical to those of strains reported previously (11, 16, 43), even though the 16S rRNA gene sequences were more than 99% identical. The 16S rRNA sequence might be similar, but the sequences that encode various enzymes might be more variable. These phenotypic characteristics may change with passage number or culture conditions.

In previous studies, the genetic diversity of *T. phagedenis*-like organisms isolated from PDD was assessed by molecular techniques such as PFGE and PCR-based restriction fragment length polymorphism (41), because epidemiological methods such as serotyping, biotyping, and outer membrane profiling have not been established for *T. phagedenis*-like spirochetes. Here, we used PFGE and RAPD to detect the heterogeneity of DNAs of *T. phagedenis*-like strains. The two typing methods revealed considerable diversity among the strains, not only from the same farm but also from the same animals. Trott et al. showed by PFGE analysis that 6 strains of *T. phagedenis*-like spirochetes were genetically heterogeneous (41). Walker et al. similarly showed that eight treponemes from various farms had different restriction patterns (43). Our result is consistent with these reports. Interestingly, we found genetic diversity among strains isolated from the same PDD lesion.

There are two possible main reasons for the DNA diversity among the strains: multiple infections with different strains and the acquisition of genetic diversity in vivo. Some evidence of genomic recombination or rearrangement in pathogens has been demonstrated (13, 40). Such mechanisms allow organisms to generate antigenic diversity, helping them to evade the immunological response of the host (17, 18). PDD is a persistent infection, and spirochetes are observed in most stages of infection (6). Although systemic antibodies against *T. phagedenis*-like spirochetes were detected (42), they may not be protective. Furthermore, recurrent infection has also been reported (26). These findings may provide further evidence for the role of these organisms in the pathogenesis of PDD. Long-term genotyping and phenotyping in experimental infection studies are needed to assess the genomic diversity among isolates over time in the same lesion.

The etiologic involvement of treponemes in PDD is incompletely known. However, the result that treponemes with very similar 16S rRNA gene sequences are common in PDD lesions in different countries implies that they may be one of the most predominant populations in the lesions and play a role in the development of PDD. Zuerner et al. showed that *T. phagedenis*-like spirochetes from PDD lesions suppress the innate immune response in cattle (47). Elliott et al. reported that *T. phagedenis*-like spirochetes induced abscesses in mice (15). These discoveries imply that *T. phagedenis*-like spirochetes have pathogenic potential. Further studies of the pathogenic features of PDD-related treponemes are required to clarify the etiology of PDD.

## ACKNOWLEDGMENTS

This study was supported by Japan Farriers Association, KAKENHI (Grant-in-Aid for Scientific Research) on Priority Areas "Applied Genomics" from the Ministry of Education, Culture, Sports, Science, and Technology of Japan, and Project for Zoonoses Education and Research, University of Miyazaki.

## REFERENCES

1. Akopyanz, N., N. O. Bukanov, T. U. Westblom, S. Kresovich, and D. E. Berg. 1992. DNA diversity among clinical isolates of *Helicobacter pylori* detected by PCR-based RAPD fingerprinting. *Nucleic Acids Res.* **20**:5137–5142.
2. Blowey, R. W., S. H. Done, and W. Cooley. 1994. Observations on the pathogenesis of digital dermatitis in cattle. *Vet. Rec.* **135**:115–117.
3. Blowey, R. W., and M. W. Sharp. 1988. Digital dermatitis in dairy cattle. *Vet. Rec.* **122**:505–508.
4. Cheli, R., and C. M. Mortellaro. 1974. La dermatite digitale del bovino, p. 208–213. In P. Gallarati (ed.), *Proceedings of the 8th International Conference on Diseases of Cattle*. International Conference on Diseases of Cattle, Piacenza, Italy.
5. Chenna, R., H. Sugawara, T. Koike, R. Lopez, T. J. Gibson, D. G. Higgins, and J. D. Thompson. 2003. Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Res.* **31**:3497–3500.
6. Choi, B. K., H. Nattermann, S. Grund, W. Haider, and U. B. Gobel. 1997. Spirochetes from digital dermatitis lesions in cattle are closely related to treponemes associated with human periodontitis. *Int. J. Syst. Bacteriol.* **47**:175–181.
7. Collighan, R. J., and M. J. Woodward. 1997. Spirochaetes and other bacterial species associated with bovine digital dermatitis. *FEMS Microbiol. Lett.* **156**:37–41.
8. Cruz, C. E., C. A. Pescador, Y. Nakajima, and D. Driemeier. 2005. Immunopathological investigations on bovine digital epidermitis. *Vet. Rec.* **157**:834–840.
9. Demirkan, I., S. D. Carter, C. A. Hart, and M. J. Woodward. 1999. Isolation and cultivation of a spirochaete from bovine digital dermatitis. *Vet. Rec.* **145**:497–498.
10. Demirkan, I., S. D. Carter, R. D. Murray, R. W. Blowey, and M. J. Woodward. 1998. The frequent detection of a treponeme in bovine digital dermatitis by immunocytochemistry and polymerase chain reaction. *Vet. Microbiol.* **60**:285–292.
11. Demirkan, I., H. F. Williams, A. Dhawi, S. D. Carter, C. Winstanley, K. D. Bruce, and C. A. Hart. 2006. Characterization of a spirochaete isolated from a case of bovine digital dermatitis. *J. Appl. Microbiol.* **101**:948–955.
12. Döpfer, D., A. Koopmans, F. A. Meijer, I. Szakall, Y. H. Schukken, W. Klee, R. B. Bosma, J. L. Cornelisse, A. J. van Asten, and A. A. ter Huurne. 1997. Histological and bacteriological evaluation of digital dermatitis in cattle, with special reference to spirochaetes and *Campylobacter faecalis*. *Vet. Rec.* **140**:620–623.
13. Dutta, C., and A. Pan. 2002. Horizontal gene transfer and bacterial diversity. *J. Biosci.* **27**:27–33.
14. Edwards, U., T. Rogall, H. Blocker, M. Emde, and E. C. Bottger. 1989. Isolation and direct complete nucleotide determination of entire genes: characterization of a gene coding for 16S rRNA. *Nucleic Acids Res.* **17**:7843–7853.
15. Elliott, M. K., D. P. Alt, and R. L. Zuerner. 2007. Lesion formation and antibody response induced by papillomatous digital dermatitis-associated spirochetes in a murine abscess model. *Infect. Immun.* **75**:4400–4408.
16. Evans, N. J., J. M. Brown, I. Demirkan, R. D. Murray, W. D. Vink, R. W. Blowey, C. A. Hart, and S. D. Carter. 2008. Three unique groups of spirochetes isolated from digital dermatitis lesions in UK cattle. *Vet. Microbiol.* **131**:141–150.
17. Futse, J. E., K. A. Brayton, M. J. Dark, D. P. Knowles, Jr., and G. H. Palmer. 2008. Superinfection as a driver of genomic diversification in antigenically variant pathogens. *Proc. Natl. Acad. Sci. USA* **105**:2123–2127.
18. Grogono-Thomas, R., M. J. Blaser, M. Ahmadi, and D. G. Newell. 2003. Role of S-layer protein antigenic diversity in the immune responses of sheep experimentally challenged with *Campylobacter fetus* subsp. *fetus*. *Infect. Immun.* **71**:147–154.
19. Hernandez, J., J. K. Shearer, and J. B. Elliott. 1999. Comparison of topical application of oxytetracycline and four nonantibiotic solutions for treatment of papillomatous digital dermatitis in dairy cows. *J. Am. Vet. Med. Assoc.* **214**:688–690.
20. Kimura, Y., M. Takahashi, H. Matsumoto, H. Tsukuda, M. Sato, K. Ohgawara, M. Kanoe, N. Goto, O. Aoki, and M. Hataya. 1992. Verrucose dermatitis and digital papillomatosis in dairy cows. *Hoof* **160**:37–38. (In Japanese.)
21. Klitgaard, K., M. Boye, N. Capion, and T. K. Jensen. 2008. Evidence of multiple *Treponema* phylotypes involved in bovine digital dermatitis as shown by 16S rDNA analysis and fluorescent in situ hybridization. *J. Clin. Microbiol.* **46**:3012–3020.

22. Koniarová, I., A. Orsag, and V. Ledecký. 1993. The role anaerobes in dermatitis digitalis et interdigitalis in cattle. *Vet. Med.* **38**:589–596.
23. Moter, A., G. Leist, R. Rudolph, K. Schrank, B. K. Choi, M. Wagner, and U. B. Gobel. 1998. Fluorescence in situ hybridization shows spatial distribution of as yet uncultured treponemes in biopsies from digital dermatitis lesions. *Microbiology* **144**(Pt. 9):2459–2467.
24. Nagai, F., H. Ota, K. Fujimoto, H. Abe, T. Munakata, T. Abe, M. Tanaka, A. Ito, Y. Yamamoto, T. Ando, M. Koiwa, H. Taniyama, N. Kikuchi, and N. Kusaba. 2000. Mass outbreak of verrucous dermatitis and digital papillomatous caused by *Treponema*-like spirochetes in daily cattle. *J. Jpn. Vet. Med. Assoc.* **53**:577–581. (In Japanese.)
25. Nordhoff, M., A. Moter, K. Schrank, and L. H. Wieler. 2008. High prevalence of treponemes in bovine digital dermatitis—A molecular epidemiology. *Vet. Microbiol.* **131**:293–300.
26. Ohtake, O. 2003. Present status of leg and hoof diseases are frequently reported from various parts in Japan. *Yamaguchi J. Vet. Med.* **30**:41–50. (In Japanese.)
27. Ohya, T., H. Yamaguchi, Y. Nii, and H. Ito. 1999. Isolation of *Campylobacter sputorum* from lesions of papillomatous digital dermatitis in dairy cattle. *Vet. Rec.* **145**:316–318.
28. Page, R. D. 1996. TreeView: an application to display phylogenetic trees on personal computers. *Comput. Appl. Biosci.* **12**:357–358.
29. Read, D. H., and R. L. Walker. 1998. Papillomatous digital dermatitis (foot-warts) in California dairy cattle: clinical and gross pathologic findings. *J. Vet. Diagn. Investig.* **10**:67–76.
30. Read, D. H., R. L. Walker, A. E. Castro, J. P. Sundberg, and M. C. Thurmond. 1992. An invasive spirochaete associated with interdigital papillomatosis of dairy cattle. *Vet. Rec.* **130**:59–60.
31. Rebhun, W. C., R. M. Payne, J. M. King, M. Wolfe, and S. N. Begg. 1980. Interdigital papillomatosis in dairy cattle. *J. Am. Vet. Med. Assoc.* **177**:437–440.
32. Rijpkema, S. G., G. P. David, S. L. Hughes, and M. J. Woodward. 1997. Partial identification of spirochaetes from two dairy cows with digital dermatitis by polymerase chain reaction analysis of the 16S rRNA gene. *Vet. Rec.* **140**:257–259.
33. Saitou, N., and M. Nei. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**:406–425.
34. Schlafer, S., M. Nordhoff, C. Wyss, S. Strub, J. Hubner, D. M. Gescher, A. Petrich, U. B. Gobel, and A. Moter. 2008. Involvement of *Guggenheimella bovis* in digital dermatitis lesions of dairy cows. *Vet. Microbiol.* **128**:118–125.
35. Schrank, K., B. K. Choi, S. Grund, A. Moter, K. Heuner, H. Nattermann, and U. B. Gobel. 1999. *Treponema brennaborensis* sp. nov., a novel spirochaete isolated from a dairy cow suffering from digital dermatitis. *Int. J. Syst. Bacteriol.* **49**(Pt. 1):43–50.
36. Schroeder, C. M., K. W. Parlor, T. L. Marsh, N. K. Ames, A. K. Goeman, and R. D. Walker. 2003. Characterization of the predominant anaerobic bacterium recovered from digital dermatitis lesions in three Michigan dairy cows. *Anaerobe* **9**:151–155.
37. Shibahara, T., T. Ohya, R. Ishii, Y. Ogihara, T. Maeda, Y. Ishikawa, and K. Kadota. 2002. Concurrent spirochaetal infections of the feet and colon of cattle in Japan. *Aust. Vet. J.* **80**:497–502.
38. Sneath, P. H., and R. R. Sokal. 1973. Numerical taxonomy: the principles and practice of numerical classification. W. H. Freeman & Co., San Francisco, CA.
39. Stamm, L. V., H. L. Bergen, and R. L. Walker. 2002. Molecular typing of papillomatous digital dermatitis-associated *Treponema* isolates based on analysis of 16S-23S ribosomal DNA intergenic spacer regions. *J. Clin. Microbiol.* **40**:3463–3469.
40. Sung, S. Y., J. V. McDowell, J. A. Carlyon, and R. T. Marconi. 2000. Mutation and recombination in the upstream homology box-flanked *ospE*-related genes of the Lyme disease spirochetes result in the development of new antigenic variants during infection. *Infect. Immun.* **68**:1319–1327.
41. Trott, D. J., M. R. Moeller, R. L. Zuerner, J. P. Goff, W. R. Waters, D. P. Alt, R. L. Walker, and M. J. Wannemuehler. 2003. Characterization of *Treponema phagedenis*-like spirochetes isolated from papillomatous digital dermatitis lesions in dairy cattle. *J. Clin. Microbiol.* **41**:2522–2529.
42. Walker, R. L., D. H. Read, K. J. Loretz, D. W. Hird, and S. L. Berry. 1997. Humoral response of dairy cattle to spirochetes isolated from papillomatous digital dermatitis lesions. *Am. J. Vet. Res.* **58**:744–748.
43. Walker, R. L., D. H. Read, K. J. Loretz, and R. W. Nordhausen. 1995. Spirochetes isolated from dairy cattle with papillomatous digital dermatitis and interdigital dermatitis. *Vet. Microbiol.* **47**:343–355.
44. Wilson, K. 1987. Preparation of genomic DNA from bacteria, p. 293–332. *In* F. M. Ausubel, R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl (ed.), *Current protocols in molecular biology*, unit 2.4. Wiley, New York, NY.
45. Wyss, C., F. E. Dewhirst, B. J. Paster, T. Thurnheer, and A. Luginbuhl. 2005. *Guggenheimella bovis* gen. nov., sp. nov., isolated from lesions of bovine dermatitis digitalis. *Int. J. Syst. Evol. Microbiol.* **55**:667–671.
46. Yeruham, I., and S. Perl. 1998. Clinical aspects of an outbreak of papillomatous digital dermatitis in a dairy cattle herd. *J. S. Afr. Vet. Assoc.* **69**:112–115.
47. Zuerner, R. L., M. Heidari, M. K. Elliott, D. P. Alt, and J. D. Neill. 2007. Papillomatous digital dermatitis spirochetes suppress the bovine macrophage innate immune response. *Vet. Microbiol.* **125**:256–264.