

Multiplex-PCR Method for Species Identification of Coagulase-Positive Staphylococci[∇]

Takashi Sasaki,^{1*} Sae Tsubakishita,¹ Yoshikazu Tanaka,² Arihito Sakusabe,³ Masayuki Ohtsuka,¹ Shintaro Hirotsuki,¹ Tetsuji Kawakami,⁴ Tsuneo Fukata,⁴ and Keiichi Hiramatsu¹

Department of Infection Control Science, Faculty of Medicine, Juntendo University, Tokyo 113-8421, Japan¹; Department of Veterinary Hygiene, Veterinary School, Nippon Veterinary and Life Science University, 1-7-1 Kyounan, Musashino, Tokyo 180-8602, Japan²; Awachuo Animal Hospital, 60 Yawata, Tateyama, Chiba 294-0047, Japan³; and Department of Veterinary Medicine, Faculty of Applied Biological Sciences, Gifu University, Gifu 501-1193, Japan⁴

Received 24 June 2009/Returned for modification 16 August 2009/Accepted 21 December 2009

In veterinary medicine, coagulase-positive staphylococci (CoPS) other than *Staphylococcus aureus* have frequently been misidentified as being *S. aureus* strains, as they have several phenotypic traits in common. There has been no reliable method to distinguish among CoPS species in veterinary clinical laboratories. In the present study, we sequenced the thermonuclease (*nuc*) genes of staphylococcal species and devised a multiplex-PCR (M-PCR) method for species identification of CoPS by targeting the *nuc* gene locus. To evaluate sensitivity and specificity, we used this M-PCR method on 374 staphylococcal strains that had been previously identified to the species level by an *hsp60* sequencing approach. We could successfully distinguish between *S. aureus*, *S. hyicus*, *S. schleiferi*, *S. intermedius*, *S. pseudintermedius*, and *S. delphini* groups A and B. The present method was both sensitive (99.8%) and specific (100%). Our M-PCR assay will allow the routine species identification of CoPS isolates from various animal species for clinical veterinary diagnosis.

The genus *Staphylococcus* is present in skin and nasal flora and causes opportunistic infections in humans and various animals. To date, seven species of coagulase-positive staphylococci (CoPS) have been identified: *Staphylococcus aureus*, *S. intermedius*, *S. schleiferi* subsp. *coagulans*, *S. hyicus*, *S. lutrae*, *S. delphini*, and *S. pseudintermedius* (10, 12). In addition to *S. aureus*, the other CoPS species can cause severe infections compared with those caused by coagulase-negative staphylococci (CoNS) (2, 13, 16, 22, 25, 28, 32).

It is known that staphylococcal species exhibit host specificity, and the species of CoPS isolated from clinical specimens differ with host animal species; for example, the predominant species in ruminants, pigs, dogs, and pigeons (*Columba livia*) are *S. aureus*, *S. hyicus*, *S. pseudintermedius*, and *S. intermedius*, respectively (11, 28). MIC breakpoints of oxacillin to determine methicillin resistance differ with species. Therefore, the identification of CoPS to the species level is an important task for veterinary diagnostic laboratories.

It is difficult to discriminate between CoPS species based on phenotypic differences because there is a lack of unique biochemical markers for species identification (12, 28). Although various molecular methods have been reported (1, 3, 6, 7, 19, 20, 24, 28), they are costly and/or time-consuming, and interpretation of the results is complicated. Thus, a simple and precise method for discriminating among CoPS species is highly desirable.

In the present study, we performed a sequence analysis of

nuc genes in CoPS and related species and developed a multiplex-PCR (M-PCR) method for the species identification of the CoPS-targeted *nuc* gene locus.

MATERIALS AND METHODS

Bacterial strains and species identification. As shown in Table 1, eight CoPS strains and six closely related CoNS species were used for phylogenetic analysis based on thermonuclease (*nuc*) genes. To evaluate the sensitivity and specificity of M-PCR for the species identification of CoPS, 374 staphylococcal strains derived from various animal species were used in the present study (Table 2).

All strains used in this study were identified to the species level by sequencing analysis based on the *hsp60* gene (19). The discrimination among *S. delphini* groups A and B was performed by using a *nuc* gene sequencing method reported previously (28). The identification of *S. schleiferi* to the subspecies level was performed by a coagulase test using rabbit serum (Eiken Chemical Co., Ltd., Tokyo, Japan).

Strains were stored in 10% skim milk at -80°C until use and were maintained on Trypticase soy agar II with 5% sheep blood (BD Japan, Co., Ltd., Tokyo, Japan).

DNA extraction. A single colony was suspended to a 1.0 McFarland standard in 100 μl of TE buffer (10 mM Tris, 1 mM EDTA [pH 8.0]) with 10 U of achromopeptidase (Wako Chemical Co., Ltd.), and the suspension was incubated at 55°C for 10 min. Supernatants were used as crude DNA extracts for PCR.

Amplification and sequence analysis of the *nuc* gene. In order to amplify the conserved regions of *nuc* genes, degenerate primers were designed by multiple alignments of amino acid sequences of the staphylococcal *nuc* genes, which were available from NCBI databases. Primers Nuc-alf1 (5'-CCNAAYACNCCNGTNCARCCN-3') and Nuc-alfR (5'-NADCCANACTANGCNARNGT-3') were used. The reaction mixture for the PCR consisted of 2 μl of DNA extract in a total volume of 50 μl composed of 2 U of Ex *Taq* (Takara Co., Ltd., Kyoto, Japan), 30 pmol each primer, 0.2 mM deoxynucleoside triphosphate mixture, and 1 \times reaction buffer (Takara). Reaction mixtures were thermally cycled once at 95°C for 2 min; 30 times at 95°C for 30 s, 52°C for 30 s, and 72°C for 30 s; and then once at 72°C for 2 min. The PCR product was cloned into plasmid pCR-4 I-TOPO (Invitrogen, Life Technologies, Carlsbad, CA) and was transformed into *Escherichia coli* TOP10 cells (Invitrogen). Insert DNA of the recombinant plasmid was sequenced by using a Big Dye Terminator (version 3.1) cycle sequencing kit (Applied Biosystems, Foster City, CA) with an ABI Prism 3100

* Corresponding author. Mailing address: Department of Infection Control Science, Faculty of Medicine, Juntendo University, 2-1-1 Hongo, Bunkyo-ku, Tokyo 113-8421, Japan. Phone: 81-3-3813-3111, ext. 3822. Fax: 81-3-5684-7830. E-mail: hiteiha@juntendo.ac.jp.

[∇] Published ahead of print on 6 January 2010.

TABLE 1. Strains used for sequencing analysis of the thermonuclease (*nuc*) gene

Species	Strain	Source	Size of ORF (bp)	GenBank accession no. for sequence of <i>nuc</i>	Reference or source
<i>S. aureus</i>	N315	Human	534	BA000018	18
<i>S. pseudintermedius</i>	LMG 22219T	Cat	507	AB327164	28
<i>S. delphini</i> group A	LMG 22190T	Dolphin	516	AB327167	This study
	CCUG 51769	Mink	507	AB327168	This study
<i>S. delphini</i> group B	P-27B	Pigeon	507	AB327166	28
	<i>S. intermedius</i>	ATCC 29663T	Pigeon	507	AB327165
<i>S. schleiferi</i> subsp. <i>schleiferi</i>	P-45C	Pigeon	504	None	This study
<i>S. schleiferi</i> subsp. <i>coagulans</i>	JCM 7470T	Dog	504	AB465334	This study
<i>S. hyicus</i>	JCM 2423T	Pig	510	AB465332	This study
<i>S. felis</i>	JCM 7469T	Cat	507	AB465335	This study
<i>S. chromogenes</i>	P-29B	Pigeon	507	AB465333	This study
<i>S. haemolyticus</i>	JCSC 1435	Human	537	AP006716	29
<i>S. epidermidis</i>	RP62A	Human	537	CP000029	15
<i>S. saprophyticus</i>	ATCC 15305T	Human	531	AP008934	17

genetic analyzer (Applied Biosystems). The 5' and 3' regions were obtained by inverse PCR, and the complete *nuc* gene sequences were determined. All staphylococcal species sequenced in previous studies harbored the *nuc* gene at a specific gene locus (*nuc* gene locus), which was located about 2 to 8 kbp downstream of the aspartate kinase gene (SA1163) (15, 17, 18, 29). For the species for which degenerate PCR of the *nuc* gene was inadequate, degenerate primers targeting the aspartate kinase gene were designed. Primers Nuc-AsdF (5'-WR NCKRTTCATNARRTAYTT-3') and AsdR (5'-ACNTAYMGNGARATGMG NGAR-3') were used to amplify the conserved regions of the aspartate kinase genes. Reaction mixtures were thermally cycled once at 95°C for 2 min; 30 times at 95°C for 30 s, 50°C for 30 s, and 72°C for 30 s; and then once at 72°C for 2 min. Downstream sequences were analyzed by inverse PCR in order to determine the complete sequence of *nuc*. Multiple alignment was carried out by using the CLUSTAL W program (30). Construction of the phylogenetic tree was performed by the neighbor-joining method (26).

M-PCR for species identification of CoPS. Primers for M-PCR were designed to amplify a portion of the *nuc* gene locus (Table 3). The reaction mixture for PCR consisted of 2 µl of DNA extract in a total volume of 50 µl composed of 2 U of Ex *Taq* (Takara Co., Ltd., Kyoto, Japan), 10 pmol each primer, 0.2 mM deoxynucleoside triphosphate mixture, and 1× reaction buffer (Takara). Reaction mixtures were thermally cycled once at 95°C for 2 min; 30 times at 95°C for 30 s, 56°C for 35 s, and 72°C for 1 min; and then once at 72°C for 2 min. DNA fragments were analyzed by electrophoresis in 1× Tris-acetate-EDTA on a 1% agarose gel stained with ethidium bromide.

In order to evaluate sensitivity and specificity, we applied the present M-PCR method to 314 CoPS and 60 CoNS strains (Table 2).

RESULTS

Sequence analysis of *nuc* genes of staphylococci. As shown in Table 1, all staphylococcal species analyzed in this study had kept the *nuc* gene. With regard to the *nuc* phylogenetic tree (Fig. 1), the relationship among CoPS species other than *S. delphini* group B agreed with that determined by the 16S rRNA and *hsp60* genes (14, 19). The *nuc* sequence for *S. delphini* group B was more closely related to that of *S. pseudintermedius* LMG 22219^T than to that of *S. delphini* LMG 22190^T (*S. delphini* group A), as previously reported (28). Five *S. schleiferi* subsp. *coagulans* strains were phylogenetically indistinguishable from 31 *S. schleiferi* subsp. *schleiferi* strains (data not shown).

The nucleotide identity of the *nuc* genes among the CoPS and closely related CoNS species ranged from 60.0 to 95.9% (mean, 71.7%). The most similar pair was *S. pseudintermedius* and *S. delphini* group B (95.9%).

M-PCR targeting the *nuc* gene locus for species identification of CoPS. By using this M-PCR method, seven species of CoPS (*S. aureus*, *S. intermedius*, *S. schleiferi* subsp. *coagulans*, *S. delphini* group A, *S. hyicus*, *S. pseudintermedius*, and *S. delphini* group B) showed a successful amplification of internal fragments with the expected sizes (359 bp, 430 bp, 526 bp, 661 bp, 793 bp, 926 bp, and 1,135 bp, respectively) with the primer pairs specific for each species (Fig. 2). Among the tested CoPS strains, all strains other than an *S. pseudintermedius* strain were correctly identified to the species level. This *S. pseudintermedius* strain had a 1.6-kbp insertion mutation (78% nucleotide identity with a partial sequence of the IS1181 transposase) within the *nuc* gene open reading frame (ORF). In addition to *S. schleiferi* subsp. *coagulans* strains, all *S. schleiferi* subsp. *schleiferi* strains had 526-bp fragments amplified by the M-PCR method. Consequently, our method had no discriminating power at the subspecies level for *S. schleiferi*. We also applied this method to *S. lutrae* CCUG 38494^T and 29 CoNS species other than *S. schleiferi* subsp. *schleiferi*, and no false-positive result was observed. Consequently, this method is both sensitive (99.8%) and specific (100%).

DISCUSSION

The species identification of CoPS needs to be performed accurately in veterinary clinical laboratories for two reasons.

The first reason is that the MIC breakpoints of oxacillin to determine methicillin resistance in staphylococci differ with species. According to Clinical and Laboratory Standards Institute (CLSI) guidelines, MIC cutoff values for oxacillin for determining methicillin resistance against *S. aureus* and *S. lugdunensis* are 4 µg/ml and differ from the values for other species (0.5 µg/ml) (8). It was previously reported that some strains exhibited oxacillin MICs of 0.5 to <4 µg/ml among *mecA*-positive *S. pseudintermedius* strains (4, 9, 27). If such strains are not identified as *S. pseudintermedius* strains but are identified as *S. aureus* strains, they could be misidentified as being methicillin-susceptible strains. Such inadequate species identification could lead to suboptimal or inappropriate treat-

TABLE 2. Staphylococcal strains used as the “gold standard” population previously identified to the species level by an *hsp60* and/or *nuc* sequencing method

Species	Total no. of isolates	Host origin(s) (no. of isolates)
CoPS		
<i>S. aureus</i>	133	Human (33), chicken (17), pig (10), horse (5), cat (44), dog (5), bovine (19)
<i>S. intermedius</i>	12	Pigeon (12)
<i>S. schleiferi</i> subsp. <i>coagulans</i>	5	Dog (3), cat (1), polar bear (1)
<i>S. delphini</i> group A	17	Pigeon (12), horse (3), mink (1), dolphin (1)
<i>S. delphini</i> group B	8	Pigeon (2), horse (6)
<i>S. hyicus</i>	27	Pig (21), chicken (6)
<i>S. pseudintermedius</i>	111	Dog (97), cat (12), human (2)
<i>S. lutrae</i>	1	Otter (1)
CoNS		
<i>S. schleiferi</i> subsp. <i>schleiferi</i>	31	Pigeon (16), dog (12), cat (3)
<i>S. felis</i>	1	Cat (1)
<i>S. chromogenes</i>	1	Pig (1)
<i>S. muscae</i>	1	Pig (1)
<i>S. sciuri</i>	1	Cat (1)
<i>S. vitulinus</i>	1	Horse (1)
<i>S. fleurettii</i>	1	Chicken (1)
<i>S. lentus</i>	1	Mouse (1)
<i>S. simulans</i>	1	Dog (1)
<i>S. piscifermentans</i>	1	Pig (1)
<i>S. pettenkoferi</i>	1	Cat (1)
<i>S. auricularis</i>	1	Pig (1)
<i>S. equorum</i>	1	Horse (1)
<i>S. xylosus</i>	1	Horse (1)
<i>S. saprophyticus</i>	1	Human (1)
<i>S. nepalensis</i>	1	Mouse (1)
<i>S. kloosii</i>	1	Chicken (1)
<i>S. gallinarum</i>	1	Chicken (1)
<i>S. succinus</i>	1	Horse (1)
<i>S. arlettae</i>	1	Horse (1)
<i>S. cohnii</i>	1	Pig (1)
<i>S. haemolyticus</i>	1	Human (1)
<i>S. pasteurii</i>	1	Common marmoset (1)
<i>S. warneri</i>	1	Human (1)
<i>S. hominis</i>	1	Human (1)
<i>S. lugdunensis</i>	1	Human (1)
<i>S. capitis</i>	1	Human (1)
<i>S. caprae</i>	1	Human (1)
<i>S. simiae</i>	1	Common marmoset (1)
<i>S. epidermidis</i>	1	Human (1)

ment decisions for methicillin-resistant staphylococcal infections (5, 23, 27).

The second reason is the public health issue of whether methicillin-resistant staphylococcal isolates from pet and farm animals are *S. aureus* strains. The isolation of methicillin-resistant *S. aureus* (MRSA) strains is also now increasingly common in veterinary medicine. There are significant concerns about the potential for household pets, horses, and food-producing animals to act as reservoirs of MRSA, with subsequent transmission to humans (21, 31). Therefore, a precise diagnosis of MRSA colonization or infection in animals is a necessary social mission for clinical veterinarians. Our M-PCR method will allow the routine identification of CoPS isolates from var-

TABLE 3. Oligonucleotide primers for M-PCR for species identification of coagulase-positive staphylococci and *S. schleiferi* subsp. *schleiferi*

Primer	Sequence (5'–3')	Size of PCR product (bp)	Species
au-F3	TCGCTTGCTATGATT GTGG	359	<i>S. aureus</i>
au-nucR	GCCAAATGTTCTACCA TAGC		
in-F	CATGTCATATTATTG CGAATGA	430	<i>S. intermedius</i>
in-R3	AGGACCATCACCATT GACATATTGA AACC		
sch-F	AATGGCTACAATGAT AATCACTAA	526	<i>S. schleiferi</i> subsp. <i>coagulans</i>
sch-R	CATATCTGTCTTTTCG GCGCG		<i>S. schleiferi</i> subsp. <i>schleiferi</i>
dea-F	TGAAGGCATATTGTA GAACAA	661	<i>S. delphini</i> group A
dea-R	CGRTACTTTTCGTTA GGTCG		
hy-F1	CATTATATGATTTGA ACGTG	793	<i>S. hyicus</i>
hy-R1	GAATCAATATCGTAA AGTTGC		
pse-F2	TRGGCAGTAGGATT CGTTAA	926	<i>S. pseudintermedius</i>
pse-R5	CTTTTGTGCTCYCMTT TTGG		
deb-F	GGAAGRTTCGTTTTT CCTAGAC	1,135	<i>S. delphini</i> group B
deb-R4	TATGCGATTCAAGAA CTGA		

ious animal species in veterinary clinical laboratories and will provide important clues for approaching the issue.

Ghebremedhin et al. previously reported a comparative analysis of interspecific similarity values of 16S rRNA, *hsp60*, *rpoB*, *sodA*, *tuf*, and *gap* gene sequences in staphylococci (90 to 99%, 74 to 93%, 71.6 to 93.6%, 81.5 to 98%, 86 to 97%, and 24 to 96%, respectively) (14), which indicate the ranges of nucle-




FIG. 1. Phylogenetic tree based on complete thermonuclease (*nuc*) gene sequences in staphylococci. The tree was constructed by the neighbor-joining method using CLUSTAL W.




FIG. 2. Electrophoresis after multiplex PCR for species identification of coagulase-positive staphylococci (CoPS) on a 1.0% agarose gel. Lanes: 1, *S. aureus*; 2, *S. intermedius*; 3, *S. schleiferi*; 4, *S. delphini* group A; 5, *S. hyicus*; 6, *S. pseudintermedius*; 7, *S. delphini* group B; 8, *S. lutrae*; 9, *S. felis*; 10, water (negative control).

otide identity scores. The *nuc* gene has been well conserved and has shown moderate diversity among members of the genus *Staphylococcus*. Therefore, we considered this gene to be a suitable PCR target for species identification.

To date, there has been no reliable method to distinguish among CoPS species in veterinary clinical laboratories. Recently, Bannoehr et al. and Blaiotta et al. reported molecular identification methods for CoPS species by PCR-restriction fragment length polymorphism (PCR-RFLP) targeting the partial *pta* (encoding phosphotransacetylase) and *kataA* (encoding catalase) genes, respectively (3, 7). Compared to these approaches, the present M-PCR method is excellent in terms of rapidity, simplicity, and cost and is better suited for clinical veterinary applications.

We previously reported that phenotypically identified *S. intermedius* strains were reclassified as being *S. intermedius*, *S. pseudintermedius*, and *S. delphini* strains by DNA-DNA hybridization and phylogenetic analysis based on partial *sodA* and *hsp60* gene sequences. In addition, *S. delphini* strains were divided into two clusters (*S. delphini* groups A and B) by *nuc* sequencing analysis (28). Although *S. delphini* group B strains were more closely related to *S. pseudintermedius* strains than to *S. delphini* group A strains (belonging to *S. delphini* LMG 22190^T) upon *nuc* phylogenetic analysis (28), Blaiotta et al. recently reported that *S. delphini* group B strains (strains h-2C and P-27B) were more closely related to *S. intermedius* than to *S. delphini* group A strains according to *kataA* gene analysis (7). *S. delphini* group B strains may therefore represent a unique evolutionary path among staphylococci.

In conclusion, we developed a single-PCR method for the species identification of CoPS, the sensitivity and specificity of which were confirmed by using other molecular-based methods such as *hsp60* and *nuc* sequencing approaches. The present method will contribute to future clinical and research findings for staphylococcal infections in veterinary medicine.

ACKNOWLEDGMENTS

This work was supported by a grant-in-aid for 21st century COE research, by a grant-in-aid for scientific research on priority areas, and

by research fellowships of the Japan Society for the Promotion of Science for Young Scientists from the Ministry of Education, Science, Sports, Culture, and Technology of Japan.

We thank Y. Nakamura, T. Sudo, K. Hayashi, M. Shibutani, and T. Sato for their help in collecting specimens.

REFERENCES

1. Aarestrup, F. M. 2001. Comparative ribotyping of *Staphylococcus intermedius* isolated from members of the Canioidea gives possible evidence for host-specificity and co-evolution of bacteria and hosts. *Int. J. Syst. Evol. Microbiol.* **51**:1343–1347.
2. Abraham, J. L., D. O. Morris, G. C. Griffith, F. S. Shofer, and S. C. Rankin. 2007. Surveillance of healthy cats and cats with inflammatory skin disease for colonization of the skin by methicillin-resistant coagulase-positive staphylococci and *Staphylococcus schleiferi* ssp. *schleiferi*. *Vet. Dermatol.* **18**:252–259.
3. Bannoehr, J., A. Franco, M. Iurescia, A. Battisti, and J. R. Fitzgerald. 2009. Molecular diagnostic identification of *Staphylococcus pseudintermedius*. *J. Clin. Microbiol.* **47**:469–471.
4. Bannoehr, J., N. L. Ben Zakour, A. S. Waller, L. Guardabassi, K. L. Thoday, A. H. M. van den Broek, and J. R. Fitzgerald. 2007. Population genetic structure of the *Staphylococcus intermedius* group: insights into *agr* diversification and the emergence of methicillin-resistant strains. *J. Bacteriol.* **189**:8685–8692.
5. Bemis, D. A., R. D. Jones, L. E. Hiatt, E. D. Ofori, B. W. Rohrbach, L. A. Frank, and S. A. Kania. 2006. Comparison of tests to detect oxacillin resistance in *Staphylococcus intermedius*, *Staphylococcus schleiferi*, and *Staphylococcus aureus* isolates from canine hosts. *J. Clin. Microbiol.* **44**:3374–3376.
6. Bes, M., S. L. Saidi, F. Becharnia, H. Meugnier, F. Vandenesch, J. Etienne, and J. Freney. 2002. Population diversity of *Staphylococcus intermedius* isolates from various host species: typing by 16S-23S intergenic ribosomal DNA spacer polymorphism analysis. *J. Clin. Microbiol.* **40**:2275–2277.
7. Blaiotta, G., V. Fusco, D. Ercolini, O. Pepe, and S. Coppola. 2010. Diversity of *Staphylococcus* species strains based on partial *kat* (catalase) gene sequences and design of a PCR-restriction fragment length polymorphism assay for identification and differentiation of coagulase-positive species (*S. aureus*, *S. delphini*, *S. hyicus*, *S. intermedius*, *S. pseudintermedius*, and *S. schleiferi* subsp. *coagulans*). *J. Clin. Microbiol.* **48**:192–201.
8. Clinical and Laboratory Standards Institute. 2008. Performance standards for antimicrobial susceptibility testing; 18th informational supplement M100-S18. Clinical and Laboratory Standards Institute, Wayne, PA.
9. Descloux, S., A. Rossano, and V. Perreten. 2008. Characterization of new staphylococcal cassette chromosome *meC* (SCC*meC*) and topoisomerase genes in fluoroquinolone- and methicillin-resistant *Staphylococcus pseudintermedius*. *J. Clin. Microbiol.* **46**:1818–1823.
10. Devriese, L. A., M. Vancanneyt, M. Baele, M. Vanechoutte, E. De Graef, C. Snauwaert, I. Cleenwerck, P. Dawyndt, J. Swings, A. Decostere, and F. Haesebrouck. 2005. *Staphylococcus pseudintermedius* sp. nov., a coagulase-positive species from animals. *Int. J. Syst. Evol. Microbiol.* **55**:1569–1573.
11. Fitzgerald, J. R., and J. R. Penades. 2008. Staphylococci of animals, p. 255–269. In J. A. Lindsay (ed.), *Staphylococcus: molecular genetics*. Caister Academic Press, Norfolk, United Kingdom.
12. Freney, J., W. E. Kloos, V. Hajek, and J. A. Webster. 1999. Recommended minimal standards for description of new staphylococcal species. *Int. J. Syst. Bacteriol.* **49**:489–502.
13. Fudaba, Y., K. Nishifuji, L. O. Andresen, T. Yamaguchi, H. Komatsuzawa, M. Amagai, and M. Sugai. 2005. *Staphylococcus hyicus* exfoliative toxins selectively digest porcine desmoglein 1. *Microb. Pathog.* **39**:171–176.
14. Ghebremedhin, B., F. Layer, W. König, and B. König. 2008. Genetic classification and distinguishing of *Staphylococcus* species based on different partial *gap*, 16S rRNA, *hsp60*, *rpoB*, *sodA*, and *tuf* gene sequences. *J. Clin. Microbiol.* **46**:1019–1025.
15. Gill, S. R., D. E. Fouts, G. L. Archer, E. F. Mongodin, R. T. Deboy, J. Ravel, I. T. Paulsen, J. F. Kolonay, L. Brinkac, M. Beanan, R. J. Dodson, S. C. Daugherty, R. Madupu, S. V. Angiuoli, A. S. Durkin, D. H. Haft, J. Vamathevan, H. Khouri, T. Utterback, C. Lee, G. Dimitrov, L. Jiang, H. Qin, J. Weidman, K. Tran, K. Kang, I. R. Hance, K. E. Nelson, and C. M. Fraser. 2005. Insights on evolution of virulence and resistance from the complete genome analysis of an early methicillin-resistant *Staphylococcus aureus* strain and a biofilm-producing methicillin-resistant *Staphylococcus epidermidis* strain. *J. Bacteriol.* **187**:2426–2438.
16. Igimi, S., E. Takahashi, and T. Mitsuoka. 1990. *Staphylococcus schleiferi* subsp. *coagulans* subsp. nov., isolated from the external auditory meatus of dogs with external ear otitis. *Int. J. Syst. Bacteriol.* **40**:409–411.
17. Kuroda, M., A. Yamashita, H. Hiramawa, M. Kumano, K. Morikawa, M. Higashide, A. Maruyama, Y. Inose, K. Matoba, H. Toh, S. Kuhara, M. Hattori, and T. Ohta. 2005. Whole genome sequence of *Staphylococcus saprophyticus* reveals the pathogenesis of uncomplicated urinary tract infection. *Proc. Natl. Acad. Sci. U. S. A.* **102**:13272–13277.
18. Kuroda, M., T. Ohta, I. Uchiyama, T. Baba, H. Yuzawa, I. Kobayashi, L. Cui, A. Oguchi, K. Aoki, Y. Nagai, J. Lian, T. Ito, M. Kanamori, H. Matsumaru, A. Maruyama, H. Murakami, A. Hosoyama, Y. Mizutani-Ui, N. K. Taka-

- hashi, T. Sawano, R. Inoue, C. Kaito, K. Sekimizu, H. Hirakawa, S. Kuhara, S. Goto, J. Yabuzaki, M. Kanehisa, A. Yamashita, K. Oshima, K. Furuya, C. Yoshino, T. Shiba, M. Hattori, N. Ogasawara, H. Hayashi, and K. Hiramatsu. 2001. Whole genome sequencing of methicillin-resistant *Staphylococcus aureus*. *Lancet* **357**:1225–1240.
19. Kwok, A. Y., and A. W. Chow. 2003. Phylogenetic study of *Staphylococcus* and *Micrococcus* species based on partial *hsp60* gene sequences. *Int. J. Syst. Evol. Microbiol.* **53**:87–92.
 20. Meyer, S. A., and K. H. Schleifer. 1978. Deoxyribonucleic acid reassociation in the classification of coagulase-positive staphylococci. *Arch. Microbiol.* **117**:183–188.
 21. Morgan, M. 2008. Methicillin-resistant *Staphylococcus aureus* and animals: zoonosis or humanosis? *J. Antimicrob. Chemother.* **62**:1181–1187.
 22. Morris, D. O., K. A. Rook, F. S. Shofer, and S. C. Rankin. 2006. Screening of *Staphylococcus aureus*, *Staphylococcus intermedius*, and *Staphylococcus schleiferi* isolates obtained from small companion animals for antimicrobial resistance: a retrospective review of 749 isolates (2003–04). *Vet. Dermatol.* **17**:332–337.
 23. Pottumarthy, S., J. M. Schapiro, J. L. Prentice, Y. B. Houze, S. R. Swanzy, F. C. Fang, and B. T. Cookson. 2004. Clinical isolates of *Staphylococcus intermedius* masquerading as methicillin-resistant *Staphylococcus aureus*. *J. Clin. Microbiol.* **42**:5881–5884.
 24. Poyart, C., G. Quesne, C. Boumaila, and P. Trieu-Cuot. 2001. Rapid and accurate species-level identification of coagulase-negative staphylococci by using the *sodA* gene as a target. *J. Clin. Microbiol.* **39**:4296–4301.
 25. Ruscher, C., A. Lübke-Becker, C. G. Wleklinski, A. Soba, L. H. Wieler, and B. Walther. 2009. Prevalence of methicillin-resistant *Staphylococcus pseud-intermedius* isolated from clinical samples of companion animals and equidae. *Vet. Microbiol.* **136**:197–201.
 26. Saitou, N., and M. Nei. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**:846–849.
 27. Sasaki, T., K. Kikuchi, Y. Tanaka, N. Takahashi, S. Kamata, and K. Hiramatsu. 2007. Methicillin-resistant *Staphylococcus pseudintermedius* in a veterinary teaching hospital. *J. Clin. Microbiol.* **45**:1118–1125.
 28. Sasaki, T., K. Kikuchi, Y. Tanaka, N. Takahashi, S. Kamata, and K. Hiramatsu. 2007. Reclassification of phenotypically identified *Staphylococcus intermedius* strains. *J. Clin. Microbiol.* **45**:2770–2778.
 29. Takeuchi, F., S. Watanabe, T. Baba, H. Yuzawa, T. Ito, Y. Morimoto, M. Kuroda, L. Cui, M. Takahashi, A. Ankaï, S. Baba, S. Fukui, J. C. Lee, and K. Hiramatsu. 2005. Whole-genome sequencing of *Staphylococcus haemolyticus* uncovers the extreme plasticity of its genome and the evolution of human-colonizing staphylococcal species. *J. Bacteriol.* **187**:7292–7308.
 30. Thompson, J. D., D. G. Higgins, and T. J. Gibson. 1994. Clustal W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
 31. Weese, J. S., and E. van Duinkerken. 5 February 2009, posting date. Methicillin-resistant *Staphylococcus aureus* and *Staphylococcus pseudintermedius* in veterinary medicine. *Vet. Microbiol.* doi:10.1016/j.vetmic.2009.01.039. [Epub ahead of print.]
 32. Yamashita, K., A. Shimizu, J. Kawano, E. Uchida, A. Haruna, and S. Igimi. 2005. Isolation and characterization of staphylococci from external auditory meatus of dogs with or without otitis externa with special reference to *Staphylococcus schleiferi* subsp. *coagulans* isolates. *Vet. Microbiol.* **107**:139–144.