

Optimization of Hexadecylpyridinium Chloride Decontamination for Culture of *Mycobacterium avium* subsp. *paratuberculosis* from Milk

L. Bradner,^a S. Robbe-Austerman,^b D. C. Beitz,^{a,c} J. R. Stabel^{c,d}

Roy J. Carver Department of Biochemistry, Biophysics, and Molecular Biology, Iowa State University, Ames, Iowa, USA^a; USDA-APHIS, Veterinary Services, Ames, Iowa, USA^b; Department of Animal Science, Iowa State University, Ames, Iowa, USA^c; USDA-ARS, National Animal Disease Center, Ames, Iowa, USA^d

A protocol was optimized for the isolation of *Mycobacterium avium* subsp. *paratuberculosis* (MAP) from milk and colostrum, with parameters including chemical decontamination, antibiotics, and different culture media. This study demonstrates that the efficiency of MAP recovery from milk is highly dependent upon the culturing protocol, and such protocols should be optimized to ensure that low concentrations of MAP in milk can be detected.

In the later stages of infection, *Mycobacterium avium* subsp. *paratuberculosis* (MAP) can disseminate throughout the body and has been isolated from milk, supramammary lymph nodes, and lymph fluid from the udder of cattle (1, 2). Calves that are fed maternal colostrum are more likely to become infected with MAP than those that are fed colostrum replacer or pasteurized colostrum, indicating that raw colostrum and milk may be the initial source of exposure for neonates (3, 4). The culture of MAP from milk presents many difficulties because of the low growth rate of MAP compared with that of other microorganisms present in milk, the fastidious nature of MAP, and the low numbers of MAP (<2 to 8 CFU/50 ml to 10 to 560 CFU/ml) naturally present in milk from infected cows (1, 5). The basic protocol for the isolation of MAP from milk includes centrifugation to concentrate bacterial cells, chemical decontamination (often using hexadecylpyridinium chloride [HPC]) to decrease contaminating microorganisms, and culture in either liquid or solid medium optimized for the recovery of MAP. Protocols for the isolation of MAP from milk vary significantly, and a comprehensive study to identify the most efficacious method for the isolation of MAP from the complex milk matrix is necessary (1, 6–11). Media most commonly used for the culture of MAP from milk include Bactec 12B, *para*-JEM, and Herrold's egg yolk (HEY) (10, 12–14). These media vary greatly in composition, including the presence of growth supplements and antibiotics, as well as growth detection methods, and therefore vary in their ability to support MAP growth (15, 16).

In the present study, triplicate samples of milk obtained from a known Johnne's disease-free cow were spiked with MAP (clinical isolate 167; National Animal Disease Center, Ames, IA) to final concentrations of 10², 10⁴, 10⁶, and 10⁸ CFU/ml. For each experiment, a positive control consisting of 10⁶ CFU of MAP/ml in phosphate-buffered saline (PBS), a negative control consisting of milk with no bacteria, and a no-treatment control consisting of 10⁶ CFU of MAP/ml of milk were included. Within each protocol, the samples (20 ml) were centrifuged at 1,865 × g for 30 min at 4°C to partition the milk unless otherwise specified. The whey layer was discarded, and the cream and pellet layers were retained for the experimental procedure. Effects of temperature (room temperature [RT], 22 to 25°C, and 39°C), centrifugation speed (1,865, 3,500, and 5,600 × g), concentration of HPC (Sigma Chemical Co., St. Louis, MO) decontaminant (0.75, 1.00, 1.25, and 1.50%), time of decontamination (5, 24, and 48 h), and antibiotics (100 µg/ml vancomycin, 50 µg/ml amphotericin B, and

100 µg/ml nalidixic acid; Sigma) on the recovery of viable MAP from milk were investigated. Two liquid culture media, Bactec 12B and *para*-JEM, and one solid medium, HEY, were compared for the ability to recover MAP from spiked milk samples. The Bactec 12B medium (Becton, Dickinson, Franklin, NJ) was supplemented with 0.5 ml of sterile unmodified raw egg yolk (in-house), 0.2 ml PANTA Plus (BD), and 0.1 ml of 50 µg/ml mycobactin J and then inoculated with 200 µl of decontaminated milk sample and placed in a 37°C incubator for 84 days. The *para*-JEM medium (Thermo Fisher Scientific [TREK Diagnostic Systems, Inc.], Cleveland, OH) was supplemented according to the manufacturer's instructions, inoculated with 500 µl of decontaminated milk sample, and placed into the ESP Culture System II machine for up to 65 days. All bottles were subjected to MAP growth confirmation procedures of Ziehl-Neelsen acid fast stain for MAP colonies and PCR of the IS900 gene target as previously described (17). HEY medium (BD) was inoculated with 100 µl of decontaminated milk sample and incubated at 39°C for 12 weeks, and colonies were counted. Statistical analysis was performed with the GLIMMIX procedure of the Statistical Analysis System (SAS Institute, Cary, NC), and a Tukey-Kramer *post hoc* test was performed if results were significant ($P < 0.05$).

The three media tested, Bactec 12B, *para*-JEM, and HEY, differed considerably in the recovery of MAP from decontaminated milk samples. Results demonstrated that HEY medium was inferior to the liquid media in recovery of MAP from milk, with limits of detection estimated to be 10⁴ CFU/ml (data not shown). These findings were comparable to a previous study using HPC decontamination prior to culture on HEY medium (6). Therefore, no further analyses with HEY medium were conducted. Bactec 12B medium consistently recovered MAP at lower concentrations and earlier in incubation ($P < 0.01$) than did the *para*-JEM medium (Fig. 1), even though the volume of inoculum added to bottles was 2.5-fold greater for *para*-JEM than for Bactec 12B. Over all experiments, Bactec 12B medium failed to recover only 3% of the milk

Received 4 February 2013 Accepted 8 February 2013

Published ahead of print 20 February 2013

Address correspondence to J. R. Stabel, judy.stabel@ars.usda.gov.

Copyright © 2013, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JCM.00333-13




FIG 1 Comparison of the Bactec 12B and *para*-JEM media for time to recovery and detection sensitivity of *Mycobacterium avium* subsp. *paratuberculosis* in raw milk spiked with 10² to 10⁸ CFU/ml. Data are expressed as means \pm standard errors of the means (SEM). Significant differences between the two media are represented by asterisks (**, $P < 0.01$).

samples spiked with 10² CFU/ml of MAP, whereas *para*-JEM medium failed to recover 71% of samples at this concentration.

The temperature of decontamination with HPC did not have a significant effect on MAP recovery ($P = 0.63$, data not shown), regardless of culture medium. However, increasing the temperature from RT to 39°C during decontamination of milk samples increased the contamination rates (15.6 to 31.1%) in Bactec 12B and (33.3 to 42.2%) *para*-JEM media. Additionally, increasing centrifugation force from 1,865 to 5,600 $\times g$ did not enhance the recovery of MAP from milk samples ($P = 0.45$, data not shown). Increasing concentration of HPC from 0.75 to 1.50% and time of exposure to HPC from 5 to 48 h resulted in significant reductions ($P < 0.05$) in the recovery of viable MAP, as seen for samples spiked with 10² CFU/ml of MAP and incubated in Bactec 12B medium (Fig. 2). Therefore, the recommended protocol for chemical decontamination of milk to maximize the recovery of MAP in culture was determined to be 0.75% HPC for 5 h at RT.

Overnight incubation of the milk samples with the antibiotic cocktail VAN (consisting of 100 μ g/ml vancomycin [V], 50 μ g/ml amphotericin B [A], and 100 μ g/ml nalidixic acid [N]), following




FIG 2 Time to recovery of *Mycobacterium avium* subsp. *paratuberculosis* in raw milk spiked with 10² CFU/ml after decontamination with 0.75, 1.00, 1.25, and 1.50% hexadecylpyridinium chloride (HPC) for 5, 24, and 48 h in Bactec 12B medium. Data are expressed as means \pm SEM. Significant differences between times of decontamination within an HPC concentration are represented by asterisks (* and **, $P < 0.05$).




FIG 3 Effects on the time to recovery of *Mycobacterium avium* subsp. *paratuberculosis* in milk cultured in Bactec 12B and *para*-JEM media after treatment with the antibiotic cocktail (VAN) consisting of 100 μ g/ml vancomycin (V), 50 μ g/ml amphotericin B (A), and 100 μ g/ml nalidixic acid (N) compared with treatment alone with hexadecylpyridinium chloride (HPC). Data are expressed as means \pm SEM. Significant differences between HPC alone or HPC plus VAN within each medium are represented by asterisks (**, $P < 0.01$; *, $P < 0.05$).

decontamination with 0.75% HPC, resulted in detrimental effects on the recovery of viable MAP (Fig. 3). There was an increase ($P < 0.01$) in the time to recovery after secondary treatment with VAN compared with HPC alone, regardless of liquid medium used or concentration of the MAP inoculum. Treatment with VAN doubled the time to recovery from 33 to 70 days for milk samples spiked with 10² CFU/ml of MAP and incubated in Bactec 12B medium. Exposure to any singular antibiotic (V, A, or N) caused a consistent decrease ($P < 0.05$) in viability in both culture media as well (data not shown).

The three culture media demonstrated different abilities to recover viable MAP from milk. HEY medium proved to be far inferior to the liquid media, with a sensitivity of detection estimated at 10⁴ CFU/ml of MAP. Bactec 12B medium consistently detected inoculation concentrations of 10² CFU/ml, had faster detection times, and had fewer incidences of machine-indicated false-positive and -negative samples compared to *para*-JEM medium. Differences in medium formulation, such as the source of egg yolk and composition of supplemental antibiotics, could have affected the recovery of viable MAP. The manufacturer of Bactec 12B medium recommends the use of PANTA PLUS, which consists of polymyxin B, amphotericin B, nalidixic acid, trimethoprim, and azlocillin. The *para*-JEM medium contains the same antibiotics as the VAN cocktail used in our decontamination protocol. Vancomycin is known to be detrimental to the viability of MAP as demonstrated by a 28- to 49-day increase in the time to detection when vancomycin (86 μ g/ml) was added to the Bactec 12B medium (18). The *para*-JEM medium also contains a reagent called *para*-JEM BLUE which has been shown to retard the growth of MAP by increasing times to detection by up to 1 week (19). The lethality of VAN on the viability of MAP may be compounded, because treatment followed exposure to HPC (6, 20). HPC can damage the MAP bacterium and may cause MAP to become more vulnerable to the antibiotics to which they are normally resistant. Further studies should investigate whether different antibiotics would have similar effects on MAP or whether decreasing the concentration of each antibiotic in the cocktail would decrease the detrimental effects observed for VAN on MAP in this study.

The optimization of a culture method allows the accurate as-

assessment of shedding of MAP into milk of naturally infected cattle, providing critical information to producers about the risk of transmitting MAP to calves via consumption of milk from infected cows.

ACKNOWLEDGMENTS

We thank Philip Dykema, Robin Swanson, and Hannah Wilson for their training, advice, and assistance. We also thank Erica Hellmich and Margaret Walker for their technical assistance in the laboratory.

REFERENCES

- Sweeney R, Whitlock R, Rosenberger A. 1992. *Mycobacterium paratuberculosis* cultured from milk and supramammary lymph nodes of infected asymptomatic cows. *J. Clin. Microbiol.* 30:166–171.
- Khol J, Pinedo P, Buergelt C, Neumann L, Baumgartner W, Rae D. 2012. The collection of lymphatic fluid from the bovine udder and its use for the detection of *Mycobacterium avium* subsp. *paratuberculosis* in the cow. *J. Vet. Diagn. Invest.* 24:23–31.
- Stabel J. 2008. Pasteurization of colostrum reduces the incidence of paratuberculosis in neonatal dairy calves. *J. Dairy Sci.* 91:3600–3606.
- Pithua P, Godden S, Wells S, Oakes M. 2009. Efficacy of feeding plasma-derived commercial colostrum replacer for the prevention of transmission of *Mycobacterium avium* subsp. *paratuberculosis* in Holstein calves. *J. Vet. Med. Assoc.* 234:1167–1176.
- Slana I, Kralik P, Kralova A, Pavlik I. 2008. On-farm spread of *Mycobacterium avium* subsp. *paratuberculosis* in raw milk studied by IS900 and F57 competitive real-time quantitative PCR and culture examination. *Int. J. Food Microbiol.* 128:250–257.
- Gao A, Odumeru J, Raymond M, Mutharia L. 2005. Development of improved method for isolation of *Mycobacterium avium* subsp. *paratuberculosis* from bulk tank milk: effect of age of milk, centrifugation, and decontamination. *Can. J. Vet. Res.* 69:81–87.
- Dundee L, Grant I, Ball H, Rowe M. 2001. Comparative evaluation of four decontamination protocols for the isolation of *Mycobacterium avium* subsp. *paratuberculosis* from milk. *Lett. Appl. Microbiol.* 33:173–177.
- Streeter R, Hoffsis F, Bech-Nielsen S, Shulaw W, Rings D. 1995. Isolation of *Mycobacterium paratuberculosis* from colostrum and milk of subclinically infected cows. *Am. J. Vet. Res.* 56:1322–1324.
- Stabel J, Wells S, Wagner B. 2002. Relationships between fecal culture, ELISA, and bulk tank milk test results for Johne's disease in US dairy herds. *J. Dairy Sci.* 85:525–531.
- Gao A, Odumeru J, Raymond M, Hendrick S, Duffield T, Mutharia L. 2009. Comparison of milk culture, direct and nested polymerase chain reaction (PCR) with fecal culture based on samples from dairy herds infected with *Mycobacterium avium* subsp. *paratuberculosis*. *Can. J. Vet. Res.* 73:58–64.
- Reddacliff L, Vadali A, Whittington R. 2003. The effect of decontamination protocols on the numbers of sheep strain *Mycobacterium avium* subsp. *paratuberculosis* isolated from tissues and faeces. *Vet. Microbiol.* 95:271–282.
- Grant I, Ball H, Rowe M. 2002. Incidence of *Mycobacterium paratuberculosis* in bulk raw and commercially pasteurized cows' milk from approved dairy processing establishments in the United Kingdom. *Appl. Environ. Microbiol.* 68:2428–2435.
- Slana I, Liapi M, Moravkova M, Kralova A, Pavlik I. 2009. *Mycobacterium avium* subsp. *paratuberculosis* in cow bulk tank milk in Cyprus detected by culture and quantitative IS900 and F57 real-time PCR. *Prev. Vet. Med.* 89:223–226.
- Ellingson J, Anderson J, Koziczkowski J, Radcliff R, Sloan S, Allen S, Sullivan N. 2005. Detection of viable *Mycobacterium avium* subsp. *paratuberculosis* in retail pasteurized whole milk by two culture methods and PCR. *J. Food Prot.* 68:966–972.
- Whittington R. 2009. Factors affecting isolation and identification of *Mycobacterium avium* subsp. *paratuberculosis* from fecal and tissue samples in a liquid culture system. *J. Clin. Microbiol.* 47:614–622.
- Tortoli E, Cichero P, Chirillo M, Gismondo M, Bono L, Gesu G, Simonetti M, Volpe G, Nardi G, Marone P. 1998. Multicenter comparison of ESP Culture System II with BACTEC 460TB and with Lowenstein-Jensen medium for recovery of mycobacteria from different clinical specimens, including blood. *J. Clin. Microbiol.* 36:1378–1381.
- Leite F, Stokes K, Robbe-Austerman S, Stabel J. 2013. Comparison of fecal DNA extraction kits for the detection of *Mycobacterium avium* subsp. *paratuberculosis* by polymerase chain reaction. *J. Vet. Diagn. Invest.* 25:27–34.
- Gumber S, Whittington R. 2007. Comparison of BACTEC 460 and MGIT 960 systems for the culture of *Mycobacterium avium* subsp. *paratuberculosis* S strain and observations on the effect of inclusion of ampicillin in culture media to reduce contamination. *Vet. Microbiol.* 119:42–52.
- Okwumabua O, Moua T, Danz T, Quinn J, O'Connor M, Gibbons-Burgener S. 2010. Growth rate retardation and inhibitory effect of para-JEM BLUE of *Mycobacterium avium* subspecies *paratuberculosis*. *J. Vet. Diagn. Invest.* 22:734–737.
- Pribylova R, Kubicova L, Babak V, Pavlik I, Kralik P. 2012. Effect of short- and long-term antibiotic exposure on the viability of *Mycobacterium avium* subsp. *paratuberculosis* as measured by propidium monoazide F57 real-time quantitative PCR and culture. *Vet. J.* 194:354–360.

ERRATUM

Optimization of Hexadecylpyridinium Chloride Decontamination for Culture of *Mycobacterium avium* subsp. *paratuberculosis* from Milk

L. Bradner, S. Robbe-Austerman, D. C. Beitz, J. R. Stabel

Roy J. Carver Department of Biochemistry, Biophysics, and Molecular Biology, Iowa State University, Ames, Iowa, USA; USDA-APHIS, Veterinary Services, Ames, Iowa, USA; Department of Animal Science, Iowa State University, Ames, Iowa, USA; USDA-ARS, National Animal Disease Center, Ames, Iowa, USA

Volume 51, no. 5, p. 1575–1577, 2013. Page 1576: Figure 2 should appear as shown below.

