




The Isolation of *Orientia tsutsugamushi* and *Rickettsia typhi* from Human Blood through Mammalian Cell Culture: a Descriptive Series of 3,227 Samples and Outcomes in the Lao People's Democratic Republic

Damien K. Ming,^{a,b} Vanheuang Phommadeecheak,^b Phonepasith Panyanivong,^b Davanh Sengdatka,^b Weerawat Phuklia,^b Vilada Chansamouth,^{b,d} Tamalee Roberts,^b Stuart D. Blacksell,^{c,d} Paul N. Newton,^{b,c,d}  Matthew T. Robinson^{b,d}

^aFaculty of Medicine, Imperial College, London, United Kingdom

^bLao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit (LOMWRU), Microbiology Laboratory, Mahosot Hospital, Vientiane, Lao People's Democratic Republic

^cMahidol Oxford Tropical Medicine Research Unit (MORU), Bangkok, Thailand

^dCentre for Tropical Medicine & Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom

ABSTRACT In the Lao People's Democratic Republic (Laos), rickettsial infections, including scrub and murine typhus, account for a significant burden of fevers. The Mahosot Hospital Microbiology Laboratory in Vientiane, Laos, routinely performs rickettsial isolation from hospitalized patients with suspected rickettsioses using mammalian cell culture systems. We review the clinical and laboratory factors associated with successful *Orientia tsutsugamushi* and *Rickettsia typhi* isolations from this laboratory over a period of 6 years between 2008 and 2014. The overall isolation success was 7.9% for all samples submitted and 17.3% for samples for which the patient had a positive *O. tsutsugamushi* or *R. typhi* rapid diagnostic test (RDT), serology, or PCR. The frequency of successful isolation was highest for samples submitted in November, at the end of the wet season (28.3%). A longer median duration of reported illness, a positive result for a concurrent *Orientia* or *Rickettsia* spp. quantitative PCR, and the use of antibiotics by the patient in the week before admission were significantly associated with isolation success ($P < 0.05$). Buffy coat inoculation and a shorter interval between sample collection and inoculation in the laboratory were associated with a higher frequency of isolation (both $P < 0.05$). This frequency was highest if cell culture inoculation occurred on the same day as blood sample collection. Factors related to the initial rickettsial bacterial concentration are likely the main contributors to isolation success. However, modifiable factors do contribute to the rickettsial isolation success, especially delays in inoculating patient samples into culture.

KEYWORDS culture, isolation, Laos, *Rickettsia*, *Orientia*, clinical methods

Within the Asia-Pacific region, the rickettsial intracellular bacterial pathogens *Orientia tsutsugamushi* and *Rickettsia typhi* are responsible for a significant burden of treatable febrile illnesses. Scrub typhus, caused by *O. tsutsugamushi*, is the main contributor to this burden and an underrecognized disease affecting predominantly rural areas with poor health infrastructure (1). This disease is transmitted through inoculation by infected chigger mites and is responsible for between 1.8% and 22.3% of fevers among inpatients, depending on region (2, 3), with a median, untreated mortality of approximately 6% (4). Murine typhus, caused by *R. typhi*, and spotted fever group (SFG) rickettsia are also present within the region. Spread through flea and tick species, these diseases tend to be milder in comparison with scrub typhus, though nonetheless severe disease occurs (5). The epidemiology of murine typhus and SFG are less well characterized but caused 11.4% of febrile illnesses in one case series (6).

Citation Ming DK, Phommadeecheak V, Panyanivong P, Sengdatka D, Phuklia W, Chansamouth V, Roberts T, Blacksell SD, Newton PN, Robinson MT. 2020. The isolation of *Orientia tsutsugamushi* and *Rickettsia typhi* from human blood through mammalian cell culture: a descriptive series of 3,227 samples and outcomes in the Lao People's Democratic Republic. *J Clin Microbiol* 58:e01553-20. <https://doi.org/10.1128/JCM.01553-20>.

Editor Bobbi S. Pritt, Mayo Clinic

Copyright © 2020 Ming et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Matthew T. Robinson, matthew.r@tropmedres.ac.

Received 19 June 2020

Returned for modification 30 July 2020

Accepted 23 September 2020

Accepted manuscript posted online 30 September 2020

Published 18 November 2020

Although efficacious treatment exists, clinical diagnosis is difficult due to nonspecific presentations, which are similar to those of other endemic infections, such as malaria and dengue. The presence of an eschar in scrub typhus is variable and diverse presentations, including pneumonitis, septic shock, and meningoenzephalitis, are seen across all rickettsial infections (7). When rickettsial infections are suspected, empirical therapy with an agent such as doxycycline is advocated (8). However, this may pose challenges in pregnancy (9) or for infections involving the central nervous system (10).

Current diagnostic strategies are based on quantitative serological testing using immunofluorescence assays (IFA). However, known limitations in IFA performance, the requirements for convalescent-phase sera, and a lack of standardization can make interpretation of serology difficult (11). Other tests, such as enzyme linked immunosorbent assays (ELISAs) and nucleic acid amplification from clinical specimens, can be helpful (12). The *in vitro* isolation of *Orientia* and *Rickettsia* spp. from clinical samples has been practiced for over 60 years. Propagation of organisms allows antibiotic susceptibility testing, genomic analysis, and study of cell-host interactions that are vital for development of vaccines and diagnostic tests (13, 14). However, as culture techniques are insensitive for diagnosis and require prolonged incubation times, their role in clinically useful diagnosis is currently limited (15).

The Lao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit (LOMWRU), based within the Mahosot Hospital Microbiology Laboratory in Vientiane, Lao People's Democratic Republic (Laos), has been routinely performing rickettsial isolation since 2008. It is currently the only laboratory within Laos equipped for this purpose. Blood samples from febrile patients with suspected rickettsiosis at Mahosot Hospital and regional hospitals in Laos undergo incubation with mammalian cell lines for up to 8 weeks, with the aim of isolating rickettsial organisms. Although there are studies which describe culture conditions used in rickettsial isolation (16), there are none to our knowledge which specifically examine the clinical or laboratory factors associated with successful isolation of *O. tsutsugamushi* and *R. typhi*. We therefore undertook a retrospective review of laboratory data at Mahosot Hospital for samples collected for rickettsial culture over 6 years between January 2008 and December 2014.

(A part of this work was presented at the 28th European Congress of Clinical Microbiology and Infectious Diseases, Madrid, Spain, in 2018 and the 2nd Asia Pacific Rickettsia Conference, Chiang Rai, Thailand, in 2019.)

MATERIALS AND METHODS

All isolation work at LOMWRU was performed in BSL-3 conditions. Up to 5 ml of blood was drawn into EDTA tubes from patients with suspected rickettsioses, all of whom provided written informed consent. Whole EDTA-anticoagulated blood (1 ml) or EDTA buffy coat fraction (200 μ l) mixed with 5 ml of RPMI 1640 medium (PAA Laboratories, USA, before 2014 and Gibco, USA, from 2014 onward) and supplemented with 10% fetal calf serum (PAA Laboratories, USA, before 2014 and concurrently with Gibco, USA, from 2014 onward) was used as an inoculum. The buffy coat layer was prepared by centrifugation of EDTA-whole blood at 1,600 $\times g$ for 10 min. Inoculation was performed directly in 12.5-cm² flasks containing L929 (ATCC number CCL-1) or Vero (ATCC number CCL-81) cell monolayers at 80% confluence, and subsequently centrifuged for 30 min at 50 $\times g$. The flask was incubated for 2 h at 35°C in a 5% CO₂ atmosphere and then the inoculum was washed and replaced with RPMI 1640-10% fetal calf serum medium. Incubation duration was at least eight weeks at 35°C in a 5% CO₂ atmosphere. Media were changed twice weekly, and subculture was performed at week four by resuspending cells and making a 1:2 or 1:10 dilution into 12.5-cm² or 75-cm² flasks containing L929 or Vero cells at 80% confluence. A scraping of the inoculated cell cultures underwent indirect IFA using pooled antibodies against scrub typhus, spotted fever, and typhus group *Rickettsia* every week after four weeks of incubation or earlier (16). Cultures positive by IFA underwent DNA extraction using DNeasy blood and tissue kit (Qiagen, UK) and growth was confirmed by quantitative PCR (qPCR) targeting the 47-kDa protein or 17-kDa outer membrane antigen (for *O. tsutsugamushi* and *Rickettsia* spp., respectively) (17). Statistical analyses of patient information and isolation factors were carried out using Stata 14 (Statacorp, USA). Variables were checked as to whether they were normally distributed by the Shapiro-Wilk or Shapiro-Francia tests before using Wilcoxon signed-rank test for nonnormally distributed variables. Median values and interquartile ranges were calculated. Pearson's chi-squared tests were used to compare frequencies of successful and unsuccessful isolations; exact 95% confidence intervals were calculated for all frequencies. Odds ratios were calculated with 95% confidence intervals and *P* values of ≤ 0.05 were considered significant. Univariate and multivariate logistical regression was used to determine the significance of variables on culture outcome. Backward multivariate analysis was done, except variables with less than 10 samples

TABLE 1 Summary of significant factors in the successful isolation of *O. tsutsugamushi* and *R. typhi*

Parameter ^b	Successful isolation		Unsuccessful isolation		P value
	Frequency (%) or median (n)	95% CI or IQR ^c	Frequency (%) or median (n)	95% CI or IQR ^c	
All patients ^a					
Median age (yrs)	30 (n = 256)	IQR: 18–40.5	26 (n = 2,957)	IQR: 16–39	0.027
Median duration of illness (days)	7 (n = 253)	IQR: 6–1	7 (n = 2,891)	IQR: 4–8	<0.001
Median duration of fever (days)	7 (n = 249)	IQR: 5–10	5 (n = 2,827)	IQR: 3–7	<0.001
History or presence of:					
Rash	43/254 (16.9%)	CI: 12.5–22.1%	273/2862 (9.5%)	CI: 8.5–10.7%	<0.001
Cough	101/240 (42.1%)	CI: 37.4–50.6%	860/2,348 (36.6%)	CI: 34.7–38.6%	0.029
CNS involvement	5/168 (3.0%)	CI: 1.0–6.8%	98/1,207 (8.1%)	CI: 6.6–9.8%	0.018
Sample type					
Buffy coat	58/291 (19.9%)	CI: 15.5–25.0%			<0.001
EDTA whole blood	193/2,907 (6.6%)	CI: 5.7–7.6%			
Median time between collection and inoculation (days)	2 (n = 248)	IQR: 1–3	3 (n = 2,766)	IQR: 1–5	<0.001
STG/TG-positive patients ^a					
Median duration of illness (days)	7 (n = 193)	IQR: 7–10	7 (n = 922)	IQR: 5–9	<0.001
History or presence of CNS involvement	4/158 (5.1%)	CI: 0.7–6.4%	75/831 (9.0%)	CI: 7.2–11.2%	0.006
Sample type					
Buffy coat	58/263 (22.1%)	CI: 17.2–27.6%			0.020
EDTA whole blood	133/841 (15.8%)	CI: 13.6–18.7%			
Median time between collection and inoculation (days)	2 (n = 191)	IQR: 1–4	3 (n = 916)	IQR: 1–7	<0.001

^aAll patients, the full data set (n = 3,227 samples); STG/TG-positive patients, the subset of samples that had either a positive qPCR, positive IgM RDT, and/or a positive IgM or IgG IFA.

^bSTG, scrub typhus group; TG, typhus group; CNS, central nervous system.

^cCI, confidence interval; IQR, interquartile range.

were discounted from analysis and variables that returned an odds ratio of 1 during multivariate analysis were also excluded.

Ethics. Ethical approvals for collection of samples for determination of the etiology of fever were granted by the Lao National Ethics Committee for Health Research and the Oxford Tropical Research Ethics Committee.

RESULTS

A total of 3,227 EDTA-anticoagulated whole blood samples were received for rickettsial isolation from 3,200 patients recruited into fever studies conducted in Vientiane (the capital) (n = 1,064; 33.0%), Luang Namtha (northern Laos) (n = 1,543; 47.8%), and Salavan (southern Laos) (n = 620; 19.2%) between January 2008 and December 2014 (see Table S1 in the supplemental material).

Successful isolation was achieved in 256/3,227 (7.9%) samples. *O. tsutsugamushi* was isolated from 231/256 (90.2%) samples and *R. typhi* from 24/256 (9.4%) samples, and isolations were confirmed through IFA and qPCR. One blood sample yielded both *O. tsutsugamushi* and *R. typhi*, while the species could not be determined for two IFA-positive isolates (0.8%). No spotted fever group rickettsia were cultured. The median duration between incubation start and a positive result was 28 days (interquartile range [IQR]: 23 to 34 days), with no significant difference between organisms cultured (Z = 0.029; P = 0.977) or the cell line used (Z = -0.364; P = 0.716) (Table S2). Cell culture contamination frequency was 373/6,452 (5.8%). The median age of patients was 26 years (IQR, 16 to 39 years; n = 3,213) and was significant between successful and unsuccessful isolations (Z = -2.212; P = 0.027). The majority of patients were male, with 45.2% female (see Table 1 and Table S1).

The isolation data set studied was an amalgamation of a number of routine diagnostic and project-based work. In order to provide a more accurate assessment of isolation success, a subset of samples (n = 1,126; 34.9%) with other laboratory evidence for rickettsial infection were selected for further analysis. These samples came from patients who had a confirmed positive result for scrub typhus group (STG) or typhus group (TG) rickettsiae by one or more of the following diagnostic tests: (i) Dip-S-Ticks murine typhus RDT (PanBio, now ImmunoDOT *Rickettsia typhi*, GenBio, adapted to detect IgM) (43.8%); (ii) scrub typhus IgM RDT (Scrub Typhus IgM and IgG Rapid

Cassette test, PanBio, Australia; Scrub Typhus Total antibody, AccessBio, USA; CareStart Scrub Typhus IgM Rapid Test Device, AccessBio, USA; or SD Bioline Tsutsugamushi IgM Test, Abbott, USA) (50.0%); (iii) *O. tsutsugamushi* 47 kDa qPCR (18) (16.0%); (iv) *Rickettsia* spp. 17 kDa qPCR (17) (4.9%); (v) IgG/IgM STG IFA (IgG, 1.2%; IgM, 1.5%); or (vi) IgG/IgM TG IFA (IgG, 2.8%; IgM, 3.3%). IgG and IgM IFA positivity was based on criteria set by studies in Thailand (19, 20) (see Text S1 in the supplemental material) using both admission and convalescent-phase sera or, where not available, only admission sera. Here, these patients are referred to as STG/TG-positive patients, or either STG-positive or TG-positive when referring to patients positive for one particular group of pathogens. The summary of significant factors associated with isolation success is displayed in Table 1 (the full data are provided in Tables S3 to S5).

The frequency of successful isolation from the STG/TG-positive subset was 17.3% (195/1,126), of which 171 were *O. tsutsugamushi*, 24 were *R. typhi*, and one isolate was unknown. The month with the greatest number of samples submitted was July (138/1,111, 12.4%), with the fewest in February (43/1,111, 3.9%). However, the frequency of successful isolations was highest from samples submitted in November (26/92, 28.3%) and lowest during February (1/43, 2.3%) ($P < 0.001$) (Fig. 1A). Isolation of *O. tsutsugamushi* from STG-positive patients was significantly associated with month ($P < 0.001$), with the highest isolation rate seen in December (41.5%) (Fig. 1B), while November had the highest isolation rate for *R. typhi* from TG-positive patients (11.6%) but with no significant relationship ($P = 0.804$) (Fig. 1C). From all STG/TG-positive patients, successful isolations were significantly associated with a longer duration of illness ($P < 0.001$), a reduced incidence of CNS involvement (i.e., symptoms of confusion, seizures, neck stiffness, or altered consciousness; $P = 0.006$), a history of taking an antibiotic during the previous week ($P = 0.022$), and sample type being buffy coat ($P = 0.020$) (Table 2). In addition to the associations described above, successful isolation of *O. tsutsugamushi* from STG-positive patients was also associated with a history or presence of a rash ($P = 0.037$) (Table 2 and Table S4). For TG-positive patients, successful isolation of *R. typhi* was only associated with a history or presence of a cough ($P = 0.011$) or a greater use of antibiotics in the preceding week ($P < 0.001$) (Table 2 and Table S5). The median whole blood volume submitted for isolation was 5.0 ml, but there was no significant association with isolation success ($P = 0.760$) (Table S4). In the 1,006 patients with data, the antibiotic classes prescribed or purchased in the week prior to sampling included beta-lactams (47.3%), tetracyclines (23.8%), quinolones (7.6%), and macrolides (4.4%), with 19.2% of patients taking combination therapy.

Running PCRs (i.e., *O. tsutsugamushi* 47 kDa qPCR or *Rickettsia* spp. 17 kDa qPCR) on a patient's EDTA buffy coat sample as a way to select samples for culturing was significantly associated with a higher rate of isolation success for the respective pathogens (65.8% and 38.9%, respectively; $P < 0.001$ for either PCR), although no significance was seen in relation to actual threshold cycle (C_T) values and isolation success ($P = 0.324$ and $P = 0.709$ for *O. tsutsugamushi* and *Rickettsia* spp., respectively). Similarly, samples from patients with positive IgM IFAs were more likely to have successful isolation for both pathogens (STG = 18.5% success, $P = 0.026$; TG = 60.0% success, $P < 0.001$), while a positive IgG IFA was only significant for *R. typhi* isolation (88.9% success, $P < 0.001$) (Table 3 and Table S6).

The majority of blood samples (72.6%) were received by the laboratory for inoculation into cell lines within 5 days. Successful isolation was associated with shorter median interval between blood collection and cell inoculation of 2 days (IQR: 1 to 4) versus 3 days (IQR: 1 to 7) ($n = 1,107$; $Z = 5.729$, $P < 0.001$) for all STG/TG-positive patients, and 2 days (IQR: 1 to 4) versus 4 days (IQR: 2 to 8) ($n = 635$; $Z = 6.922$, $P < 0.001$) for all STG-positive patients. For all samples from STG/TG-positive patients, the frequency of successful isolation was highest if the blood sample was inoculated on the same day as collection (29.7%, 19/64) and decreased with samples inoculated on subsequent days ($n = 938$; $P = 0.032$) (Fig. 2). Although not significant, there were also differences between the nine different operators who carried out the isolations, with their overall rates of isolation success ranging from 5.6% to 33.3% ($P = 0.809$).




FIG 1 Total number of blood samples submitted for rickettsial isolation (bars) and percentage of those with successful isolations (line) by month between 2008 and 2014. (A) All STG/TG-positive patients ($n = 1,111$). (B) *O. tsutsugamushi* (Ot) isolations from STG-positive patients ($n = 636$). (C) *R. typhi* (Rt) isolations from TG-positive patients ($n = 510$).

A multivariate logistic regression model for isolation of either organism from STG/TG-positive patients identified that a positive PCR was significantly associated with isolation success, although the confidence interval was large (odds ratio [OR], 49.3; 95% CI 9.79 to 248.20; $P < 0.001$). Isolation of *O. tsutsugamushi* from STG-positive patients was significantly associated with a positive *O. tsutsugamushi* PCR (OR, 9.64; 95% CI 5.42 to 17.16; $P < 0.001$), while the presence of a cough was significant for *R. typhi* isolation from TG-positive patients (OR, 3.17; 95% CI 1.08 to 9.35; $P = 0.036$) (see Tables S7 to S10).

DISCUSSION

The *in vitro* isolation of rickettsial organisms from clinical samples is increasingly important in the context of reports of reduced antibiotic susceptibility (21) and the

TABLE 2 Summary of significant factors in the successful isolation of *O. tsutsugamushi* or *R. typhi* from samples from STG- or TG-positive patients

Parameter ^b	Successful isolation ^a		Unsuccessful isolation		P value
	Frequency (%) or median (n)	95% CI or IQR ^c	Frequency (%) or median (n)	95% CI or IQR	
STG-positive patients					
Median duration of illness (days)	7 (n = 163)	IQR: 7–11	7 (n = 486)	IQR: 5–10	0.002
History or presence of rash	26/161 (16.1%)	CI: 10.8–22.8%	48/477 (10.1%)	CI: 7.5–13.1%	0.037
History or presence of CNS involvement	3/129 (2.3%)	CI: 0.5–6.6%	49/421 (11.6%)	CI: 8.7–15.1%	0.002
Sample type					
Buffy coat	45/89 (50.6%)	CI: 39.8–61.3%			<0.001
EDTA whole blood	115/554 (20.8%)	CI: 17.3–24.3%			
Median time between collection and inoculation (days)	2 (n = 159)	IQR: 1–4	4 (n = 476)	IQR: 2–8	<0.001
TG-positive patients					
History or presence of cough	14/21 (66.6%)	CI: 43.0–85.4%	187/480 (39.0%)	CI: 34.6–43.5%	0.011
Took an antibiotic in the previous wk	10/16 (62.5%)	CI: 35.4–84.8%	151/397 (38.1%)	CI: 33.2–43.0%	<0.001

^aSuccessful isolation of *O. tsutsugamushi* from STG-positive patients and *R. typhi* from TG-positive patients.

^bSTG, scrub typhus group; TG, typhus group; CNS, central nervous system.

^cCI, confidence interval; IQR, interquartile range.

need for understanding the genomic diversity of rickettsial pathogens for development of vaccines and diagnostics.

O. tsutsugamushi was isolated more frequently than *R. typhi*. Although murine typhus is present in Vientiane (6), it is less common than scrub typhus and therefore this is likely the reason for fewer isolations. In addition, optimal culture conditions may vary between *Rickettsia* spp. (22) and *Orientia tsutsugamushi* (23) and the cell culture conditions used here may not have been optimal for *R. typhi*. The optimization of *R. typhi* isolation should be investigated in greater detail.

The initial rickettsial bacterial load within the blood sample submitted is the likely main predictor of isolation outcome. Overall, C_T values were high (mean = 38.14; IQR = 36.03 to 39.29; $n = 69$), indicating low bacterial loads. Although comparisons between C_T values of successful and unsuccessful isolations do not show any statistical significance, there is a significant relationship between overall qPCR positivity and isolation success in the regression model, as well as a higher isolation frequency through the use of the buffy coat, suggesting an association with bacterial load. The dynamics of rickettsial infection are complex. An early, limited bacteremic phase for *O. tsutsugamushi* before intracellular entry has been described from animal models (24), consistent with a relatively low median number of organisms (<100 copies/ μ l) within human blood (25, 26). Within our cohort, a longer duration of illness was associated with a successful isolation, probably because prolonged illness is associated with a higher bacterial burden. The severity of illness has been associated with higher bacterial burdens in *O. tsutsugamushi* and *R. typhi* infections (27) and potentially why we see that the presence of a cough (for *R. typhi*) or a rash (for *O. tsutsugamushi*) is significantly associated with successful isolation. We also found a significant association between isolation and antibiotic use in the preceding week before admission, although, unexpectedly, this was related to an increase in successful isolations. One possible explanation for this is that patients with more severe disease had a higher chance of seeking

TABLE 3 Diagnostic tests that were significantly associated with a successful isolation of *O. tsutsugamushi* or *R. typhi* when positive for the target organism

Category	Test(s) ^a (P value)
Associated with successful isolation for the whole population	qPCR (<0.001)
Associated with successful isolation of <i>O. tsutsugamushi</i>	<i>O. tsutsugamushi</i> qPCR (<0.001), STG IgM RDT (<0.001), STG IgM IFA (0.026)
Associated with successful isolation of <i>R. typhi</i>	<i>Rickettsia</i> spp. qPCR (<0.001), TG IgM RDT (0.001), TG IgM IFA (<0.001), TG IgG IFA (<0.001)

^aSTG, scrub typhus group; TG, typhus group; IFA, immunofluorescence assay; RDT, rapid diagnostic test; qPCR, quantitative PCR.




FIG 2 Days between sample collection and inoculation in cell culture versus isolation success. (A) Percentage successful isolation for all STG/TG-positive patients ($R^2=0.673$; $P = 0.032$). (B) Percentage successful isolation of *O. tsutsugamushi* from STG-positive patients ($R^2=0.640$; $P = 0.252$). (C) Percentage successful isolation of *R. typhi* from TG-positive patients ($R^2=0.272$, $P = 0.222$).

antibiotics, but the majority (74%) of those antibiotics taken would not be efficacious against either rickettsial pathogen (28, 29).

The use of PCR to screen for possible positive patients showed a significant association with isolation success when positive. This may be expected, as PCRs rely on the detection of bacterial DNA in the sample, indicative of a current infection. IFAs were not used to select samples for isolation in any of the associated studies here and were completed after samples had been sent for isolation. As would be expected, positive IgM IFAs are significantly associated with a successful isolation, as IgM is indicative of a current or recent infection. Although positive RDTs are also significantly associated with successful isolations (see Table S6 in the supplemental material), the majority of attempted isolations were carried out based on positive RDT results; therefore, it is

difficult to interpret their usefulness without discounting the risk of bias in the data. This information highlights the benefit of using diagnostic tests to guide the selection of samples for isolation.

Temporal changes in isolation success for *O. tsutsugamushi* reflect the observed seasonality of scrub typhus, which appears after onset of the annual monsoon season (6, 8). Isolation success frequency was highest for blood samples collected at the end of the wet season (28.3% for all rickettsial pathogens) and lowest for those submitted in the middle of the dry season (2.3%). The seasonal increase in isolation rate during the wet season may result through increased caseload from chigger exposure in activities such as rice harvest, as well as ecological factors which tend to increase vector density and alter feeding behavior during that period (30). A similar pattern, although on a much smaller scale, is seen with *R. typhi* isolation. Although not statistically significant, seasonal changes are most likely due to the same factors because seasonality in regional flea abundance has been previously identified (31).

A shorter time between blood collection and inoculation was associated with higher rates of isolation and may relate to decline in viability of the bacteria in *ex vivo* blood with time. Delays occur because samples collected from study sites have to be transported to the laboratory in Vientiane, and rickettsial isolation was not performed during weekends. The impact of delayed inoculation has been demonstrated for *R. conorii*, for which the frequency of isolation through shell-vial culture was reduced when clinical samples were not inoculated on the same day as blood draw (32). Systems aimed to shorten the time to inoculation or maintain viability through use of hemoculture fluid (33) may improve isolation success; this is important as the initial bacterial concentration in clinical samples is likely to be low. We also found variation in success rates between different technicians performing the isolation, most likely due to an individual's level of training and experience. Further evaluation or competency training, along with routine audit, would be essential for standardizing practices. The findings within the multivariate model that PCR was the most significant factor at predicting isolation success suggest that current culture success depends largely on initial patient and bacterial factors. Laboratory data, in respect to isolations, were complete and 3,145/3,227 (97.5%) patients had at least some clinical information available, although this was highly variable. Prehospital antibiotic use data were only available for 1,767/3,200 (55.2%) patients and qPCR data available for 1,610/3,227 samples (49.9%).

In summary, we describe the clinical and laboratory factors associated with *O. tsutsugamushi* and *R. typhi* isolation success in Laos. Similar to the culture of conventional bacteria from blood, we found that host factors such as illness duration and factors indicative of initial blood bacterial concentration are strongly associated with isolation success. It will be important to further define the role of culture technique to aid staff training and reduce delayed sample inoculation to improve the isolation yield and support research in this group of diseases.

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

SUPPLEMENTAL FILE 1, PDF file, 0.05 MB.

SUPPLEMENTAL FILE 2, PDF file, 0.1 MB.

ACKNOWLEDGMENTS

We are grateful to the director of Mahosot Hospital, the director and staff of the Microbiology Laboratory, and the ward doctors and nurses at Mahosot Hospital, especially the Microbiology Laboratory. We thank the Adult Infectious Diseases Ward, Sethathirat Hospital, the Luang Namtha and Salavan Hospitals, and colleagues in MORU-Bangkok who made this work possible. We thank Catrin Moore and Sabine Dittrich for their past supervision of rickettsial culture. We are also grateful to Phoudthasone Kouaykesone for her help with data collection.

The work was supported by the Wellcome Trust. S.D.B., P.N.N., and M.T.R. are funded by the Wellcome Trust.

REFERENCES

- Bonell A, Lubell Y, Newton PN, Crump JA, Paris DH. 2017. Estimating the burden of scrub typhus: a systematic review. *PLoS Negl Trop Dis* 11: e0005838. <https://doi.org/10.1371/journal.pntd.0005838>.
- Kasper MR, Blair PJ, Touch S, Sokhal B, Yasuda CY, Williams M, Richards AL, Burgess TH, Wierzbza TF, Putnam SD. 2012. Infectious etiologies of acute febrile illness among patients seeking health care in south-central Cambodia. *Am J Trop Med Hyg* 86:246–253. <https://doi.org/10.4269/ajtmh.2012.11-0409>.
- Blacksell SD, Sharma NP, Phumratanaprapin W, Jenjaroen K, Peacock SJ, White NJ, Pukrittayakamee S, Day NP. 2007. Serological and blood culture investigations of Nepalese fever patients. *Trans R Soc Trop Med Hyg* 101:686–690. <https://doi.org/10.1016/j.trstmh.2007.02.015>.
- Taylor AJ, Paris DH, Newton PN. 2015. A systematic review of mortality from untreated scrub typhus (*Orientia tsutsugamushi*). *PLoS Negl Trop Dis* 9:e0003971. <https://doi.org/10.1371/journal.pntd.0003971>.
- Dumler JS, Taylor JP, Walker DH. 1991. Clinical and laboratory features of murine typhus in south Texas, 1980 through 1987. *JAMA* 266:1365–1370. <https://doi.org/10.1001/jama.1991.03470100057033>.
- Phongmany S, Rolain J-M, Phetsouvanh R, Blacksell SD, Soukhaseum V, Rasachack B, Phiasakha K, Soukhaseum S, Frichithavong K, Chu V, Keouangkhout V, Martinez-Aussel B, Chang K, Darasavath C, Rattanavong O, Sisouphone S, Mayxay M, Vidamaly S, Parola P, Thammanong C, Heuangvongsy M, Syhavong B, Raoult D, White NJ, Newton PN. 2006. Rickettsial infections and fever, Vientiane, Laos. *Emerg Infect Dis* 12:256–262. <https://doi.org/10.3201/eid1202.050900>.
- Xu G, Walker DH, Jupiter D, Melby PC, Arcari CM. 2017. A review of the global epidemiology of scrub typhus. *PLoS Negl Trop Dis* 11:e0006062. <https://doi.org/10.1371/journal.pntd.0006062>.
- Mayxay M, Castonguay-Vanier J, Chansamouth V, Dubot-Peres A, Paris DH, Phetsouvanh R, Tangkhabuanbutra J, Douangdala P, Inthath S, Souvannasing P, Slesak G, Tongyoo N, Chanthongthip A, Panyanouvong P, Sibounheuang B, Phommason K, Dohnt M, Phonekeo D, Hongvanthong B, Xayadeth S, Ketmayoon P, Blacksell SD, Moore CE, Craig SB, Burns MA, von Sonnenburg F, Corwin A, de Lamballerie X, Gonzalez JJ, Christophel EM, Cawthorne A, Bell D, Newton PN. 2013. Causes of non-malarial fever in Laos: a prospective study. *Lancet Glob Health* 1:e46–e54. [https://doi.org/10.1016/S2214-109X\(13\)70008-1](https://doi.org/10.1016/S2214-109X(13)70008-1).
- Cross R, Ling C, Day NP, McGready R, Paris DH. 2016. Revisiting doxycycline in pregnancy and early childhood—time to rebuild its reputation? *Expert Opin Drug Saf* 15:367–382. <https://doi.org/10.1517/14740338.2016.1133584>.
- Dittrich S, Rattanavong S, Lee SJ, Panyanivong P, Craig SB, Tulsiani SM, Blacksell SD, Dance DAB, Dubot-Pères A, Sengduangphachanh A, Phoumin P, Paris DH, Newton PN. 2015. *Orientia*, rickettsia, and leptospira pathogens as causes of CNS infections in Laos: a prospective study. *Lancet Glob Health* 3:e104–e112. [https://doi.org/10.1016/S2214-109X\(14\)70289-X](https://doi.org/10.1016/S2214-109X(14)70289-X).
- Lim C, Paris DH, Blacksell SD, Laongnualpanich A, Kantipong P, Chierakul W, Wuthiekanun V, Day NP, Cooper BS, Limmathurotsakul D. 2015. How to determine the accuracy of an alternative diagnostic test when it is actually better than the reference tests: a re-evaluation of diagnostic tests for scrub typhus using Bayesian LCMs. *PLoS One* 10:e0114930. <https://doi.org/10.1371/journal.pone.0114930>.
- Dhawan S, Robinson MT, Stenos J, Graves SR, Wangrangsimakul T, Newton PN, Day NPJ, Blacksell SD. 2020. Selection of diagnostic cutoffs for murine typhus IgM and IgG immunofluorescence assay: a systematic review. *Am J Trop Med Hyg* 103:55–63. <https://doi.org/10.4269/ajtmh.19-0818>.
- Watt G, Chouriyagune C, Ruangwearayud R, Watcharapichat P, Phulsuksombati D, Jongsakul K, Teja-Isavadharm P, Bhodhidatta D, Corcoran KD, Dasch GA, Strickman D. 1996. Scrub typhus infections poorly responsive to antibiotics in northern Thailand. *Lancet* 348:86–89. [https://doi.org/10.1016/S0140-6736\(96\)02501-9](https://doi.org/10.1016/S0140-6736(96)02501-9).
- Valbuena G, Walker DH. 2012. Approaches to vaccines against *Orientia tsutsugamushi*. *Front Cell Infect Microbiol* 2:170. <https://doi.org/10.3389/fcimb.2012.00170>.
- Gouriet F, Fenollar F, Patrice JY, Drancourt M, Raoult D. 2005. Use of shell-vial cell culture assay for isolation of bacteria from clinical specimens: 13 years of experience. *J Clin Microbiol* 43:4993–5002. <https://doi.org/10.1128/JCM.43.10.4993-5002.2005>.
- Luksameetanasan R, Blacksell S, Kalambaheti T, Wuthiekanun V, Chierakul W, Chueasuwanchai S, Apiwattanaporn A, Stenos J, Graves S, Peacock S, Day N. 2007. Patient and sample-related factors that effect the success of in vitro isolation of *Orientia tsutsugamushi*. *Southeast Asian J Trop Med Public Health* 38:91–96.
- Jiang J, Chan T-C, Temenak JJ, Dasch GA, Ching W-M, Richards AL. 2004. Development of a quantitative real-time polymerase chain reaction assay specific for *Orientia tsutsugamushi*. *Am J Trop Med Hyg* 70: 351–356. <https://doi.org/10.4269/ajtmh.2004.70.351>.
- Horinouchi H, Murai K, Okayama A, Nagatomo Y, Tachibana N, Tsubouchi H. 1996. Genotypic identification of *Rickettsia tsutsugamushi* by restriction fragment length polymorphism analysis of DNA amplified by the polymerase chain reaction. *Am J Trop Med Hyg* 54:647–651. <https://doi.org/10.4269/ajtmh.1996.54.647>.
- Lim C, Blacksell SD, Laongnualpanich A, Kantipong P, Day NP, Paris DH, Limmathurotsakul D. 2015. Optimal cutoff titers for indirect immunofluorescence assay for diagnosis of scrub typhus. *J Clin Microbiol* 53: 3663–3666. <https://doi.org/10.1128/JCM.01680-15>.
- Phakhounthong K, Mukaka M, Dittrich S, Tanganuchitcharnchai A, Day NPJ, White LJ, Newton PN, Blacksell SD. 2020. The temporal dynamics of humoral immunity to *Rickettsia typhi* infection in murine typhus patients. *Clin Microbiol Infect* 26:781.e9–781.e16. <https://doi.org/10.1016/j.cmi.2019.10.022>.
- Tantibhedhyangkul W, Angelakis E, Tongyoo N, Newton PN, Moore CE, Phetsouvanh R, Raoult D, Rolain JM. 2010. Intrinsic fluoroquinolone resistance in *Orientia tsutsugamushi*. *Int J Antimicrob Agents* 35: 338–341. <https://doi.org/10.1016/j.ijantimicag.2009.11.019>.
- Ives TJ, Marston EL, Regnery RL, Butts JD. 2001. In vitro susceptibilities of *Bartonella* and *Rickettsia* spp. to fluoroquinolone antibiotics as determined by immunofluorescent antibody analysis of infected Vero cell monolayers. *Int J Antimicrob Agents* 18:217–222. [https://doi.org/10.1016/S0924-8579\(01\)00388-0](https://doi.org/10.1016/S0924-8579(01)00388-0).
- Tamura A, Makisaka Y, Enatsu T, Urakami H, Okubo K, Fukuhara M, Mahara F. 1999. Isolation of *Orientia tsutsugamushi* from patients in Shikoku and finding of a strain which grows preferentially at low temperatures. *Microbiol Immunol* 43:979–981. <https://doi.org/10.1111/j.1348-0421.1999.tb03359.x>.
- Sunyakumthorn P, Paris DH, Chan TC, Jones M, Luce-Fedrow A, Chattopadhyay S, Jiang J, Anantatat T, Turner GD, Day NP, Richards AL. 2013. An intradermal inoculation model of scrub typhus in Swiss CD-1 mice demonstrates more rapid dissemination of virulent strains of *Orientia tsutsugamushi*. *PLoS One* 8:e54570. <https://doi.org/10.1371/journal.pone.0054570>.
- Sonthayanon P, Chierakul W, Wuthiekanun V, Phimda K, Pukrittayakamee S, Day NP, Peacock SJ. 2009. Association of high *Orientia tsutsugamushi* DNA loads with disease of greater severity in adults with scrub typhus. *J Clin Microbiol* 47:430–434. <https://doi.org/10.1128/JCM.01927-08>.
- Paris DH, Aukkanit N, Jenjaroen K, Blacksell SD, Day NP. 2009. A highly sensitive quantitative real-time PCR assay based on the groEL gene of contemporary Thai strains of *Orientia tsutsugamushi*. *Clin Microbiol Infect* 15:488–495. <https://doi.org/10.1111/j.1469-0691.2008.02671.x>.
- Dittrich S, Castonguay-Vanier J, Moore CE, Thongyoo N, Newton PN, Paris DH. 2014. Loop-mediated isothermal amplification for *Rickettsia typhi* (the causal agent of murine typhus): problems with diagnosis at the limit of detection. *J Clin Microbiol* 52:832–838. <https://doi.org/10.1128/JCM.02786-13>.
- Kim YS, Yun HJ, Shim SK, Koo SH, Kim SY, Kim S. 2004. A comparative trial of a single dose of azithromycin versus doxycycline for the treatment of mild scrub typhus. *Clin Infect Dis* 39:1329–1335. <https://doi.org/10.1086/425008>.
- Brown GW, Saunders JP, Singh S, Huxsol DL, Shirai A. 1978. Single dose doxycycline therapy for scrub typhus. *Trans R Soc Trop Med Hyg* 72: 412–416. [https://doi.org/10.1016/0035-9203\(78\)90138-4](https://doi.org/10.1016/0035-9203(78)90138-4).
- Kuo CC, Lee PL, Chen CH, Wang HC. 2015. Surveillance of potential hosts

- and vectors of scrub typhus in Taiwan. *Parasit Vectors* 8:611. <https://doi.org/10.1186/s13071-015-1221-7>.
31. Kim H-C, Yang Y-C, Chong S-T, Ko S-J, Lee S-E, Klein TA, Chae J-S. 2010. Detection of *Rickettsia typhi* and seasonal prevalence of fleas collected from small mammals in the Republic of Korea. *J Wildl Dis* 46:165–172. <https://doi.org/10.7589/0090-3558-46.1.165>.
32. La Scola B, Raoult D. 1996. Diagnosis of Mediterranean spotted fever by cultivation of *Rickettsia conorii* from blood and skin samples using the centrifugation-shell vial technique and by detection of *R. conorii* in circulating endothelial cells: a 6-year follow-up. *J Clin Microbiol* 34: 2722–2727. <https://doi.org/10.1128/JCM.34.11.2722-2727.1996>.
33. Dittrich S, Card E, Phuklia W, Rudgard WE, Silousok J, Phoumin P, Bouthasavong L, Azarian S, Davong V, Dance DA, Vongsouvath M, Phetsouvanh R, Newton PN. 2016. Survival and growth of *Orientia tsutsugamushi* in conventional hemocultures. *Emerg Infect Dis* 22: 1460–1463. <https://doi.org/10.3201/eid2208.151259>.